

# SOME CHARACTERIZATIONS OF 0-DISTRIBUTIVE SEMILATTICES

H.S. CHAKRABORTY AND M.R. TALUKDER

ABSTRACT. In this paper we discuss prime down-sets of a semilattice. We give a characterization of prime down-sets of a semilattice. We also give some characterizations of 0-distributive semilattices and a characterization of minimal prime ideals containing an ideal of a 0-distributive semilattice. Finally, we give a characterization of minimal prime ideals of a pseudocomplemented semilattice.

## 1. INTRODUCTION

Semilattices have been studied by many authors. The class of distributive semilattices is an important subclass of semilattices. We refer the readers to [4, 9, 10] for distributive semilattices. We also refer the monograph [5] for the background of distributive semilattices. The class of 0-distributive semilattices is a nice extension of the class of distributive semilattices. This extension is useful for the study of pseudocomplemented semilattices. For pseudocomplemented semilattices we refer the readers to [2, 3, 5, 6]. We also refer the readers to [7, 8] for 0-distributive semilattices (see [1, 11] for 0-distributive lattices). In this paper we study 0-distributive semilattices. By semilattice we mean meet-semilattice.

A semilattice  $\mathbf{S}$  with 0 is called **0-distributive** if for any  $a, b, c \in S$  such that  $a \wedge b = 0 = a \wedge c$  implies  $a \wedge d = 0$  for some  $d \geq b, c$ . The **pentagonal lattice**  $\mathcal{P}_5$  (see Figure 1) as a semilattice is 0-distributive but the **diamond lattice**  $\mathcal{M}_3$  (see Figure 1) as a semilattice is not 0-distributive. A semilattice  $\mathbf{S}$  is called **directed above** if for all  $x, y \in S$  there exists  $z \in S$  such that  $z \geq x, y$ . Every 0-distributive semilattice is directed above.

---

*Date:* September 9, 2012.

*2010 Mathematics Subject Classification.* 06A12, 06A99, 06B10.

*Key words and phrases.* Semilattices, distributive semilattice, 0-distributive semilattice, pseudocomplemented semilattice, ideal, filter.




FIGURE 1. 0-distributive and non-0-distributive

Minimal prime ideals and maximal filters play an important role in semilattices. In Section 2, we introduce a notion of minimal prime down-set and maximal filters in semilattices. Here we give a characterization of minimal prime down-sets and maximal filters in semilattices.

Like as a distributive semilattice (or distributive lattice) Stone's version separation theorem is not true for 0-distributive semilattice. For example, if we consider the pentagonal lattice  $\mathcal{P}_5$  (see Figure 1) as a 0-distributive semilattice, then  $F = [c]$  is a filter and  $I = [a]$ , is an ideal such that  $F \cap I = \emptyset$  but there is no prime filter containing  $F$  and disjoint from  $I$ . In Section 3 we discuss Stone's version separation theorem for 0-distributive semilattices. In this section we give some characterizations of 0-distributive semilattices.

In Section 4 we discuss the pseudocomplementation in semilattices. We close the paper with a characterization of a minimal prime ideals of a pseudocomplemented 0-distributive semilattice.

## 2. PRIME DOWN-SETS AND MAXIMAL FILTERS

Let  $\mathbf{S}$  be a semilattice. A non-empty subset  $D$  of  $S$  is called a **down-set** if  $a \in D, b \in S$  with  $b \leq a$  implies that  $b \in D$ . A down-set  $D$  of  $\mathbf{S}$  is called a **proper down-set** if  $D \neq S$ . A **prime down-set** is a proper down-set  $P$  of  $\mathbf{S}$  such that  $a \wedge b \in P$  implies  $a \in P$  or  $b \in P$ . A prime down-set  $P$  is called **minimal** if there is a prime down-set  $Q$  such that  $Q \subseteq P$ , then  $P = Q$ .

**Theorem 2.1.** *Any prime down-set of a semilattice contains a minimal prime down-set.*

*Proof.* Let  $\mathbf{S}$  be a semilattice with 0. Let  $P$  be a prime down-set of  $\mathbf{S}$  and let  $\mathcal{P}$  be the set of all prime down-sets contained in  $P$ . Then  $\mathcal{P}$

is non-empty since  $P \in \mathcal{P}$ . Let  $\mathcal{C}$  be a chain in  $\mathcal{P}$  and let

$$M := \bigcap \{X \mid X \in \mathcal{C}\}.$$

We claim that  $M$  is a prime down-set. Clearly  $M$  is non-empty as  $0 \in M$ . Let  $a \in M$  and  $b \leq a$ . Then  $a \in X$  for all  $X \in \mathcal{C}$ . Hence  $b \in X$  for all  $X \in \mathcal{C}$  as  $X$  is a down-set. Thus  $b \in M$ . Now let  $x \wedge y \in M$  for some  $x, y \in S$ . Then  $x \wedge y \in X$  for all  $X \in \mathcal{C}$ . Since  $X$  is a prime down-set for all  $X \in \mathcal{C}$ , we have either  $x \in X$  or  $y \in X$  for all  $X \in \mathcal{C}$ . This implies that either  $x \in M$  or  $y \in M$ . Hence  $M$  is a prime down-set.

Thus by applying the dual form of Zorn's Lemma to  $\mathcal{P}$ , there is a minimal member of  $\mathcal{P}$ .  $\square$

Let  $\mathbf{S}$  be a semilattice. A non-empty subset  $F$  of  $S$  is called a **filter** if

- (i)  $a, b \in F$  implies  $a \wedge b \in F$
- (ii)  $a \in S, b \in F$  with  $a \geq b$  implies  $a \in F$ .

A filter  $F$  of a semilattice  $\mathbf{S}$  is called **proper filter** if  $F \neq S$ . A **maximal filter**  $F$  of  $\mathbf{S}$  is a proper filter which is not contained in any other proper filter, that is, if there is a proper filter  $G$  such that  $F \subseteq G$ , then  $F = G$ .

Following result is due to [8].

**Lemma 2.2.** *Let  $M$  be a proper filter of  $\mathbf{S}$  with  $0$ . Then  $M$  is maximal if and only if for all  $a \in S \setminus M$ , there is some  $b \in M$  such that  $a \wedge b = 0$ .*

Now we have the following result.

**Theorem 2.3.** *Let  $F$  be a non-empty proper subset of a semilattice  $\mathbf{S}$ . Then  $F$  is a filter if and only if  $S \setminus F$  is a prime down-set.*

*Proof.* Let  $F$  be a filter of a semilattice  $\mathbf{S}$ . Let  $x \in S \setminus F$  and  $y \leq x$ . Then  $x \notin F$  and hence  $y \notin F$  as  $F$  is a filter. This implies  $y \in S \setminus F$ . Thus  $S \setminus F$  is a down-set. Since  $F$  is a filter  $S \setminus F \neq S$ . Thus  $S \setminus F$  is a proper down-set. To prove  $S \setminus F$  is a prime down-set, let  $a, b \in S$  such that  $a \wedge b \in S \setminus F$ . Then  $a \wedge b \notin F$  and hence either  $a \notin F$  or  $b \notin F$  as  $F$  is filter. This implies either  $a \in S \setminus F$  or  $b \in S \setminus F$ . Therefore,  $S \setminus F$  is a prime down-set.

Conversely, let  $S \setminus F$  be a prime down-set and  $x, y \in F$ . Then clearly,  $x, y \notin S \setminus F$  and hence  $x \wedge y \notin S \setminus F$  as  $S \setminus F$  is a prime down-set. Thus  $x \wedge y \in F$ . Suppose  $x \in F$  and  $x \leq y$ . Then  $x \notin S \setminus F$ . Since

$S \setminus F$  is a down-set, we have  $y \notin S \setminus F$ . Hence  $y \in F$ . This implies  $F$  is a filter.  $\square$

**Theorem 2.4.** *Let  $F$  be a non-empty subset of a semilattice  $\mathbf{S}$ . Then  $F$  is a maximal filter if and only if  $S \setminus F$  is a minimal prime down-set.*

*Proof.* Let  $F$  be a maximal filter and  $S \setminus F$  is not a minimal prime down-set. Then there exists a prime down-set  $I$  such that  $I \subseteq S \setminus F$  which implies  $F \subseteq S \setminus I$  which contradict to the maximality of  $F$ . Hence  $S \setminus F$  is minimal prime down-set.

Conversely, let  $S \setminus F$  be a minimal prime down-set and  $F$  is not a maximal filter. Thus there exists a proper filter  $G$  such that  $F \subseteq G$  which implies  $S \setminus G \subseteq S \setminus F$  which contradict the minimality of  $S \setminus F$ . Hence  $F$  is a maximal filter.  $\square$

### 3. MINIMAL PRIME IDEALS

Let  $\mathbf{S}$  be a semilattice. A down-set  $I$  of  $S$  is called an **ideal** if  $a, b \in I$  implies the existence of  $c \in I$  such that  $a, b \leq c$ . The set of all ideals of  $S$  is denoted by  $\mathcal{I}(S)$ . An ideal  $I$  of  $\mathbf{S}$  is called a **proper ideal** if  $I \neq S$ . A **prime ideal**  $P$  is a proper ideal of  $\mathbf{S}$  such that  $a \wedge b \in P$  implies either  $a \in P$  or  $b \in P$ . A prime ideal  $P$  is called **minimal** if there is a prime ideal  $Q$  such that  $Q \subseteq P$ , then  $P = Q$ . A filter  $F$  of  $\mathbf{S}$  is called a **prime filter** if  $F \neq S$  and  $S \setminus F$  is a prime ideal.

We shall often use the following lemma in this paper.

**Lemma 3.1.** *Let  $\mathbf{S}$  be a directed above semilattice with 0. If  $\mathbf{S}$  is not 0-distributive, then the set*

$$F := \{x \in S \mid x \geq a \wedge y \neq 0 \text{ for all } y \geq b, c\},$$

where  $a, b, c \in S$  such that  $a \wedge b = a \wedge c = 0$ , is a proper filter.

*Proof.* Since  $\mathbf{S}$  is not 0-distributive, there are  $p, q, r \in S$  such that  $p \wedge q = p \wedge r = 0$  and  $p \wedge d \neq 0$  for all  $d \geq q, r$ . Now we have  $p \geq p \wedge d$ . Thus  $p \in F$ . Hence  $F$  is nonempty. Clearly  $0 \notin F$ . It is enough to show that  $F$  is a filter. Let  $x \in F$  and  $z \geq x$ . Then  $x \geq a \wedge y$  for all  $y \geq b, c$  and by transitivity  $z \geq a \wedge y$  for all  $y \geq b, c$ . Hence  $z \in F$ . Again let  $x, z \in F$ . Then  $x \geq a \wedge y$  and  $z \geq a \wedge y$  for all  $y \geq b, c$ . Thus  $x \wedge z \geq a \wedge y$  for all  $y \geq b, c$ . Hence  $x \wedge z \in F$ . This implies  $F$  is a filter.  $\square$

Now we have the following result.

**Theorem 3.2.** *Every maximal filter of a 0-distributive semilattice is a prime filter.*

*Proof.* Let  $\mathbf{S}$  be a 0-distributive semilattice. Again let  $Q$  be a maximal filter of  $S$ . We shall show that  $Q$  is prime. It is sufficient to show that  $S \setminus Q$  is a prime ideal. By Theorem 2.4 we have  $S \setminus Q$  is a minimal prime down-set. Now let  $x, y \in S \setminus Q$ . Then by Lemma 2.2 we have  $a \wedge x = 0 = b \wedge y$  for some  $a, b \in Q$ . Let  $c = a \wedge b$ . Clearly  $c \wedge x = 0 = c \wedge y$  and  $c \in Q$ . Hence by the 0-distributivity of  $\mathbf{S}$  there exists  $z \in S$  such that  $z \geq x, y$  and  $c \wedge z = 0$ . Hence  $z \in S \setminus Q$ . Thus  $S \setminus Q$  is a prime ideal which implies  $Q$  is prime.  $\square$

Let  $A$  be non-empty subset of a semilattice  $\mathbf{S}$  with 0. Set

$$A^\perp := \{x \in S \mid a \wedge x = 0 \text{ for all } a \in A\}.$$

Then  $A^\perp$  is called the **annihilator** of  $A$ . If  $A = S$  then  $A^\perp = S^\perp = (0)$ . For  $a \in S$ , the annihilator of  $\{a\}$  is simply denoted by  $a^\perp$  and hence  $a^\perp = \{x \in S \mid a \wedge x = 0\}$ . We can easily show that

$$A^\perp = \bigcap_{a \in A} a^\perp.$$

Let  $\mathbf{S}$  be a semilattice with 0. An ideal  $I$  of  $\mathbf{S}$  is called an **annihilator ideal** if  $I = A^\perp$  for some non-empty subset  $A$  of  $S$ .

Our aim is to prove a Stone's version separation theorem for 0-distributive semilattices. The following result due to [8, Theorem 7].

**Theorem 3.3.** *Let  $\mathbf{S}$  be a semilattice with 0. Then  $\mathbf{S}$  is 0-distributive if and only if for any filter  $F$  of  $S$  such that  $F \cap x^\perp = \emptyset$  ( $x \in S$ ), there exists a prime filter containing  $F$  and disjoint from  $x^\perp$ .*

Our conjecture is:

**Conjecture 3.4.** *Let  $\mathbf{S}$  be a directed above semilattice with 0. Then  $\mathbf{S}$  is 0-distributive if and only if for any filter  $F$  and any annihilator ideal  $I$  of  $S$  such that  $F \cap I = \emptyset$ , there exists a prime filter containing  $F$  and disjoint from  $I$ .*

The necessary conditions of a directed above semilattice to be 0-distributive is given below, but unfortunately, we could not prove or disprove the condition is sufficient or not.

**Theorem 3.5.** *Let  $\mathbf{S}$  be a directed above semilattice with 0. If for any filter  $F$  and any annihilator ideal  $I$  of  $S$  such that  $F \cap I = \emptyset$ , there exists a prime filter containing  $F$  and disjoint from  $I$ , then  $\mathbf{S}$  is 0-distributive.*

*Proof.* Suppose the condition holds. If  $\mathbf{S}$  is not 0-distributive, then there are  $a, b, c \in S$  such that  $a \wedge b = 0 = a \wedge c$  and  $a \wedge d \neq 0$  for all  $d \geq b, c$  (such  $d$  exists as  $\mathbf{S}$  is directed above). Let

$$F := \{x \in S \mid x \geq a \wedge y \text{ for all } y \geq b, c\}.$$

Then by Lemma 3.1, we have  $F$  is a proper filter.

Let  $I$  be an annihilator ideal such that  $a \wedge d \notin I$  (such annihilator exists as  $a \wedge d \notin S^\perp$ ). We shall show that  $I \cap F = \emptyset$ . If  $x \in I \cap F$ , then  $x \geq a \wedge y$  for all  $y \geq b, c$  which implies  $a \wedge d \in I$ , which is a contradiction. Hence  $I \cap F = \emptyset$ . Thus by the assumption, there is a prime filter  $Q$  such that  $F \subseteq Q$  and  $I \cap Q = \emptyset$ . This implies  $a \in Q$  and  $y \in Q$  for all  $y \geq b, c$ . We shall show that either  $b \in Q$  or  $c \in Q$ . If  $b, c \notin Q$  then  $b, c \in S \setminus Q$ . Since  $Q$  is a prime filter,  $S \setminus Q$  is a prime ideal. So, there is  $e \in S \setminus Q$  such that  $e \geq b, c$  which is a contradiction. Hence either  $b \in Q$  or  $c \in Q$ . This implies, either  $a \wedge b \in Q$  or  $a \wedge c \in Q$ . Hence  $0 \in Q$ , which contradicts the fact that  $Q$  is a prime filter. Therefore,  $a \wedge d = 0$  for some  $d \geq b, c$  and hence  $\mathbf{S}$  is 0-distributive.  $\square$

Let  $\mathbf{S}$  be a semilattice. For  $a \in S$ , the ideal  $(a]$  is called the **ideal generated by  $a$** . It can be easily seen that  $(a]^\perp = a^\perp$  for any  $a \in S$ . An ideal  $I$  of  $S$  is called an  $\alpha$ -**ideal** if  $(i^\perp)^\perp \subseteq I$  for any  $i \in I$ .

Now we shall give some characterizations of 0-distributive semilattice. The following lemma is due to [1]

**Lemma 3.6.** *Every proper filter of a semilattice with 0 is contained in a maximal filter.*

We have the following result which is a generalization of [1, Theorem 3.1].

**Theorem 3.7.** *Let  $\mathbf{S}$  be a semilattice with 0. Then the following statements (i)–(iv) are equivalent and any one of them implies (v) and (vi).*

- (i)  $\mathbf{S}$  is 0-distributive;
- (ii) every maximal filter of  $S$  is prime;
- (iii) every minimal prime down-set of  $S$  is a minimal prime ideal;

- (iv) every proper filter of  $S$  is disjoint from a minimal prime ideal;
- (v) for each element  $a \in S$  such that  $a \neq 0$ , there is a minimal prime ideal not containing  $a$ ;
- (vi) each element  $a \in S$  such that  $a \neq 0$  is contained in a prime filter.

Proof. (i) $\Rightarrow$ (ii). This follows by the Lemma 3.2.

(ii) $\Rightarrow$ (iii). Let  $N$  be a minimal prime down-set. Then by Lemma 2.4 we have  $S \setminus N$  is a maximal filter. Hence by (ii)  $S \setminus N$  is a prime filter. Thus  $N$  is a prime ideal.

(iii) $\Rightarrow$ (iv). Let  $F$  be a proper filter of  $S$ . By Lemma 3.6 there is a maximal filter  $M$  such that  $F \subseteq M$ . Hence by Lemma 2.4 we have  $S \setminus M$  is a minimal prime down-set. Thus by (iii)  $S \setminus M$  is a minimal prime ideal. Clearly,  $F \cap (S \setminus M) = \emptyset$ .

(iv) $\Rightarrow$ (i). Suppose  $\mathbf{S}$  is not 0-distributive. Then there are  $a, b, c \in S$  such that  $a \wedge b = a \wedge c = 0$  and  $a \wedge d \neq 0$  for all  $d \geq b, c$ . Now set

$$F = \{x \in S \mid x \geq a \wedge y \text{ for all } y \geq b, c\}.$$

Then by Lemma 3.1, we have  $F$  is a proper filter and hence by (iv) there exists a prime ideal  $Q$  such that  $F \cap Q = \emptyset$ . Thus  $a \wedge p \notin Q$  for any  $p \geq b, c$ . This implies  $a, p \notin Q$  for any  $p \geq b, c$ . Now  $a \notin Q$  implies  $b, c \in Q$ . Then there is  $m \geq b, c$  such that  $m \in Q$  which is a contradiction. Therefore,  $a \wedge d = 0$  for some  $d \geq b, c$  and hence  $\mathbf{S}$  is 0-distributive.

(iv) $\Rightarrow$ (v). Let  $a \in S$  such that  $a \neq 0$ . Then  $[a]$  is a proper filter. Then by (iv)  $[a]$  is disjoint from a minimal prime ideal  $N$  of  $S$ . Thus  $a \notin N$ .

(v) $\Rightarrow$ (vi). Let  $a \in S$  such that  $a \neq 0$ . Then by (v) there is a minimal prime ideal  $P$  such that  $a \notin P$  which implies  $a \in S \setminus P$ . By the definition of prime filter we have  $S \setminus P$  is a prime filter.  $\square$

Now we have following result which is a generalization of [1, Lemma 1.8].

**Lemma 3.8.** *Let  $A$  be a non-empty subset of a semilattice  $\mathbf{S}$  with 0. Then  $A^\perp$  is the intersection of all the minimal prime down-set not containing  $A$ .*

Proof. Let  $\mathbf{S}$  be a semilattice with 0 and  $\emptyset \neq A \subseteq S$ . Suppose

$$X := \bigcap \{P \mid A \not\subseteq P \text{ and } P \text{ is a minimal prime down-set}\}$$

Let  $x \in A^\perp$ . Then  $x \wedge y = 0$  for all  $y \in A$ . This implies there is  $z \notin P$  such that  $x \wedge z = 0 \in P$ . As  $P$  is prime, we have  $x \in P$ . Hence  $x \in X$ .

Conversely let,  $x \in X$ . If  $x \notin A^\perp$ . Then  $x \wedge q \neq 0$  for some  $q \in A$ . Let  $D = [x \wedge q]$ . Then  $0 \notin D$ . Hence,  $D \neq S$ . Then by Lemma 3.6 we have  $D \subseteq M$  for some maximal filter  $M$ . Hence by Lemma 2.4 we have  $S \setminus M$  is a minimal prime down-set. Now  $x \notin S \setminus M$  as  $x \in D$  implies  $x \in M$ . Moreover  $A \not\subseteq S \setminus M$  as  $q \in A$  but  $q \in M$  implies  $q \notin S \setminus M$ , which is a contradiction to  $x \in X$ . Hence  $x \in A^\perp$ . Thus the lemma is proved.  $\square$

**Theorem 3.9.** *Let  $\mathbf{S}$  be a 0-distributive semilattice. If  $A$  is a nonempty subset of  $S$  and  $F$  is a proper filter intersecting  $A$ , there is a minimal prime ideal containing  $A^\perp$  and disjoint from  $F$ .*

*Proof.* Let  $\mathbf{S}$  be a directed above semilattice with 0. Again let  $A$  be a nonempty subset of  $\mathbf{S}$  and  $F$  be a proper filter such that  $F \cap A \neq \emptyset$ . Then Lemma 2.3  $S \setminus F$  is a prime down-set and by Lemma 2.1  $N \subseteq S \setminus F$  for some minimal prime down-set  $N$ . Clearly,  $N \cap F = \emptyset$ . Also  $A \not\subseteq S \setminus F$  and so  $A \not\subseteq N$ . By Lemma 3.8  $A^\perp \subseteq N$ . Since  $\mathbf{S}$  is 0-distributive, by theorem 3.7(iv)  $N$  is a minimal prime ideal.  $\square$

#### 4. PSEUDOCOMPLEMENTATION FOR 0-DISTRIBUTIVE SEMILATTICES

Let  $\mathbf{S}$  be a semilattice with 0. An element  $d \in S$  is called the **pseudocomplement** of  $x \in S$ , if  $x \wedge d = 0$  and  $y \in S, x \wedge y = 0$  implies  $y \leq d$ . The pseudocomplement of  $x$  is denoted by  $x^*$ . A semilattice  $\mathbf{S}$  is called pseudocomplemented if each element of  $S$  has a pseudocomplement. The pseudocomplement of 0 is the largest element 1. Thus a pseudocomplemented semilattice contains both the smallest element and the largest element.

**Theorem 4.1.** *Every pseudocomplemented semilattice is 0-distributive but the converse is not true.*

*Proof.* Let  $\mathbf{S}$  be a pseudocomplemented semilattice. Suppose  $a, b, c \in S$  with  $a \wedge b = 0 = a \wedge c$ . By the definition of pseudocomplemented,  $b \leq a^*, c \leq a^*$  and  $a \wedge a^* = 0$ . Thus  $\mathbf{S}$  is a 0-distributive semilattice.



To prove the converse is not true, consider the semilattice,  $\mathcal{M}_2$  shown in the Figure 2, which is clearly 0-distributive but not pseudocomplemented as  $a^*$  does not exist.




FIGURE 2

□

**Theorem 4.2.** *Let  $S$  be a pseudocomplemented semilattice and let  $J$  be an ideal of  $S$ . Then a prime ideal  $P$  containing  $J$  is a minimal prime ideal containing  $J$  if and only if for each  $x \in P$  there is  $y \in S \setminus P$  such that  $x \wedge y \in J$ .*

*Proof.* Let  $P$  be a prime ideal of  $S$  containing  $J$  such that the given condition holds. We shall show that  $P$  is a minimal prime ideal containing  $J$ . Let  $K$  be a prime ideal containing  $J$  such that  $K \subseteq P$ . Let  $x \in P$ . Then there is  $y \in S \setminus P$  such that  $x \wedge y \in J$ . Hence  $x \wedge y \in K$  as  $K$  containing  $J$ . Since  $K$  is prime and  $y \notin K$  implies  $x \in K$ . Hence  $P \subseteq K$ . Thus  $K = P$ . Therefore,  $P$  is a minimal prime ideal containing  $J$ .

Conversely, let  $P$  be a minimal prime ideal containing  $J$ . Let  $x \in P$ . Suppose for all  $y \in S \setminus P$ ,  $x \wedge y \notin J$ . Set  $D = (S \setminus P) \vee [x]$ . We claim that  $0 \notin D$ . For if  $0 \in D$ , then  $0 = q \wedge x$  for some  $q \in S \setminus P$ . Thus,  $x \wedge q = 0 \in J$  which is a contradiction. Therefore,  $0 \notin D$ . Since  $(0] = 1^\perp$  by Theorem 3.3, there is a prime filter  $Q$  such that  $D \subseteq Q$  and  $0 \notin Q$ . Let  $M = S \setminus Q$ . Then by the definition of prime filter of a semilattice,  $M$  is a prime ideal. We claim that  $M \cap D = \emptyset$ . If  $a \in M \cap D$ , then  $a \in M$  and hence  $a \notin Q$ . Thus  $a \notin D$  which is a contradiction. Hence  $M \cap D = \emptyset$ . Therefore,  $M \cap (S \setminus P) = \emptyset$  and hence  $M \subseteq P$ . Also  $M \neq P$ , because  $x \in D$  implies  $x \in Q$  and hence  $x \notin M$ .

but  $x \in P$ . This shows that  $P$  is not minimal which is a contradiction. Hence the given condition holds.  $\square$

We enclose the paper with the following useful characterization of minimal prime ideal.

**Theorem 4.3.** *Let  $\mathbf{S}$  be a pseudocomplemented semilattice and let  $P$  be a prime ideal of  $S$ . Then the followings are equivalent:*

- (i)  $P$  is minimal.
- (ii)  $x \in P$  implies that  $x^* \notin P$ .

Proof. (i) $\Rightarrow$ (ii). Let  $P$  be a minimal prime ideal and let  $x^* \in P$  for some  $x \in P$ . Set  $D = (S \setminus P) \vee [x]$ . We claim that  $0 \notin D$ . For if  $0 \in D$ , then  $0 = q \wedge x$  for some  $q \in S \setminus P$ , which implies  $q \leq x^* \in P$  which is a contradiction. Therefore,  $0 \notin D$ . Since  $(0] = 1^\perp$  by Theorem 3.3, there is a prime filter  $Q$  such that  $D \subseteq Q$  and  $0 \notin Q$ . Let  $M = S \setminus Q$ . Then by the definition of prime filter of a semilattice,  $M$  is a prime ideal. We claim that  $M \cap D = \emptyset$ . If  $a \in M \cap D$ , then  $a \in M$  and hence  $a \notin Q$ . Thus  $a \notin D$  which is a contradiction. Hence  $M \cap D = \emptyset$ . Therefore,  $M \cap (S \setminus P) = \emptyset$  and hence  $M \subseteq P$ . Also  $M \neq P$ , because  $x \in D$  implies  $x \in Q$  and hence  $x \notin M$  but  $x \in P$ . This shows that  $P$  is not minimal which is a contradiction. Hence (ii) holds.

(ii) $\Rightarrow$ (i). Let  $P$  be a prime ideal of  $S$  such that (ii) holds. We shall show that  $P$  is a minimal prime ideal. Let  $K$  be a prime ideal satisfying (ii) such that  $K \subseteq P$ . Let  $x \in P$ . Then  $x \wedge x^* = 0 \in K$ . Since  $K$  is prime and  $x^* \notin K$  implies  $x \in K$ . Hence  $P \subseteq K$ . Thus  $K = P$ . Therefore,  $P$  is a minimal prime ideal.  $\square$

## REFERENCES

- [1] P. Balasubramani and P. V. Venkatanarasimhan, *Characterizations of the 0-Distributive Lattice*, Indian J. pure appl. Math. **32**(3) 315-324, March (2001).
- [2] S.N. Begum and A.S.A. Noor, *Congruence kernels of distributive JP-semilattices*, Mathematica Bohemica, in press.
- [3] T.S. Blyth, *Ideals and Filters of Pseudocomplemented Semilattices*, *Proceedings of the Edinburgh Mathematical Society*, **23** (1980), 301–316.
- [4] W.H. Cornish, *Characterization of distributive and modular semilattices*, Math. Japonica, **22** (1977), 159–174.
- [5] G. Grätzer, *General Lattice Theory*, Birkhäuser Verlag Basel, 1998.
- [6] G. Grätzer, *Lattice Theory First Concept and Distributive Lattice*, Sanfrancisco W. H. Freeman, (1971).

- [7] V. V. Joshi and B. N. Waphare, *Characterizations of 0-distributive posets*, *Mathematica Bohemica* Vol. **01** (2005), 73-80.
- [8] Y. S. Pawar and N. K. Thakare, *0-Distributive semilattices*, *Canad. Math. Bull.* Vol. **21(4)** (1978), 469-475.
- [9] P. V. Ramana Murty and M. Krishna Murty, *Some remarks on certain classes of semilattices*, *Internat. J. Math. & Math. Sci.*, Vol. **05** No. **01**, (1982), 21-30.
- [10] J.B. Rhodes, *Modular and distributive semilattice*. *Trans. Amer. Math. Society.* Vol. 201 (1975), 31-41.
- [11] J.C. Varlet, *A Generalization of the Notion of Pseudo-complementedness*. *Bulletin de ln Société des Sciences de Liège*, Vol. **36** (1968), 149-158

DEPARTMENT OF MATHEMATICS, BANGABANDHU SHEIKH MUJIBUR RAHMAN  
SCIENCE AND TECHNOLOGY UNIVERSITY, GOPALGANJ-8100, BANGLADESH  
*E-mail address:* apu.012@yahoo.com

DEPARTMENT OF MATHEMATICS, SHAHJALAL UNIVERSITY OF SCIENCE AND  
TECHNOLOGY, SYLHET-3114, BANGLADESH  
*E-mail address:* r.talukder-mat@sust.edu