

Ore extensions over right strongly Hopfian rings

Ouyang Lunqun and Liu Jinwang

Department of Mathematics, Hunan University of Science and Technology

Xiangtan, Hunan 411201, P.R. China

Xiang Yueming

Department of Mathematics and Applied Mathematics,

Huaihua University, Huaihua, 418000, P.R. China

Abstract An associative ring is said to be right strongly Hopfian if the chain of right annihilators $r_R(a) \subseteq r_R(a^2) \subseteq \dots$ stabilizes for each $a \in R$. In this article, we are interested in the class of right strongly Hopfian rings and the transfer of this property from an associative ring R to the Ore extension $R[x; \alpha, \delta]$ and the monoid ring $R[M]$. It is proved that if R is (α, δ) -compatible and $R[x; \alpha, \delta]$ is reversible, then the Ore extension $R[x; \alpha, \delta]$ is right strongly Hopfian if and only if R is right strongly Hopfian, and it is also showed that if M is a strictly totally ordered monoid and $R[M]$ is a reversible ring, then the monoid ring $R[M]$ is right strongly Hopfian if and only if R is right strongly Hopfian. Consequently, several known results regarding strongly Hopfian rings are extended to a more generally setting.

Keywords strongly Hopfian ring; Ore extension; monoid ring.

2000 MR Subject Classification 16S99.

1. Introduction

Throughout this paper all rings are associative with identity. For a nonempty subset X of a ring R , $l_R(X) = \{a \in R \mid aX = 0\}$ and $r_R(X) = \{a \in R \mid Xa = 0\}$ denote the left and the right annihilator of X in R , respectively. Following A. Hmaimou et al [5], a ring R is left strongly Hopfian if for every endomorphism f of R , the chain $\ker f \subseteq \ker f^2 \subseteq \dots$ stabilizes. Equivalently, R is left strongly Hopfian if the chain of left annihilators $l_R(a) \subseteq l_R(a^2) \subseteq \dots$ stabilizes for each $a \in R$. The class of left strongly Hopfian rings is very large. It contains Noetherian rings, Laskerian rings, rings

satisfying *acc* on d-annihilators and those satisfying *acc* on d-colons, and so on [4]. If R is a commutative ring, then a left strongly Hopfian ring is also called a strongly Hopfian ring. A. Hmaimou et al [5] showed that for a commutative ring R , the ring R is strongly Hopfian if and only if the polynomial ring $R[x]$ is strongly Hopfian if and only if the Laurent polynomial ring $R[x; x^{-1}]$ is strongly Hopfian. Let R be a commutative ring. In [4], S. Hizem provided an example of a strongly Hopfian ring R such that the power series ring $R[[x]]$ is not necessary strongly Hopfian, and also gave some necessary and sufficient conditions for $R[[x]]$ to be strongly Hopfian. For more details and properties of left strongly Hopfian rings, see [2, 4, 5, 7, 8].

Let α be an endomorphism, and δ an α -derivation of R , that is, δ is an additive map such that $\delta(ab) = \delta(a)b + \alpha(a)\delta(b)$, for $a, b \in R$. According to Annin [1], a ring R is said to be α -compatible if for each $a, b \in R, ab = 0 \Leftrightarrow a\alpha(b) = 0$. Clearly, this may only happen when the endomorphism α is injective. Moreover, R is said to be δ -compatible if for each $a, b \in R, ab = 0 \Rightarrow a\delta(b) = 0$. A ring R is (α, δ) -compatible if it is both α -compatible and δ -compatible. Recall that a ring R is reversible if $ab = 0 \Rightarrow ba = 0$ for all $a, b \in R$, and a ring R is semicommutative if $ab = 0$ implies $aRb = 0$ for any $a, b \in R$. Clearly, any subring of a reversible ring is also reversible, and if R is a reversible ring, then for any $n \in \mathbb{N}$ and any permutation $\sigma \in S_n, x_1x_2 \cdots x_n = 0$ implies $x_{\sigma(1)}Rx_{\sigma(2)}R \cdots Rx_{\sigma(n)}R = 0$ for any $x_i \in R, 1 \leq i \leq n$. Reversible rings are semicommutative, but the reverse is not true in general [6, Example 1.5]. Let $f(x) = a_0 + a_1x + \cdots + a_nx^n \in R[x; \alpha, \delta]$, $\{a_0, a_1, \dots, a_n\}$ denotes the subset of R comprised of the coefficients of $f(x)$.

In this article, we are interested in the class of right strongly Hopfian rings and the transfer of this property from an associative ring R to the Ore extension $R[x; \alpha, \delta]$ and the monoid ring $R[M]$. We first provide some examples of right strongly Hopfian rings. We next show that: (1) if R is (α, δ) -compatible and $R[x; \alpha, \delta]$ is reversible, then the Ore extension $R[x; \alpha, \delta]$ is right strongly Hopfian if and only if R is right strongly Hopfian. (2) If M is a strictly totally ordered monoid and $R[M]$ a reversible ring, then the monoid ring $R[M]$ is right strongly Hopfian if and only if R is right strongly Hopfian.

2. Extensions of right strongly Hopfian rings

Definition 2.1 *A ring R is right strongly Hopfian if the chain of right annihilators $r_R(a) \subseteq r_R(a^2) \subseteq \cdots$ stabilizes for each $a \in R$.*

The next Lemma is known and very useful. We leave the proof for the reader.

Lemma 2.2 *Let $a \in R$. Then the chain $r_R(a) \subseteq r_R(a^2) \subseteq \cdots$ stabilizes if and only if there exists $n > m$ such that $r_R(a^n) = r_R(a^m)$.*

Lemma 2.3 *Let $A \subset B$ be an extension of rings. If B is right strongly Hopfian, then so is A .*

Proof Let $a \in A$. Then $r_A(a) = r_B(a) \cap A$.

Proposition 2.4 *Let $T_n(R)$ denote the $n \times n$ upper triangular matrix ring over a ring R . Then the following conditions are equivalent:*

- (1) R is right strongly Hopfian;
- (2) $T_n(R)$ is right strongly Hopfian.

Proof (1) \Rightarrow (2). Suppose R is right strongly Hopfian and let

$$A = \begin{pmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ 0 & a_{22} & \cdots & a_{2n} \\ \cdots & \cdots & \cdots & \cdots \\ 0 & 0 & \cdots & a_{nn} \end{pmatrix} \in T_n(R).$$

We proceed by induction on n to show that $T_n(R)$ is right strongly Hopfian. Let $n = 2$. Put $\alpha = \begin{pmatrix} a & b \\ 0 & c \end{pmatrix} \in T_2(R)$. Since R is right strongly Hopfian, there exists $m \in \mathbb{N}$ such that for any $n > m$, $r_R(a^n) = r_R(a^m)$ and $r_R(c^n) = r_R(c^m)$. Now we show that $r_{T_2(R)}(\alpha^{2m+1}) = r_{T_2(R)}(\alpha^{2m})$. If $\beta = \begin{pmatrix} x & y \\ 0 & z \end{pmatrix} \in r_{T_2(R)}(\alpha^{2m+1})$, then

$$\begin{aligned} & \alpha^{2m+1}\beta \\ &= \begin{pmatrix} a^{2m+1} & a^{2m}b + a^{2m-1}bc + \cdots + a^m b c^m + \cdots + b c^{2m} \\ 0 & c^{2m+1} \end{pmatrix} \begin{pmatrix} x & y \\ 0 & z \end{pmatrix} \\ &= \begin{pmatrix} a^{2m+1}x & a^{2m+1}y + (a^{2m}b + a^{2m-1}bc + \cdots + a^m b c^m + \cdots + b c^{2m})z \\ 0 & c^{2m+1}z \end{pmatrix} = 0. \end{aligned}$$

Thus $x \in r_R(a^{2m+1}) = r_R(a^{2m})$ and $z \in r_R(c^{2m+1}) = r_R(c^{2m}) = \cdots = r_R(c^m)$. Hence the equation

$$a^{2m+1}y + (a^{2m}b + a^{2m-1}bc + \cdots + a^m b c^m + \cdots + b c^{2m})z = 0$$

becomes

$$\begin{aligned} & a^{2m+1}y + (a^{2m}b + a^{2m-1}bc + \cdots + a^{m+1}b c^{m-1})z \\ &= a^{m+1}(a^m y + (a^{m-1}b + a^{m-2}bc + \cdots + b c^{m-1})z) = 0. \end{aligned}$$

Then

$$a^m y + (a^{m-1}b + a^{m-2}bc + \cdots + b c^{m-1})z \in r_R(a^{m+1}),$$

and so

$$a^m y + (a^{m-1}b + a^{m-2}bc + \cdots + b c^{m-1})z \in r_R(a^m).$$

Hence

$$\begin{aligned} & a^m(a^m y + (a^{m-1}b + a^{m-2}bc + \cdots + bc^{m-1})z) \\ &= a^{2m}y + (a^{2m-1}b + a^{2m-2}bc + \cdots + a^m bc^{m-1})z = 0. \end{aligned}$$

Then

$$\begin{aligned} & a^{2m}\beta \\ &= \begin{pmatrix} a^{2m} & a^{2m-1}b + a^{2m-2}bc + \cdots + abc^{2m-2} + bc^{2m-1} \\ 0 & c^{2m} \end{pmatrix} \begin{pmatrix} x & y \\ 0 & z \end{pmatrix} \\ &= \begin{pmatrix} a^{2m}x & a^{2m}y + (a^{2m-1}b + a^{2m-2}bc + \cdots + a^m bc^{m-1} + \cdots + bc^{2m-1})z \\ 0 & c^{2m}z \end{pmatrix} \\ &= \begin{pmatrix} 0 & a^{2m}y + (a^{2m-1}b + a^{2m-2}bc + \cdots + a^m bc^{m-1})z \\ 0 & 0 \end{pmatrix} = 0. \end{aligned}$$

Hence $r_{T_2(R)}(\alpha^{2m+1}) \subseteq r_{T_2(R)}(\alpha^{2m})$ and so $r_{T_2(R)}(\alpha^{2m+1}) = r_{T_2(R)}(\alpha^{2m})$. Therefore $T_2(R)$ is right strongly Hopfian.

Next, we assume that the result is true for $n-1$, $n > 2$, and let

$$A = \begin{pmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ 0 & a_{22} & \cdots & a_{2n} \\ \cdots & \cdots & \cdots & \cdots \\ 0 & 0 & \cdots & a_{nn} \end{pmatrix} \in T_n(R).$$

We show that $r_{T_n(R)}(A) \subseteq r_{T_n(R)}(A^2) \subseteq \cdots$ stabilizes. Put

$$A = \begin{pmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ 0 & a_{22} & \cdots & a_{2n} \\ \cdots & \cdots & \cdots & \cdots \\ 0 & 0 & \cdots & a_{nn} \end{pmatrix} = \begin{pmatrix} A_{n-1} & B \\ 0 & a_{nn} \end{pmatrix}.$$

By the induction hypothesis, we can find $m \in \mathbb{N}$ such that for any $s > m$, $r_{T_{n-1}(R)}(A_{n-1}^s) = r_{T_{n-1}(R)}(A_{n-1}^m)$ and $r_R(a_{nn}^s) = r_R(a_{nn}^m)$. Then using the same way as above, we can show that $r_{T_n(R)}(A^{2m+1}) = r_{T_n(R)}(A^{2m})$ and so $T_n(R)$ is right strongly Hopfian by induction.

(2) \Rightarrow (1) This follows easily from Lemma 2.3.

Corollary 2.5 *Let $L_n(R)$ denote the lower triangular matrix ring over R . Then the following conditions are equivalent:*

- (1) R is right strongly Hopfian;
- (2) $L_n(R)$ is right strongly Hopfian.

Let

$$S_n(R) = \left\{ \begin{pmatrix} a & a_{12} & \cdots & a_{1n} \\ 0 & a & \cdots & a_{2n} \\ \cdots & \cdots & \cdots & \cdots \\ 0 & 0 & \cdots & a \end{pmatrix} \mid a, a_{ij} \in R \right\},$$

$$G_n(R) = \left\{ \left(\begin{array}{cccc} a_1 & a_2 & \cdots & a_n \\ 0 & a_1 & \cdots & a_{n-1} \\ \cdots & \cdots & \cdots & \cdots \\ 0 & 0 & \cdots & a_1 \end{array} \right) \mid a_i \in R, 1 \leq i \leq n \right\},$$

and let $R \bowtie R$ denote the trivial extension of R by R .

Corollary 2.6 *The following conditions are equivalent:*

- (1) R is right strongly hopfian;
- (2) $S_n(R)$ is right strongly Hopfian;
- (3) $G_n(R)$ is right strongly Hopfian;
- (4) $R[x]/(x^n)$ is right strongly Hopfian;
- (5) $R \bowtie R$ is right strongly Hopfian.

Proof Note that $R[x]/(x^n) \cong G_n(R)$ and $R \bowtie R \cong G_2(R)$.

Let R be a ring. Immediately, we deduce that the lower triangular matrix ring over R is right strongly Hopfian if and only if the upper triangular matrix ring is right strongly Hopfian. Let R be a ring, and let $W(R) = \left\{ \left(\begin{array}{ccc} a_{11} & 0 & 0 \\ a_{21} & a_{22} & a_{23} \\ 0 & 0 & a_{33} \end{array} \right) \mid a_{ij} \in R \right\}$.

Then $W(R)$ is a 3×3 subring of $M_3(R)$ under usual matrix addition and multiplication. A natural problem asks if the right strongly Hopfian property of such a ring coincides with that of R . This inspires us to consider the right strongly Hopfian property of $W(R)$.

Proposition 2.7 *Let R be a ring. Then $W(R)$ is right strongly Hopfian if and only if R is right strongly Hopfian.*

Proof Suppose R is right strongly Hopfian and let

$$\alpha = \begin{pmatrix} a & 0 & 0 \\ x & b & y \\ 0 & 0 & c \end{pmatrix} \in W(R).$$

Then there exists $m \in \mathbb{N}$ such that for any $n > m$, $r_R(a^n) = r_R(a^m)$, $r_R(b^n) = r_R(b^m)$, and $r_R(c^n) = r_R(c^m)$. Now we show that $r_{W(R)}(\alpha^{2m+1}) = r_{W(R)}(\alpha^{2m})$. If

$$\beta = \begin{pmatrix} d & 0 & 0 \\ s & e & t \\ 0 & 0 & f \end{pmatrix} \in r_{W(R)}(\alpha^{2m+1}),$$

then

$$\alpha^{2m+1}\beta = \begin{pmatrix} a^{2m+1}d & 0 & 0 \\ ud + b^{2m+1}s & b^{2m+1}e & b^{2m+1}t + vf \\ 0 & 0 & c^{2m+1}f \end{pmatrix} = 0,$$

where

$$u = xa^{2m} + bxa^{2m-1} + \cdots + b^m xa^m + b^{m+1} xa^{m-1} + \cdots + b^{2m-1} xa + b^{2m} x,$$

and

$$v = b^{2m} y + b^{2m-1} yc + b^{2m-2} yc^2 + \cdots + byc^{2m-1} + yc^{2m}.$$

Hence

$$d \in r_R(a^{2m+1}) = r_R(a^{2m}) = \cdots = r_R(a^m),$$

$$e \in r_R(b^{2m+1}) = r_R(b^{2m}) = \cdots = r_R(b^m),$$

and

$$f \in r_R(c^{2m+1}) = r_R(c^{2m}) = \cdots = r_R(c^m).$$

Then

$$\begin{aligned} 0 &= ud + b^{2m+1}s \\ &= (xa^{2m} + bxa^{2m-1} + \cdots + b^m xa^m + b^{m+1} xa^{m-1} + \cdots + b^{2m} x)d + b^{2m+1}s \\ &= (b^{m+1} xa^{m-1} + b^{m+2} xa^{m-2} + \cdots + b^{2m} x)d + b^{2m+1}s \\ &= b^{m+1}((xa^{m-1} + bxa^{m-2} + \cdots + b^{m-1} x)d + b^m s), \end{aligned}$$

and

$$\begin{aligned} 0 &= b^{2m+1}t + vf \\ &= b^{2m+1}t + (b^{2m}y + b^{2m-1}yc + \cdots + b^{m+1}yc^{m-1} + b^m yc^m + \cdots + yc^{2m})f \\ &= b^{2m+1}t + (b^{2m}y + b^{2m-1}yc + \cdots + b^{m+1}yc^{m-1})f \\ &= b^{m+1}(b^m t + (b^{m-1}y + b^{m-2}yc + \cdots + yc^{m-1})f). \end{aligned}$$

Hence

$$(xa^{m-1} + bxa^{m-2} + \cdots + b^{m-1}x)d + b^m s \in r_R(b^{m+1}) = r_R(b^m)$$

and

$$b^m t + (b^{m-1}y + b^{m-2}yc + \cdots + yc^{m-1})f \in r_R((b^{m+1})) = r_R(b^m).$$

So

$$\begin{aligned} &b^m((xa^{m-1} + bxa^{m-2} + \cdots + b^{m-1}x)d + b^m s) \\ &= b^m xa^{m-1} + b^{m+1} xa^{m-2} + \cdots + b^{2m-1} x)d + b^{2m} s = 0. \end{aligned}$$

and

$$\begin{aligned} &b^m(b^m t + (b^{m-1}y + b^{m-2}yc + \cdots + yc^{m-1})f) \\ &= b^{2m} t + (b^{2m-1}y + b^{2m-2}yc + \cdots + b^m yc^{m-1})f = 0. \end{aligned}$$

Then by a routine computations, we can show that $\alpha^{2m}\beta = 0$ and so $\beta \in r_{W(R)}(\alpha^{2m})$. Hence $r_{W(R)}(\alpha^{2m+1}) = r_{W(R)}(\alpha^{2m})$. Therefore $W(R)$ is right strongly Hopfian.

Conversely, if $W(R)$ is right strongly Hopfian, then by Lemma 2.3, R is right strongly Hopfian.

Let α be an endomorphism and δ an α -derivation of R . We denote by $R[x; \alpha, \delta]$ the Ore extension whose elements are the polynomials over R , the addition is defined

as usual and the multiplication is subject to the relation $xa = \alpha(a)x + \delta(a)$ for any $a \in R$. From this rule, an inductive argument can be made in order to calculate an expression for $x^j a$, for all $j \in \mathbb{N}$ and $a \in R$. To recall this result, we shall use some convenient notation introduced in [9].

Notation 2.8 Let δ be an α -derivation of R . For integers i, j with $0 \leq i \leq j$, $f_i^j \in \text{End}(R, +)$ will denote the map which is the sum of all possible words in α, δ built with i letters α and $j - i$ letters δ . For instance, $f_0^0 = 1, f_j^j = \alpha^j, f_0^j = \delta^j$ and $f_{j-1}^j = \alpha^{j-1}\delta + \alpha^{j-2}\delta\alpha + \dots + \delta\alpha^{j-1}$. Using recursive formulas for the f_i^j 's and induction, as done in [9], one can show with a routine computation that

$$x^j a = \sum_{i=0}^j f_i^j(a) x^i.$$

The following Lemma is well known and we omit the proof (see [3, Lemma 2.1]).

Lemma 2.9 Let R be an (α, δ) -compatible ring. Then we have the following:

- (1) If $ab = 0$, then $a\alpha^n(b) = \alpha^n(a)b = 0$ for all positive integers n .
- (2) If $\alpha^k(a)b = 0$ for some positive integer k , then $ab = 0$.
- (3) If $ab = 0$, then $\alpha^n(a)\delta^m(b) = 0 = \delta^m(a)\alpha^n(b)$ for all positive integers m, n .
- (4) If $ab = 0$, then $a f_i^j(b) = 0$ and $f_i^j(a)b = 0$ for all i, j .

Lemma 2.10 Let R be an α -compatible ring. If $\alpha^{k_1}(a_1)\alpha^{k_2}(a_2)\dots\alpha^{k_n}(a_n) = 0$ for some positive integers, then $a_1 a_2 \dots a_n = 0$.

Proof Using induction, for $n = 1$, the result is true by the injectivity of α . Now suppose $\alpha^{k_1}(a_1)\alpha^{k_2}(a_2)\dots\alpha^{k_n}(a_n) = 0$. Then $\alpha^{k_1}(a_1)\alpha^{k_2}(a_2)\dots\alpha^{k_{n-1}}(a_{n-1})a_n = 0$, and so $\alpha^{k_1}(a_1)\alpha^{k_2}(a_2)\dots\alpha^{k_{n-1}}(a_{n-1}a_n) = 0$. Then $a_1 a_2 \dots a_n = 0$.

Lemma 2.11 Let R be an (α, δ) -compatible ring, $f(x) = a_0 + a_1x + \dots + a_nx^n$ and $g(x) = b_0 + b_1x + \dots + b_mx^m$ be two polynomials in $R[x; \alpha, \delta]$. Then we have the following:

- (1) If for all $0 \leq i \leq n$ and $0 \leq j \leq m$, $a_i b_j = 0$, then $f(x)g(x) = 0$.
- (2) If R is semicommutative and $c \in R$ is such that for all $0 \leq j \leq m$, $cb_j = 0$, then $cf(x)g(x) = 0$.

Proof (1) We have

$$\begin{aligned} f(x)g(x) &= (a_0 + a_1x + \dots + a_nx^n)(b_0 + b_1x + \dots + b_mx^m) \\ &= \sum_{l=0}^{m+n} \left(\sum_{s+t=l} (\sum_{i=s}^n a_i f_s^i(b_t)) \right) x^l. \end{aligned}$$

By Lemma 2.9, $a_i b_t = 0$ implies $a_i f_s^i(b_t) = 0$. Thus it is easy to see that $f(x)g(x) = 0$.

(2) Since R is semicommutative, for all $0 \leq i \leq n$ and $0 \leq j \leq m$, $cb_j = 0$ implies $ca_i b_j = 0$. Thus by (1) we complete the proof.

For two polynomials $f(x)$ and $g(x)$ in $R[x; \alpha, \delta]$, in order to calculate a expression for $(f(x) + g(x))^n$, for all $n \in \mathbb{N}$, we denote by $[Q_i^n f(x)g(x)]$ the polynomial which is the sum of all possible terms, which each term is a product of i polynomials $f(x)$ and $n - i$ polynomials $g(x)$. Using this convenient notation, we have $(f(x) + g(x))^n = f(x)^n + [Q_{n-1}^n f(x)g(x)] + [Q_{n-2}^n f(x)g(x)] + \cdots + [Q_1^n f(x)g(x)] + g(x)^n$.

Lemma 2.12 *Let R be an (α, δ) -compatible semicommutative ring, ax^r , $f(x) = b_0 + b_1x + \cdots + b_mx^m$, $g(x) = c_0 + c_1x + \cdots + c_qx^q$ be three polynomials in $R[x; \alpha, \delta]$. If $c_j \in r_R(a^n)$ for all $0 \leq j \leq q$, then for any $p > n$, $[Q_n^p(ax^r)f(x)]g(x) = 0$.*

Proof It is easy to check that the coefficients of $[Q_n^p(ax^r)f(x)]$ can be written as sums of monomials of length p in $f_s^t(a)$ and $f_u^v(b_j)$, where $b_j \in \{b_0, b_1, \dots, b_m\}$ and $t \geq s \geq 0$, $v \geq u \geq 0$ are nonnegative positive integers. Consider each monomial $f_{s_1}^{t_1}(v_1)f_{s_2}^{t_2}(v_2) \cdots f_{s_p}^{t_p}(v_p)$ where $v_1, v_2, \dots, v_p \in \{a, b_0, b_1, \dots, b_m\}$. It would contains n letters a . Suppose $v_{r_1} = v_{r_2} = \cdots = v_{r_n} = a$ for some $1 \leq r_1 < r_2 < \cdots < r_n \leq p$. Then we write the monomial $f_{s_1}^{t_1}(v_1)f_{s_2}^{t_2}(v_2) \cdots f_{s_p}^{t_p}(v_p)$ as

$$f_{s_1}^{t_1}(v_1) \cdots f_{s_{r_1}}^{t_{r_1}}(a)f_{s_{r_1+1}}^{t_{r_1+1}}(v_{r_1+1}) \cdots f_{s_{r_n-1}}^{t_{r_n-1}}(v_{r_n-1})f_{s_{r_n}}^{t_{r_n}}(a)f_{s_{r_n+1}}^{t_{r_n+1}}(v_{r_n+1}) \cdots f_{s_p}^{t_p}(v_p),$$

where $v_s \in \{b_0, b_1, \dots, b_m\}$ if $s \notin \{r_1, r_2, \dots, r_n\}$. For each $0 \leq j \leq q$, since R is (α, δ) -compatible and semicommutative, $a^n c_j = aa \cdots ac_j = 0$ implies

$$f_{s_{r_1}}^{t_{r_1}}(a)f_{s_{r_2}}^{t_{r_2}}(a) \cdots f_{s_{r_n}}^{t_{r_n}}(a)c_j = 0,$$

and so

$$f_{s_1}^{t_1}(v_1) \cdots f_{s_{r_1}}^{t_{r_1}}(a)f_{s_{r_1+1}}^{t_{r_1+1}}(v_{r_1+1}) \cdots f_{s_{r_n-1}}^{t_{r_n-1}}(v_{r_n-1})f_{s_{r_n}}^{t_{r_n}}(a)f_{s_{r_n+1}}^{t_{r_n+1}}(v_{r_n+1}) \cdots f_{s_p}^{t_p}(v_p)c_j = 0.$$

Thus by Lemma 2.11, we complete the proof.

The same idea can be used to prove the following.

Corollary 2.13 *Let R be an (α, δ) -compatible semicommutative ring, ax^r , $f(x) = b_0 + b_1x + \cdots + b_mx^m$, $g(x) = c_0 + c_1x + \cdots + c_qx^q$ be three polynomials in $R[x; \alpha, \delta]$. If $c_j \in r_R(a^n)$ for all $0 \leq j \leq q$, Then we have the following:*

- (1) For any $p > n + l$, $[Q_{n+l}^p(ax^r)f(x)]g(x) = 0$.
- (2) $R[x; \alpha, \delta](a^{i_1}x^{n_1})R[x; \alpha, \delta](a^{i_2}x^{n_2})R[x; \alpha, \delta] \cdots (a^{i_k}x^{n_k})R[x; \alpha, \delta]g(x) = 0$ if $i_1 + i_2 + \cdots + i_k \geq n$.

Proposition 2.14 *Let R be (α, δ) -compatible and $R[x; \alpha, \delta]$ be reversible. Then the following conditions are equivalent:*

- (1) R is right strongly Hopfian;
- (2) $R[x; \alpha, \delta]$ is right strongly Hopfian.

Proof (1) \Rightarrow (2) Let $f(x) = a_0 + a_1x + \cdots + a_nx^n \in R[x; \alpha, \delta]$. Since R is right strongly Hopfian, there exists $k \in \mathbb{N}$ such that for all $l > k$ and all $0 \leq i \leq n$, $r_R(a_i^l) = r_R(a_i^k)$. Now we show that $r_{R[x; \alpha, \delta]}(f(x)^{(n+1)k+1}) = r_{R[x; \alpha, \delta]}(f(x)^{(n+1)k})$. If

$$g(x) = b_0 + b_1x + \cdots + b_mx^m \in r_{R[x; \alpha, \delta]}(f(x)^{(n+1)k+1}),$$

then

$$\begin{aligned} 0 &= f(x)^{(n+1)k+1}g(x) = (a_0 + a_1x + \cdots + a_nx^n)^{(n+1)k+1}(b_0 + b_1x + \cdots + b_mx^m) \\ &= a_n\alpha^n(a_n)\alpha^{2n}(a_n) \cdots \alpha^{(n+1)kn}(a_n)\alpha^{[(n+1)k+1]n}(b_m)x^{[(n+1)k+1]n+m} + \text{lower terms.} \end{aligned}$$

Hence

$$a_n\alpha^n(a_n)\alpha^{2n}(a_n) \cdots \alpha^{(n+1)kn}(a_n)\alpha^{[(n+1)k+1]n}(b_m) = 0.$$

By Lemma 2.10, we obtain $a_n^{(n+1)k+1}b_m = 0$. Hence

$$b_m \in r_R(a_n^{(n+1)k+1}) = r_R(a_n^k).$$

From $f(x)^{(n+1)k+1}g(x) = 0$, we have $a_n^k f(x)^{(n+1)k+1}g(x) = 0$. Then by Lemma 2.11, we obtain

$$\begin{aligned} 0 &= a_n^k f(x)^{(n+1)k+1}g(x) = a_n^k(a_0 + a_1x + \cdots + a_nx^n)^{(n+1)k+1}(b_0 + b_1x + \cdots + b_mx^m) \\ &= a_n^k(a_0 + a_1x + \cdots + a_nx^n)^{(n+1)k+1}(b_0 + b_1x + \cdots + b_{m-1}x^{m-1}) \\ &\quad + a_n^k(a_0 + a_1x + \cdots + a_nx^n)^{(n+1)k+1}b_mx^m \\ &= a_n^k(a_0 + a_1x + \cdots + a_nx^n)^{(n+1)k+1}(b_0 + b_1x + \cdots + b_{m-1}x^{m-1}) \\ &= a_n^k a_n \alpha^n(a_n) \cdots \alpha^{(n+1)kn}(a_n)\alpha^{[(n+1)k+1]n}(b_{m-1})x^{[(n+1)k+1]n+m-1} + \text{lower terms.} \end{aligned}$$

Hence

$$a_n^{k+1}\alpha^n(a_n)\alpha^{2n}(a_n) \cdots \alpha^{(n+1)kn}(a_n)\alpha^{[(n+1)k+1]n}(b_{m-1}) = 0$$

and so

$$b_{m-1} \in r_R(a_n^{(n+2)k+1}) = r_R(a_n^k).$$

Using the same method repeatedly, we obtain

$$b_j \in r_R(a_n^{(n+1)k+1}) = r_R(a_n^k) \text{ for all } 0 \leq j \leq m.$$

Consider the polynomial $f(x)$ as the sum of two polynomials a_nx^n and $h(x) = a_{n-1}x^{n-1} + a_{n-2}x^{n-2} + \cdots + a_0$. Then by Corollary 2.13, we obtain

$$\begin{aligned} 0 &= f(x)^{(n+1)k+1}g(x) = (a_nx^n + h(x))^{(n+1)k+1}g(x) \\ &= (a_nx^n)^{(n+1)k+1}g(x) + \left[Q_{(n+1)k}^{(n+1)k+1}(a_nx^n)h(x) \right] g(x) + \cdots \\ &\quad + \left[Q_k^{(n+1)k+1}(a_nx^n)h(x) \right] g(x) + \left[Q_{k-1}^{(n+1)k+1}(a_nx^n)h(x) \right] g(x) \\ &\quad + \cdots + \left[Q_1^{(n+1)k+1}(a_nx^n)h(x) \right] g(x) + h(x)^{(n+1)k+1}g(x) \\ &= \left[Q_{k-1}^{(n+1)k+1}(a_nx^n)h(x) \right] g(x) + \left[Q_{k-2}^{(n+1)k+1}(a_nx^n)h(x) \right] g(x) + \cdots \\ &\quad + \left[Q_1^{(n+1)k+1}(a_nx^n)h(x) \right] g(x) + h(x)^{(n+1)k+1}g(x). \end{aligned} \quad (1)$$

Multiplying equation (1) on the left side by $(a_n x^n)^{k-1}$, then by Lemma 2.12 and Corollary 2.13, we obtain

$$(a_n x^n)^{k-1} h(x)^{(n+1)k+1} g(x) = 0.$$

Since $R[x; \alpha, \delta]$ is reversible, this implies

$$\left[Q_{k-1}^{(n+1)k+1} (a_n x^n) h(x) \right] g(x) h(x)^{k-1} = 0. \quad (2)$$

Multiplying equation (1) on the right side by $h(x)^{k-1}$, we obtain

$$\begin{aligned} & \left[Q_{k-2}^{(n+1)k+1} (a_n x^n) h(x) \right] g(x) h(x)^{k-1} + \left[Q_{k-3}^{(n+1)k+1} (a_n x^n) h(x) \right] g(x) h(x)^{k-1} \\ & + \cdots + h(x)^{(n+1)k+1} g(x) h(x)^{k-1} = 0. \end{aligned} \quad (3)$$

Multiplying equation (3) on the left side by $(a_n x^n)^{k-2}$, we obtain

$$(a_n x^n)^{k-2} \left[Q_1^{(n+1)k+1} (a_n x^n) h(x) \right] g(x) h(x)^{k-1} + (a_n x^n)^{k-2} h(x)^{(n+1)k+1} g(x) h(x)^{k-1} = 0. \quad (4)$$

By equation $(a_n x^n)^{k-1} h(x)^{(n+1)k+1} g(x) = 0$ and $R[x; \alpha, \delta]$ is reversible, it is easy to see that $(a_n x^n)^{k-2} \left[Q_1^{(n+1)k+1} (a_n x^n) h(x) \right] g(x) h(x)^{k-1} = 0$. Hence equation (4) becomes

$$(a_n x^n)^{k-2} h(x)^{(n+1)k+1} g(x) h(x)^{k-1} = 0.$$

Since $R[x; \alpha, \delta]$ is reversible, this implies

$$\left[Q_{k-2}^{(n+1)k+1} (a_n x^n) h(x) \right] g(x) h(x)^{k-1} h(x)^{k-2} = 0.$$

Multiplying equation (3) on the right side by $h(x)^{k-2}$, we obtain

$$\left[Q_{k-3}^{(n+1)k+1} (a_n x^n) h(x) \right] g(x) h(x)^{k-1} h(x)^{k-2} + \cdots + h(x)^{(n+1)k+1} g(x) h(x)^{k-1} h(x)^{k-2} = 0.$$

Continue this process yields that

$$h(x)^{(n+1)k+1} g(x) h(x)^{k-1} h(x)^{k-2} \cdots h(x) = 0,$$

and so

$$\begin{aligned} & h(x)^{(n+1)k+1 + \frac{k(k-1)}{2}} g(x) \\ & = (a_0 + a_1 x + \cdots + a_{n-1} x^{n-1})^{(n+1)k+1 + \frac{k(k-1)}{2}} (b_0 + b_1 x + \cdots + b_m x^m) = 0, \end{aligned}$$

since $R[x; \alpha, \delta]$ is reversible. Now by the same way as above, we obtain

$$b_j \in r_R(a_{n-1}^{(n+1)k+1 + \frac{k(k-1)}{2}}) = r_R(a_{n-1}^k)$$

for all $0 \leq j \leq m$. Using induction on n , we obtain

$$b_j \in r_R(a_i^{(n+1)k+1}) = r_R(a_i^k)$$

for all $0 \leq j \leq m$ and $0 \leq i \leq n$. It is now easy to check that $f(x)^{(n+1)k} g(x) = 0$. Hence $r_{R[x; \alpha, \delta]}(f(x)^{(n+1)k+1}) = r_{R[x; \alpha, \delta]}(f(x)^{(n+1)k})$. Therefore $R[x; \alpha, \delta]$ is right strongly Hopfian.

(2) \Rightarrow (1) It is trivial.

Corollary 2.15 *We have the following:*

(1) *If R is α -compatible and the skew polynomial ring $R[x; \alpha]$ is reversible, then the skew polynomial ring $R[x; \alpha]$ is right strongly Hopfian if and only if R is right strongly Hopfian.*

(2) *If R is δ -compatible and the differential polynomial ring $R[x; \delta]$ is reversible, then the differential polynomial ring $R[x; \delta]$ is right strongly Hopfian if and only if R is right strongly Hopfian.*

Proof It is an immediate consequence of Proposition 2.14.

Corollary 2.16 ([5, Theorem 5.1]). *Let R be a commutative strongly Hopfian ring, then the polynomial ring $R[x]$ is strongly Hopfian.*

Let M be a multiplicative monoid. In the following, e will always stand for the identity of M . Then $R[M]$ will denote the monoid ring over R consisting of all elements of the form $\sum_{i=1}^n r_i g_i$ with $r_i \in R$, $g_i \in M$, $i = 1, 2, \dots, n$, where the addition is given naturally and the multiplication is given by

$$\left(\sum_{i=1}^n r_i g_i\right)\left(\sum_{j=1}^m s_j h_j\right) = \sum_{i=1}^n \sum_{j=1}^m (r_i s_j)(g_i h_j).$$

Recall that the ordered monoid (M, \leq) is a strictly ordered monoid if for any $g, g', h \in M$, $g < g'$ implies that $gh < g'h$ and $hg < hg'$.

For two elements α and β in $R[M]$, in order to calculate a expression for $(\alpha + \beta)^n$, for all $n \in \mathbb{N}$, we denote by $[Q_i^n \alpha \beta]$ the sum of all possible terms which each term is a product of i elements α and $n - i$ elements β . Using this convenient notation, we have $(\alpha + \beta)^n = \alpha^n + [Q_{n-1}^n \alpha \beta] + [Q_{n-2}^n \alpha \beta] + \dots + [Q_1^n \alpha \beta] + \beta^n$.

Lemma 2.17 *Let (M, \leq) be a strictly totally ordered monoid and R a semicommutative ring, $\alpha = ag$, $\beta = b_1 h_1 + b_2 h_2 + \dots + b_n h_n$ and $\gamma = c_1 v_1 + c_2 v_2 + \dots + c_m v_m$ be three elements in $R[M]$. If there exists a positive integer $n \in \mathbb{Z}$ such that $c_j \in r_R(a^n)$ for all $1 \leq j \leq m$, then for any $p > n$, $[Q_n^p \alpha \beta] \gamma = 0$.*

Proof The coefficients of $[Q_n^p \alpha \beta]$ can be written as sums of monomials of length p in a and b_j , where $b_j \in \{b_1, b_2, \dots, b_n\}$. Consider one of such monomials, $d_1 d_2 \dots d_p$, where $d_i \in \{a, b_1, b_2, \dots, b_n\}$, $0 \leq i \leq p$. It would contain n letters a . Suppose $d_{r_1} = d_{r_2} = \dots = d_{r_n} = a$ for some $1 \leq r_1 < r_2 < \dots < r_n \leq p$. Then we can written the monomial as $d_1 d_2 \dots d_{r_1-1} a d_{r_1+1} \dots d_{r_2-1} a d_{r_2+1} \dots d_{r_n-1} a d_{r_n+1} \dots d_p$. Since R is semicommutative and $c_j \in r_R(a^n)$ for all $1 \leq j \leq m$, $a^n c_j = a a \dots a c_j = 0$ implies

$$d_1 d_2 \dots d_{r_1-1} a d_{r_1+1} \dots d_{r_n-1} a d_{r_n+1} \dots d_p c_j = 0$$

for all $1 \leq j \leq m$. Hence each monomial appears in $[Q_n^p \alpha \beta] \gamma$ is equal to 0. Therefore $[Q_n^p \alpha \beta] \gamma = 0$.

The same idea can be used to prove the following.

Corollary 2.18 *Let (M, \leq) be a strictly totally ordered monoid and R a semicommutative ring, $\alpha = ag$, $\beta = b_1h_1 + b_2h_2 + \cdots + b_nh_n$ and $\gamma = c_1v_1 + c_2v_2 + \cdots + c_mv_m$ be three elements in $R[M]$. If there exists a positive integer $n \in \mathbb{Z}$ such that $c_j \in r_R(a^n)$ for all $1 \leq j \leq m$, then for any $p > n + l$, we have the following:*

- (1) $[Q_{n+l}^p \alpha \beta] \gamma = 0$.
- (2) $R[M](ag)^{n_1} R[M](ag)^{n_2} R[M] \cdots (ag)^{n_k} R[M] \gamma R[M] = 0$ if $n_1 + n_2 + \cdots + n_k \geq n$.

Proposition 2.19 *Let M be a strictly totally ordered monoid and $R[M]$ a reversible ring. Then the following conditions are equivalent:*

- (1) R is a right strongly Hopfian ring;
- (2) $R[M]$ is a right strongly Hopfian ring.

Proof (1) \Rightarrow (2). Let $\alpha = a_1g_1 + a_2g_2 + \cdots + a_ng_n \in R[M]$ with $g_i < g_j$ if $i < j$. Since R is right strongly Hopfian, there exists $k \in \mathbb{N}$ such that for all $l > k$ and all $1 \leq i \leq n$, $r_R(a_i^l) = r_R(a_i^k)$. Now we show that $r_{R[M]}(\alpha^{nk+1}) = r_{R[M]}(\alpha^{nk})$. If $\beta = b_1h_1 + b_2h_2 + \cdots + b_mh_m \in r_{R[M]}(\alpha^{nk+1})$ with $h_s < h_t$ if $s < t$. Then

$$0 = \alpha^{nk+1}\beta = (a_1g_1 + a_2g_2 + \cdots + a_ng_n)^{nk+1}(b_1h_1 + b_2h_2 + \cdots + b_mh_m).$$

Consider the coefficient of the largest element $g_n^{nk+1}h_m$ in $\alpha^{nk+1}\beta$, we obtain $a_n^{nk+1}b_m = 0$. Hence $b_m \in r_R(a_n^{nk+1}) = r_R(a_n^k)$. From $\alpha^{nk+1}\beta = 0$, we have

$$\begin{aligned} 0 &= (a_n^k e) \alpha^{nk+1} \beta = (a_n^k e) (a_1g_1 + a_2g_2 + \cdots + a_ng_n)^{nk+1} (b_1h_1 + b_2h_2 + \cdots + b_mh_m) \\ &= (a_n^k e) (a_1g_1 + a_2g_2 + \cdots + a_ng_n)^{nk+1} (b_1h_1 + b_2h_2 + \cdots + b_{m-1}h_{m-1}) \\ &\quad + (a_n^k e) (a_1g_1 + a_2g_2 + \cdots + a_ng_n)^{nk+1} b_m h_m \\ &= (a_n^k e) (a_1g_1 + a_2g_2 + \cdots + a_ng_n)^{nk+1} (b_1h_1 + b_2h_2 + \cdots + b_{m-1}h_{m-1}) \\ &= (a_n^k e) \alpha^{nk+1} (\beta - b_m h_m). \end{aligned}$$

Consider the coefficient of the largest element $g_n^{nk+1}h_{m-1}$ in $(a_n^k e) \alpha^{nk+1} (\beta - b_m h_m)$, we obtain

$$b_{m-1} \in r_R(a_n^{(n+1)k+1}) = r_R(a_n^k).$$

Continue this process yields that $b_j \in r_R(a_n^{nk+1}) = r_R(a_n^k)$ for all $1 \leq j \leq m$. Consider the element α as the sum of two elements $a_n g_n$ and $\gamma = a_1g_1 + a_2g_2 + \cdots + a_{n-1}g_{n-1}$. Then by Lemma 2.17 and Corollary 2.18, we obtain

$$\begin{aligned} 0 &= \alpha^{nk+1}\beta = (a_n g_n + \gamma)^{nk+1}\beta \\ &= (a_n g_n)^{nk+1}\beta + [Q_{nk}^{nk+1}(a_n g_n)\gamma] \beta + \cdots + [Q_{k-1}^{nk+1}(a_n g_n)\gamma] \beta + \cdots + \gamma^{nk+1}\beta \\ &= [Q_{k-1}^{nk+1}(a_n g_n)\gamma] \beta + [Q_{k-2}^{nk+1}(a_n g_n)\gamma] \beta + \cdots + \gamma^{nk+1}\beta. \quad (5) \end{aligned}$$

Multiplying equation (5) on the left side by $(a_n g_n)^{k-1}$, then by Corollary 2.18, we obtain $(a_n g_n)^{k-1} \gamma^{nk+1} \beta = 0$ and so $[Q_{k-1}^{nk+1}(a_n g_n)\gamma] \beta \gamma^{k-1} = 0$ since $R[M]$ is reversible. Multiplying equation (5) on the right side by γ^{k-1} , we obtain

$$[Q_{k-2}^{nk+1}(a_n g_n)\gamma] \beta \gamma^{k-1} + [Q_{k-3}^{nk+1}(a_n g_n)\gamma] \beta \gamma^{k-1} + \cdots + \gamma^{nk+1} \beta \gamma^{k-1} = 0. \quad (6)$$

Multiplying equation (6) on the left side by $(a_n g_n)^{k-2}$, we obtain

$$(a_n g_n)^{k-2} [Q_1^{nk+1}(a_n g_n)\gamma] \beta \gamma^{k-1} + (a_n g_n)^{k-2} \gamma^{nk+1} \beta \gamma^{k-1} = 0.$$

Since $R[M]$ is reversible, $(a_n g_n)^{k-1} \gamma^{nk+1} \beta = 0$ implies

$$(a_n g_n)^{k-2} [Q_1^{nk+1}(a_n g_n)\gamma] \beta \gamma^{k-1} = 0.$$

Hence we obtain $(a_n g_n)^{k-2} \gamma^{nk+1} \beta \gamma^{k-1} = 0$, and so $[Q_{k-2}^{nk+1}(a_n g_n)\gamma] \beta \gamma^{k-1} \gamma^{k-2} = 0$ since $R[M]$ is reversible. Then multiplying equation (6) on the right side by γ^{k-2} , we obtain

$$[Q_{k-3}^{nk+1}(a_n g_n)\gamma] \beta \gamma^{k-1} \gamma^{k-2} + [Q_{k-4}^{nk+1}(a_n g_n)\gamma] \beta \gamma^{k-1} \gamma^{k-2} + \dots + \gamma^{nk+1} \beta \gamma^{k-1} \gamma^{k-2} = 0.$$

Continue this process we obtain $\gamma^{nk+1} \beta \gamma^{\frac{k(k-1)}{2}} = 0$ and so $\gamma^{nk+1+\frac{k(k-1)}{2}} \beta = 0$. Using the same way as above, we can show that

$$b_j \in r_R(a_{n-1}^{nk+1+\frac{k(k-1)}{2}}) = r_R(a_{n-1}^k)$$

for all $1 \leq j \leq m$. Using induction on n , we obtain

$$b_j \in r_R(a_i^{nk+1}) = r_R(a_i^k)$$

for all $1 \leq j \leq m$ and $1 \leq i \leq n$. Then it is easy to check that $\alpha^{nk} \beta = 0$. Hence $\beta \in r_{R[M]}(\alpha^{nk})$, and so $r_{R[M]}(\alpha^{nk+1}) = r_{R[M]}(\alpha^{nk})$. Therefore $R[M]$ is right strongly Hopfian.

(2) \Rightarrow (1) It is trivial.

Corollary 2.20 ([5, Corollary 5.4]) *Let R be a commutative strongly Hopfian ring, then $R[x, x^{-1}]$ is strongly Hopfian.*

Acknowledgements: The authors wish to express their hearty thanks to the referees for their helpful comments and suggestions that greatly improved the exposition.

References

- [1] S. Annin, Associated primes over skew polynomials, *Comm. Algebra*, **30** (2002), no. 5, 2511-2528.
- [2] R. Gilmer, W. Heinzer, The Laskerian property, power series rings, and Noetherian spectra, *Proc. Amer. Math. Soc.* **79** (1980), no. 1, 13-16.
- [3] E. Hashemi, A. Moussavi, Polynomial extensions of quasi-Baer rings, *Acta Math Hungar*, **107** (2005), no. 3, 207-224.
- [4] S. Hizem, Formal power series over strongly Hopfian rings, *Comm. Algebra*, **39** (2011), 279-291.

- [5] A. Hmaimou, A. Kaidi, E. Sanchez Campos, Generalized fitting modules and rings, *J. Algebra*, **308** (2007), 199-214.
- [6] N. K. Kim, Y. Lee, Extensions of reversible rings, *J. pure Appl. Algebra*, **185** (2003), 207-223.
- [7] Chin-Pi Lu, Modules satisfying acc on certain type of colons, *Pacific Journal of Mathematics*, **131** (1988), no. 2, 303-318.
- [8] Chin-Pi Lu, Modules and rings satisfying (accr), *Proc. Ams*, **117** (1993), no. 1, 5-10.
- [9] T. Y. Lam, A. Leroy, J. Matczuk, Primeness, semiprimeness and prime radical of ore extensions, *Comm. Algebra*, **25** (1997), no. 8, 2459-2506.