

Bayesian Estimation of Exponentiated Weibull Distribution under Partially Acceleration Life Tests

¹Abd EL-Baset A. Ahmad, ²Ahmed A. Soliman, ¹Manal M. Yousef;

¹*Department of Statistics, Faculty of Science, Assiut University,*

²*Department of Statistics, Faculty of Science, Sohage University,*

Abstract. This paper presents statistical methods for analyzing partially accelerated life test from constant-stress test. The lifetime of items under use condition follows the two parameter exponentiated Weibull distribution. Based on progressively Type-II censored sample, the classical maximum likelihood method as well as a fully Bayesian method based on Lindley (1980) approximation form and the Markov chain Monte Carlo technique are developed for inference about model parameters and acceleration factor. Furthermore, approximate confidence intervals and credible intervals are presented. A Monte Carlo simulation is given to study the precision of different estimators.

Keywords: Constant-stress partially accelerated life test; Exponentiated Weibull; Progressive Type-II censoring; Maximum likelihood method; Bayesian estimation; Lindley's approximation; Markov chain Monte Carlo; Monte Carlo simulation.

1 Introduction

Accelerated life test (ALT) is a popular experimental strategy to get information on life distribution of highly reliable product. Test units under normal operating conditions are often extremely reliable with large mean times to fail. To overcome such a problem, ALT experiments preferred to obtain the reliability information of product components in a short period of time by subjecting it to higher than usual stress (pressure, temperature, voltage,....,etc.). In ALT, items are tested under stress levels that are far more than those normally applied in practice with the conclusive aim of inducing early failure. At the accelerated conditions, the test data obtained are analyzed in terms of a mathematical model, and then extrapolated to the normal stress to estimate the life distribution parameters. In some cases, such mathematical model which relating the lifetime of the unit and the stress is not known, i.e.,

the data obtained from ALT cannot be extrapolated to normal stress. So, in such cases, partially accelerated life test (PALT) is better criterion to use. The objective of a PALT is to collect more failure data in a limited time without necessarily using high stresses to all test units.

In ALT, all units are run only at accelerated conditions, while in PALT they are run at both normal and accelerated conditions. The stress can be applied in various ways: commonly used methods are constant-stress and step-stress. In constant-stress PALT, the total test units are first divided into two groups. Items of group 1 are allocated to normal condition and items of group 2 are allocated to accelerated condition. Each unit is run at a constant level of stress until the unit fails or censored.

Several authors have dealt with problems of constant-stress ALT. Based on progressively Type-II censored samples, Abdel-Hamid (2009) considered constant-stress PALT when the lifetime of units under use condition follows the Burr Type-XII distribution. Bayes and maximum likelihood (ML) methods of estimation are applied on constant-stress ALTs to finite mixtures of distributions by AL-Hussaini and Abdel -Hamid (2004,2006).

In the step-stress PALT, a test unit is first run at use condition and if it does not fail in a specified time, then it is run at accelerated condition until failures occurs or the observation censored, see, for example, Abd-Elfattah et al. (2008). Most of the current literature on the accelerated life testing problem is based on the sample theoretic methods as discussed in Nelson (1990).

In Bayesian estimation, we consider several types of loss functions. The squared error, LINEX and entropy loss functions are widely used in literature. These loss functions were used by several authors, among them Pandey (1997), Soliman (2001), Jaheen (2005) and Guure et al. (2013).

The exponentiated Weibull distribution with two parameters (α, β) ($EW(\alpha, \beta)$) was introduced by Mudholkar and Srivastava (1993) as an extension of the Weibull family of distribution by adding an additional shape parameter. Jiang and Murthy (1999) obtained the parameter estimation of the model by graphical approaches. New statistical measures on the EW provided by Nassar and Eissa (2003).

The cumulative distribution function (CDF), probability density function (PDF), and the hazard rate function (HRF) of the EW are given, respectively, by

$$\begin{aligned}
F_1(t) &= (1 - e^{-t^\alpha})^\beta, & t > 0, \quad \alpha, \beta > 0, \\
f_1(t) &= \alpha\beta t^{\alpha-1} e^{-t^\alpha} (1 - e^{-t^\alpha})^{\beta-1}, \\
h_1(t) &= \frac{\alpha\beta t^{\alpha-1} e^{-t^\alpha} (1 - e^{-t^\alpha})^{\beta-1}}{1 - (1 - e^{-t^\alpha})^\beta}.
\end{aligned} \tag{1}$$

Bayes estimators for the two shape parameters, reliability and hazard rate functions of the EW distribution based on Type-II censored and complete samples were discussed by Nassar and Eissa (2004). Choudhury (2005) obtained a simple derivation for the moments of the EW distribution.

Several recent monographs provide more complete reviews of Markov chain Monte Carlo (MCMC) methods. The Metropolis-Hastings (MH) is one of the most important MCMC methods. It is a general method for constructing a Markov chain that draws samples from transition distributions for arbitrary posterior distributions. The Gibbs sampler may be one of the best known MCMC sampling algorithms in the Bayesian computational literature. For more details see, Chib and Greenberg (1995), Robert and Casella (1999), Liu (2001) and Glasserman (2003).

The most common censoring schemes are Type-I and Type-II censoring. The conventional Type-I and Type-II censoring schemes do not have the flexibility of allowing removal units at points other than the terminal point of the experiment. One censoring scheme known as the progressive Type-II censoring scheme, which has this advantage, see the details from Balakrishnan and Aggarwala (2000).

In addition to this introductory section, this article includes five sections. Model description and assumptions, the ML estimates (MLEs) of model parameters, acceleration factor and asymptotic variance-covariance matrix of the estimators are given in Section 2. Section 3 provides the Bayes estimates of model parameters and acceleration factor using Lindley's approximation and the MCMC procedure. Simulation studies are given in Section 4. Finally, Section 5 is devoted to the concluding remarks.

2 Model description and maximum likelihood estimation

In a constant-stress PALT, the total sample size n of test units is divided into two groups, n_1 items of group 1 are allocated to use condition and $n_2 = n - n_1$ items of group 2 are allocated to accelerated condition.

Lifetimes $T_{ji}, i = 1, \dots, n_j, j = 1, 2$ denoted two progressively Type-II censored samples from EW distribution, where $T_{1i}, i = 1, \dots, n_1$ be the items allocated at use condition and $T_{2i}, i = 1, \dots, n_2$ at accelerated condition. At the first failure time t_{j1}, R_{j1} units are randomly removed from the remaining $n_j - 1$ surviving units. At the second failure time t_{j2}, R_{j2} units are randomly removed from the remaining $n_j - R_{j1} - 1$. The test continues until the m_j th failure at which time, all remaining $R_{jm_j} = n_j - m_j - R_{j1} - R_{j2} - \dots - R_{j(m_j-1)}$ items are withdrawn. The R_{ji} are fixed prior to the study, $m_j \leq n_j$.

2.1 Assumptions

- (1) The lifetime of a test unit is $EW(\alpha, \beta)$ distribution at use condition.
- (2) At accelerated condition the hazard rate of a test unit is increased to $\lambda h_1(t)$, where λ is an acceleration factor. Therefore CDF, PDF and HRF are given, respectively, by

$$\begin{aligned}
 F_2(t) &= 1 - [1 - (1 - e^{-t^\alpha})^\beta]^\lambda, & t > 0, & (\alpha, \beta > 0, \lambda > 1), \\
 f_2(t) &= \alpha\beta\lambda t^{\alpha-1} e^{-t^\alpha} (1 - e^{-t^\alpha})^{\beta-1} [1 - (1 - e^{-t^\alpha})^\beta]^{\lambda-1}, \\
 h_2(t) &= \frac{\alpha\beta\lambda t^{\alpha-1} e^{-t^\alpha} (1 - e^{-t^\alpha})^{\beta-1}}{1 - (1 - e^{-t^\alpha})^\beta}.
 \end{aligned} \tag{2}$$

- (3) The lifetimes of test units are independent and identically distributed.

2.2 Maximum likelihood estimation

Let t_{ji, m_j, n_j} be two progressively Type-II censored samples from EW (α, β) distribution with censoring schemes $R_j = (R_{j1}, \dots, R_{jm_j}), j = 1, 2$. The likelihood function is given by

$$\begin{aligned}
l(\alpha, \beta, \lambda | \mathbf{t}) &= \prod_{j=1}^2 A_j \prod_{i=1}^{m_j} f_j(t_{ji, m_j, n_j}) [1 - F_j(t_{ji, m_j, n_j})]^{R_{ji}} \\
&= \prod_{j=1}^2 A_j (\alpha\beta)^{m_j} \lambda^{(j-1)m_j} \prod_{i=1}^{m_j} t_{ji}^{\alpha-1} e^{-t_{ji}^\alpha} (1 - e^{-t_{ji}^\alpha})^{\beta-1} [1 - (1 - e^{-t_{ji}^\alpha})^\beta]^{r_{ji}\lambda^{j-1}-1}, \\
&3
\end{aligned} \tag{2}$$

where $A_j = n_j(n_j - 1 - R_{j1})(n_j - 2 - R_{j1} - R_{j2}) \dots (n_j - (m_j - 1) - R_{j1} - \dots - R_{j(m_j-1)})$ and $r_{ji} = (R_{ji} + 1)$.

The log-likelihood function may be written as

$$\begin{aligned}
L(\alpha, \beta, \lambda | \mathbf{t}) &= \sum_{j=1}^2 \ln A_j + (m_1 + m_2) \ln(\alpha\beta) + m_2 \ln \lambda \\
&+ \sum_{j=1}^2 \sum_{i=1}^{m_j} [(\alpha - 1) \ln t_{ji} - t_{ji}^\alpha + (\beta - 1)\omega_1(\alpha; t_{ji}) + [r_{ji}\lambda^{j-1} - 1]\omega_2(\alpha, \beta; t_{ji})], \quad (4)
\end{aligned}$$

where, for simplicity, we use

$$\begin{aligned}
\omega_1(\alpha; t_{ji}) &= \ln(1 - e^{-t_{ji}^\alpha}), \\
\omega_2(\alpha, \beta; t_{ji}) &= \ln(1 - (1 - e^{-t_{ji}^\alpha})^\beta). \quad 5
\end{aligned} \tag{3}$$

For determining the MLEs of (α, β, λ) , we first have the partial derivatives of (4) with respect to α, β and λ as

$$\begin{aligned}
\frac{\partial L}{\partial \alpha} &= \frac{m_1 + m_2}{\alpha} + \sum_{j=1}^2 \sum_{i=1}^{m_j} [\ln t_{ji} - \omega_3(\alpha; t_{ji}) + (\beta - 1) \frac{\partial \omega_1(\alpha; t_{ji})}{\partial \alpha} + (r_{ji}\lambda^{j-1} - 1) \frac{\partial \omega_2(\alpha, \beta; t_{ji})}{\partial \alpha}], \\
\frac{\partial L}{\partial \beta} &= \frac{m_1 + m_2}{\beta} + \sum_{j=1}^2 \sum_{i=1}^{m_j} [\omega_1(\alpha; t_{ji}) + (r_{ji}\lambda^{j-1} - 1) \frac{\partial \omega_2(\alpha, \beta; t_{ji})}{\partial \beta}], \\
\frac{\partial L}{\partial \lambda} &= \frac{m_2}{\lambda} + \sum_{i=1}^{m_2} r_{2i} \omega_2(\alpha, \beta; t_{2i}), \quad 6
\end{aligned} \tag{4}$$

where

$$\begin{aligned}
\frac{\partial \omega_1(\alpha; t_{ji})}{\partial \alpha} &= \omega_3(\alpha; t_{ji})\omega_4(\alpha; t_{ji}), \\
\frac{\partial \omega_2(\alpha, \beta; t_{ji})}{\partial \alpha} &= -\beta\omega_3(\alpha; t_{ji})\omega_4(\alpha; t_{ji})\omega_5(\alpha, \beta; t_{ji}), \\
\frac{\partial \omega_2(\alpha, \beta; t_{ji})}{\partial \beta} &= -\omega_1(\alpha; t_{ji})\omega_5(\alpha, \beta; t_{ji}), 7
\end{aligned} \tag{5}$$

with

$$\begin{aligned}
\omega_3(\alpha; t_{ji}) &= t_{ji}^\alpha \ln t_{ji}, \\
\omega_4(\alpha; t_{ji}) &= \frac{e^{-t_{ji}^\alpha}}{1 - e^{-t_{ji}^\alpha}}, \\
\omega_5(\alpha, \beta; t_{ji}) &= \frac{(1 - e^{-t_{ji}^\alpha})^\beta}{1 - (1 - e^{-t_{ji}^\alpha})^\beta}, 8
\end{aligned} \tag{6}$$

MLEs of α, β and λ are obtained by equating the partial derivatives (6) to zero. Since the MLEs of α, β and λ can not be expressed explicitly, we apply the Newton-Raphson method to obtain $\hat{\alpha}, \hat{\beta}$ and $\hat{\lambda}$.

2.3 Observed Fisher's information matrix

The observed Fisher's information matrix is given by

$$F = -\left(\frac{\partial^2 \hat{L}(\alpha, \beta, \lambda | \mathbf{t})}{\partial K_p \partial K_q}\right), \quad p, q = 1, 2, 3, \quad K_1 = \alpha, K_2 = \beta, K_3 = \lambda,$$

where the second derivatives of $L(\alpha, \beta, \lambda | \mathbf{t})$ are given in Appendix A and the caret $\hat{\cdot}$ indicates that the derivative is calculated at $(\hat{\alpha}, \hat{\beta}, \hat{\lambda})$.

Consequently, the asymptotic variance covariance matrix for the MLEs of the parameters α, β and λ defined by inverting the observed Fisher's information matrix defined above.

Based on the asymptotic normality of the MLEs, we compute the approximate confidence intervals for α, β and λ respectively. The $100(1 - \gamma)\%$ confidence intervals for α, β and λ are given by

$$(\hat{\alpha} \pm z_{1-\gamma/2} \sqrt{\text{var}(\hat{\alpha})}), (\hat{\beta} \pm z_{1-\gamma/2} \sqrt{\text{var}(\hat{\beta})}) \text{ and } (\hat{\lambda} \pm z_{1-\gamma/2} \sqrt{\text{var}(\hat{\lambda})}),$$

where $z_{1-\gamma/2}$ is the upper $(\gamma/2)$ percentile of the standard normal distribution.

3 Bayes estimation

We suggest the joint prior density of α, β and λ as follows:

$$\pi(\alpha, \beta, \lambda) = \pi_1(\beta|\alpha)\pi_2(\alpha)\pi_3(\lambda),$$

where α and β are dependent with bivariate prior density suggested by Nassar and Eissa (2004), where $\beta|\alpha \sim \text{Gamma}(\mu, \alpha)$ and $\alpha \sim \text{Exp}(\nu)$, and λ is independent of them with non-informative prior $\pi_3(\lambda) \propto \frac{1}{\lambda}$, $\lambda > 0$. So, the joint prior of α, β and λ is written as

$$\pi(\alpha, \beta, \lambda) \propto \frac{\alpha^{-\mu}}{\lambda} \beta^{\mu-1} e^{-\left(\frac{\alpha}{\nu} + \frac{\beta}{\alpha}\right)}. \quad (9)$$

The joint posterior density function of α, β and λ can be written from (3) and (9) as

$$\begin{aligned} \pi^*(\alpha, \beta, \lambda|\mathbf{t}) &\propto l(\alpha, \beta, \lambda|\mathbf{t}) \pi(\alpha, \beta, \lambda) \\ &= \alpha^{m_1+m_2-\mu} \beta^{m_1+m_2+\mu-1} \lambda^{m_2-1} e^{-\frac{\alpha^2+\beta\nu}{\alpha\nu}} \\ &\quad \times \prod_{j=1}^2 \prod_{i=1}^{m_j} t_{ji}^{\alpha-1} e^{-t_{ji}^\alpha} (1 - e^{-t_{ji}^\alpha})^{\beta-1} [1 - (1 - e^{-t_{ji}^\alpha})^\beta]^{r_{ji}\lambda^{j-1}-1}. \end{aligned}$$

Based on squared-error loss function, the Bayes estimate of a function of the model parameters and acceleration factor is its posterior expectation. If $U = U(\theta) \equiv U(\alpha, \beta, \lambda)$ then

$$\hat{U}_{BS} = E(U(\theta)|\mathbf{t}) = \int_{\theta} U(\theta) \pi^*(\theta|\mathbf{t}) d\theta.$$

Unfortunately, we can not compute explicitly this integral in most cases. Therefore, we adopt two different procedures to approximate this integral; Lindley's approximation and MCMC procedure.

3.1 Lindley's approximation

Based on Lindley's approximation, the approximate Bayes estimates of α, β and λ for the squared error loss function are given, respectively, by

$$\begin{aligned} \tilde{\alpha}_{BS} &= \alpha + a_1 + \frac{1}{2}(\psi_1\sigma_{11} + \psi_2\sigma_{12} + \psi_3\sigma_{13}) \downarrow_{\hat{\alpha}, \hat{\beta}, \hat{\lambda}}, \\ \tilde{\beta}_{BS} &= \beta + a_2 + \frac{1}{2}(\psi_1\sigma_{21} + \psi_2\sigma_{22} + \psi_3\sigma_{23}) \downarrow_{\hat{\alpha}, \hat{\beta}, \hat{\lambda}}, \\ \tilde{\lambda}_{BS} &= \lambda + a_3 + \frac{1}{2}(\psi_1\sigma_{31} + \psi_2\sigma_{32} + \psi_3\sigma_{33}) \downarrow_{\hat{\alpha}, \hat{\beta}, \hat{\lambda}}. \end{aligned}$$

For the detailed derivations, see Appendix B.

3.2 Markov chain Monte Carlo

For generating samples from the posterior density function, we consider hybrid algorithm, which combines the MH algorithm and Gibbs sampler. The conditional posterior distributions of the parameters α, β and λ can be written, respectively, by

$$\begin{aligned}\pi^*(\alpha|\beta, \lambda, \mathbf{t}) &\propto \alpha^{m_1+m_2-\mu} e^{-\frac{\alpha^2+\beta\nu}{\alpha\nu}} \prod_{j=1}^2 \prod_{i=1}^{m_j} t_{ji}^\alpha e^{-t_{ji}^\alpha} (1 - e^{-t_{ji}^\alpha})^{\beta-1} (1 - (1 - e^{-t_{ji}^\alpha})^\beta)^{r_{ji}\lambda^{j-1}-1}, \\ \pi^*(\beta|\alpha, \lambda, \mathbf{t}) &\propto \beta^{m_1+m_2+\mu-1} e^{-\frac{\beta}{\alpha}} \prod_{j=1}^2 \prod_{i=1}^{m_j} (1 - e^{-t_{ji}^\alpha})^{\beta-1} (1 - (1 - e^{-t_{ji}^\alpha})^\beta)^{r_{ji}\lambda^{j-1}-1}, \\ \pi^*(\lambda|\alpha, \beta, \mathbf{t}) &\propto \lambda^{m_2-1} \prod_{i=1}^{m_2} (1 - (1 - e^{-t_{2i}^\alpha})^\beta)^{r_{2i}\lambda},\end{aligned}$$

where, $\pi^*(\lambda|\alpha, \beta, \mathbf{t})$ is a Gamma($m_2, -\sum_{i=1}^{m_2} r_{2i} \ln(1 - (1 - e^{-t_{2i}^\alpha})^\beta)$) distribution, so we use the Gibbs sampler to generate random numbers of λ . Moreover, the conditional posterior distribution of α and β cannot be reduced analytically to well known distributions and therefore it is not possible to sample directly by standard methods. So, we use the MH method with normal proposal distribution to generate random numbers from these distributions.

The following algorithm can be used to evaluate samples for α, β and λ .

Step 1. Choose the MLEs $\hat{\alpha}, \hat{\beta}$ and $\hat{\lambda}$, as the starting values ($\alpha^{(0)}, \beta^{(0)}, \lambda^{(0)}$) of α, β and λ .

Step 2. Set $i = 1$

Step 3. Generate $\lambda^{(i)}$ from $\pi^*(\lambda|\alpha^{(i-1)}, \beta^{(i-1)}, \mathbf{t})$.

Step 4. Generate $\alpha^{(i)}$ from $N(\alpha^{(i-1)}, \sigma_{11}^2)$.

Step 5. Generate $\beta^{(i)}$ from $N(\beta^{(i-1)}, \sigma_{22}^2)$.

Step 6. Set $i = i + 1$

Step 7. Repeat steps 3-6 N times.

Step 8. The approximate means of $U(\alpha, \beta, \lambda)$ are given, respectively, by

$$E(U|\mathbf{t}) = \frac{1}{N - M} \sum_{i=M+1}^N U(\alpha^{(i)}, \beta^{(i)}, \lambda^{(i)}),$$

where M is the burn-in period.

Step 9. $(U_{(N-M)\gamma/2}(\alpha^{(i)}, \beta^{(i)}, \lambda^{(i)}), U_{(N-M)(1-\gamma/2)}(\alpha^{(i)}, \beta^{(i)}, \lambda^{(i)}))$ yields an approximate $100(1 - \gamma)\%$ credible interval for $U(\alpha^{(i)}, \beta^{(i)}, \lambda^{(i)})$.

4 Simulation Study

A Monte Carlo simulation study was conducted in order to assess the performance of all methods of estimations (ML and the approximate Bayes estimations). The simulation study is carried out according to the following algorithm.

Step 1. We choose the values $n_1, n_2, m_1, m_2, c, \lambda$ and the censored scheme $R_{ji}, i = 1, \dots, m_j, j = 1, 2$.

Step 2. For given values of the prior parameters μ and ν generate α and β from $\text{Exp}(\nu)$ and $\text{Gamma}(\mu, \alpha)$.

Step 3. Generate two independent random samples of sizes m_1 and m_2 from $\text{Uniform}(0,1)$ distribution.

Step 4. Set

$$U_{ji} = V_{ji}^{1/(i+\sum_{q=m_j-i+1}^{m_j} R_{jq})}, \quad j = 1, 2, \quad i = 1, 2, \dots, m_j.$$

Step 5. $W_{j1}, W_{j2}, \dots, W_{jm_j}, j = 1, 2$ are two progressive Type-II censored samples of sizes $m_j, j = 1, 2$, from $U(0, 1)$ where

$$W_{ji} = 1 - \prod_{q=m_j-i+1}^{m_j} U_{jq}, \quad i = 1, \dots, m_j.$$

Step 6. Generate two random samples $t_{j1}, t_{j2}, \dots, t_{jm_j}, j = 1, 2$, from CDFs $F_1(t)$ and $F_2(t)$

$$\begin{aligned} t_{1i} &= (-\ln(1 - W_{ji}^{\frac{1}{\beta}})^{\frac{1}{\alpha}}), \quad i = 1, \dots, m_1, \\ t_{2i} &= (-\ln(1 - (1 - W_{ji}^{\frac{1}{\lambda}})^{\frac{1}{\beta}})^{\frac{1}{\alpha}}), \quad i = 1, \dots, m_2. \end{aligned}$$

Step 7. For each sample, the MLEs of acceleration factor and the parameters of EW distribution, (λ, α, β) , are obtained by solving the nonlinear Eq.(6) using the Newton-Raphson method.

Step 8. The approximate confidence intervals are constructed, through asymptotic variance covariance matrix of the estimators.

Step 9. The Bayes estimates of (λ, α, β) are computed based on squared error loss function.

Step 10. Using MH and Gibbs sampling algorithms, to generate a sequence of 5500 random samples iteratively with $N = 5500$ and $M = 500$.

Step 11. The squared deviations $(\theta^* - \theta)^2$ are computed where θ^* stands for an estimate (ML, Bayes) of the parameter θ .

Step 12. Steps 2-11 were repeated 1,000 times for different sample sizes and different censoring schemes. The mean square errors (MSE) of all the estimates is computed.

Tables 2 and 3 represent the MSE's of the ML and Bayes estimates of the population parameters α, β and acceleration factor λ in two different cases. While, Tables 4 and 5 present the 95% confidence intervals of the parameters α, β and λ .

We consider several progressive Type-II censoring schemes on data as in Table 1 with notation that $(2 * 0, 1)$ means $(0, 0, 1)$.

TABLE 1: Censoring schemes

n_1	n_2	Scheme	$R_j, \quad j = 1, 2$
m_1	m_2		
40	50	I	$R_1 = (2 * 0, 1, 0, 3, 0, 2, 0, 2, 0, 1, 2 * 0, 1, 16 * 0)$
30	40		$R_2 = (3 * 0, 1, 2 * 0, 2, 2 * 0, 2, 2 * 0, 3, 2 * 0, 1, 2 * 0, 1, 21 * 0)$
		II	$R_1 = (12 * 0, 1, 2, 3, 2, 1, 1, 12 * 0)$ $R_2 = (17 * 0, 1, 2, 3, 2, 1, 1, 17 * 0)$
		III	$R_1 = (29 * 0, 10)$ $R_2 = (39 * 0, 10)$
60	70	I	$R_1 = (2 * 0, 2, 0, 5, 0, 2, 0, 3, 0, 5, 2 * 0, 3, 26 * 0)$
40	50		$R_2 = (3 * 0, 2, 2 * 0, 5, 2 * 0, 2, 2 * 0, 3, 2 * 0, 5, 2 * 0, 3, 31 * 0)$
		II	$R_1 = (17 * 0, 2, 1, 5, 3, 2, 7, 17 * 0)$ $R_2 = (22 * 0, 2, 1, 5, 3, 2, 7, 22 * 0)$
		III	$R_1 = (39 * 0, 20)$ $R_2 = (49 * 0, 20)$
80	90	I	$R_1 = (2 * 0, 4, 0, 5, 0, 2, 0, 3, 0, 7, 2 * 0, 9, 36 * 0)$
50	60		$R_2 = (3 * 0, 4, 2 * 0, 5, 2 * 0, 2, 2 * 0, 3, 2 * 0, 7, 2 * 0, 9, 41 * 0)$
		II	$R_1 = (22 * 0, 3, 6, 2, 8, 4, 7, 22 * 0)$ $R_2 = (27 * 0, 3, 6, 2, 8, 4, 7, 27 * 0)$
		III	$R_1 = (49 * 0, 30)$ $R_2 = (59 * 0, 30)$

TABLE 2: MSE of the estimates for parameters

$$\alpha = 2.91363, \beta = 2.61485, \lambda = 1.56$$

			ML		Bayes	
					Lindley's	MCMC
n_1	n_2	Scheme				
m_1	m_2					
40	50	I	α	0.09608	0.08946	0.08968
30	40		β	0.13969	0.12842	0.12872
			λ	0.16494	0.1623	0.16223
		II	α	0.11028	0.10251	0.10338
			β	0.13166	0.12085	0.12175
			λ	0.18043	0.17751	0.17789
		III	α	0.12055	0.11351	0.11405
			β	0.15081	0.13883	0.14
			λ	0.17104	0.16873	0.16851
60	70	I	α	0.07915	0.07491	0.07553
40	50		β	0.10084	0.09464	0.09512
			λ	0.14583	0.14463	0.14503
		II	α	0.07649	0.07236	0.07291
			β	0.10073	0.09459	0.09537
			λ	0.12111	0.11895	0.11945
		III	α	0.11109	0.10676	0.10718
			β	0.09892	0.09277	0.09378
			λ	0.13794	0.13671	0.13779
80	90	I	α	0.0582	0.05576	0.05577
50	60		β	0.08022	0.07633	0.07643
			λ	0.0962	0.09538	0.09485
		II	α	0.06122	0.05853	0.05908
			β	0.07974	0.07583	0.07588
			λ	0.10501	0.10412	0.10434
		III	α	0.0798	0.07762	0.07832
			β	0.07651	0.07298	0.07383
			λ	0.10059	0.10008	0.10038

TABLE 3: MSE of the estimates for parameters

$$\alpha = 3.51913, \beta = 1.51795, \lambda = 1.54$$

			ML			
			Bayes			
			Lindley's MCMC			
n_1	n_2	Scheme				
m_1	m_2					
40	50	I	α	0.26772	0.21731	0.22157
30	40		β	0.07094	0.06704	0.06712
			λ	0.16354	0.1748	0.1744
		II	α	0.27833	0.22175	0.2277
			β	0.07265	0.06766	0.06783
			λ	0.1678	0.17742	0.17602
		III	α	0.38876	0.31002	0.3152
			β	0.06923	0.06212	0.06222
			λ	0.1624	0.17002	0.16907
60	70	I	α	0.17404	0.14759	0.15116
40	50		β	0.04734	0.04503	0.045
			λ	0.11643	0.12308	0.1232
		II	α	0.17926	0.1521	0.1554
			β	0.05077	0.04771	0.04771
			λ	0.12008	0.12594	0.12575
		III	α	0.31782	0.27029	0.27377
			β	0.0544	0.04934	0.04962
			λ	0.12723	0.13235	0.13236
80	90	I	α	0.12219	0.10833	0.10969
60	70		β	0.03456	0.03355	0.03365
			λ	0.08652	0.09019	0.0902
		II	α	0.15898	0.13899	0.13996
			β	0.037	0.3515	0.03527
			λ	0.09065	0.0939	0.09384
		III	α	0.31309	0.27505	0.28105
			β	0.05012	0.04605	0.04654
			λ	0.10882	0.11211	0.11261

TABLE 4: Confidence bounds of the estimates (in 95 %) for
 $\alpha = 2.91363, \beta = 2.61485, \lambda = 1.56$

				approximate interval	credible interval
n_1	n_2	Scheme		(L,U)	(L,U)
m_1	m_2				
40	50	I	α	(2.42914,3.56727)	(2.45555,3.56322)
30	40		β	(1.92734,3.36041)	(1.98595,3.39017)
			λ	(0.84663,2.37104)	(0.99169,2.53311)
		II	α	(2.42067,3.59626)	(2.4505,3.59577)
			β	(1.91509,3.32584)	(1.97231,3.35346)
			λ	(0.84876,2.38166)	(0.99397,2.54389)
		III	α	(2.36833,3.63959)	(2.41476,3.65651)
			β	(1.93459,3.3572)	(1.98604,3.38101)
			λ	(0.85237,2.41005)	(0.99855,2.56936)
60	70	I	α	(2.49309,3.48263)	(2.5116,3.48074)
40	50		β	(2.00992,3.22434)	(2.05513,3.25065)
			λ	(0.9375,2.27111)	(1.04934,2.39522)
		II	α	(2.47114,3.50648)	(2.49333,3.50547)
			β	(2.01149,3.20584)	(2.05565,3.23178)
			λ	(0.91842,2.22797)	(1.02893,2.35049)
		III	α	(2.42225,3.57175)	(2.46066,3.59027)
			β	(2.0436,3.24956)	(2.08138,3.26725)
			λ	(0.94556,2.31234)	(1.05878,2.43386)
80	90	I	α	(2.5326,3.41207)	(2.54671,3.41255)
50	60		β	(2.07706,3.15737)	(2.1141,3.18075)
			λ	(0.98829,2.17199)	(1.07816,2.2707)
		II	α	(2.50628,3.43489)	(2.52523,3.43551)
			β	(2.088,3.1472)	(2.12148,3.16645)
			λ	(0.99276,2.18866)	(1.08322,2.28708)
		III	α	(2.46144,3.51901)	(2.49502,3.53592)
			β	(2.09437,3.15597)	(2.12531,3.17224)
			λ	(0.99477,2.20548)	(1.08571,2.30263)

TABLE 5: Confidence bounds of the estimates (in 95 %) for
 $\alpha = 3.51913, \beta = 1.51795, \lambda = 1.54$

				approximate interval	credible interval
n_1	n_2	Scheme		(L,U)	(L,U)
m_1	m_2				
40	50	I	α	(2.80596,4.54147)	(2.82148,4.48682)
30	40		β	(1.02471,2.00564)	(1.11111,2.09251)
			λ	(0.85838,2.33891)	(1.01536,2.55539)
		II	α	(2.78717,4.60404)	(2.8097,4.54947)
			β	(1.01792,2.00088)	(1.10399,2.08719)
			λ	(0.84788,2.32189)	(1.00435,2.53589)
		III	α	(2.63416,4.79423)	(2.72592,4.78266)
			β	(0.98885,2.01067)	(1.07853,2.09005)
			λ	(0.83846,2.32045)	(0.99609,2.5288)
60	70	I	α	(2.83853,4.42044)	(2.85716,4.3823)
40	50		β	(1.09197,1.94035)	(1.15874,2.00573)
			λ	(0.94251,2.24411)	(1.0636,2.40462)
		II	α	(2.8316,4.41398)	(2.85014,4.37716)
			β	(1.08475,1.9297)	(1.15096,1.99492)
			λ	(0.93673,2.22988)	(1.0575,2.38777)
		III	α	(2.68038,4.68962)	(2.77759,4.71312)
			β	(1.05018,1.94595)	(1.11922,2.00432)
			λ	(0.92761,2.23222)	(1.04937,2.38713)
80	90	I	α	(2.94807,4.26461)	(2.95458,4.23693)
50	60		β	(1.14108,1.89164)	(1.19382,1.94589)
			λ	(0.99234,2.14016)	(1.08954,2.26649)
		II	α	(2.90035,4.32394)	(2.91276,4.29198)
			β	(1.13326,1.88811)	(1.18894,1.94221)
			λ	(0.98899,2.14478)	(1.08801,2.27084)
		III	α	(2.74908,4.62927)	(2.83724,4.6602)
			β	(1.0967,1.9062)	(1.15389,1.95326)
			λ	(0.98987,2.16831)	(1.09042,2.29018)

5 Conclusion

In this article, we have considered the constant-stress PALT when the observed data comes from EW distribution under progressive Type-II censoring. We have obtained the MSE of ML and Bayes methods. Furthermore, the confidence intervals for the parameters are obtained. For the Bayesian analysis, we perform the Lindley's approximation and the MCMC algorithm. For different censoring schemes, we list the MSE from the ML and Bayes methods and 95% confidence intervals in Tables 2,3,4 and 5.

It may be noticed, from Tables 2 and 3 that :

1. The MSE of the ML and Bayes estimates and confidence intervals decreases as the sample sizes increases.
2. The Bayes estimates have the smallest MSE as compared with their corresponding ML estimates for all the considered cases .
3. For the Bayesian estimates, the MSE's of the estimates calculated by Lindley's approximation are quite close to their corresponding MSE's of the estimates calculated by MCMC.
4. The credible intervals have the smallest length as compared with their corresponding approximate confidence intervals.

Appendix A

The second and mixed partial derivatives of α, β and λ are obtained from (6) as the following

$$\begin{aligned} \frac{\partial^2 L}{\partial \alpha^2} &= -\frac{m_1+m_2}{\alpha^2} - \sum_{j=1}^2 \sum_{i=1}^{m_j} \left[\frac{\partial \omega_3(\alpha; t_{ji})}{\partial \alpha} - (\beta - 1) \frac{\partial^2 \omega_1(\alpha; t_{ji})}{\partial \alpha^2} - [r_{ji} \lambda^{j-1} - 1] \frac{\partial^2 \omega_2(\alpha, \beta; t_{ji})}{\partial \alpha^2} \right], \\ \frac{\partial^2 L}{\partial \alpha \partial \beta} &= \sum_{j=1}^2 \sum_{i=1}^{m_j} \left[\frac{\partial \omega_1(\alpha; t_{ji})}{\partial \alpha} + [r_{ji} \lambda^{j-1} - 1] \frac{\partial^2 \omega_2(\alpha, \beta; t_{ji})}{\partial \alpha \partial \beta} \right], \\ \frac{\partial^2 L}{\partial \alpha \partial \lambda} &= \sum_{i=1}^{m_2} r_{2i} \frac{\partial \omega_2(\alpha, \beta; t_{2i})}{\partial \alpha}, \\ \frac{\partial^2 L}{\partial \beta^2} &= -\frac{m_1+m_2}{\beta^2} + \sum_{j=1}^2 \sum_{i=1}^{m_j} [r_{ji} \lambda^{j-1} - 1] \frac{\partial^2 \omega_2(\alpha, \beta; t_{ji})}{\partial \beta^2}, \\ \frac{\partial^2 L}{\partial \beta \partial \lambda} &= \sum_{i=1}^{m_2} r_{2i} \frac{\partial \omega_2(\alpha, \beta; t_{2i})}{\partial \beta}, \\ \frac{\partial^2 L}{\partial \lambda^2} &= -\frac{m_2}{\lambda^2}. \end{aligned}$$

where

$$\begin{aligned}
\frac{\partial \omega_3(\alpha; t_{ji})}{\partial \alpha} &= \omega_3(\alpha; t_{ji}) \ln t_{ji}, \\
\frac{\partial^2 \omega_1(\alpha; t_{ji})}{\partial \alpha^2} &= D_1(\alpha; t_{ji}) \frac{\partial \omega_1(\alpha; t_{ji})}{\partial \alpha}, \\
\frac{\partial^2 \omega_2(\alpha, \beta; t_{ji})}{\partial \alpha^2} &= D_2(\alpha, \beta; t_{ji}) \frac{\partial \omega_2(\alpha, \beta; t_{ji})}{\partial \alpha}, \\
\frac{\partial^2 \omega_2(\alpha, \beta; t_{ji})}{\partial \beta^2} &= D_3(\alpha, \beta; t_{ji}) \frac{\partial \omega_2(\alpha, \beta; t_{ji})}{\partial \beta}, \\
\frac{\partial^2 \omega_2(\alpha, \beta; t_{ji})}{\partial \alpha \partial \beta} &= \frac{\partial \omega_2(\alpha, \beta; t_{ji})}{\partial \alpha} \left(\frac{1}{\beta} + D_3(\alpha, \beta; t_{ji}) \right), \\
D_1(\alpha; t_{ji}) &= \ln t_{ji} - \omega_3(\alpha; t_{ji}) [1 + \omega_4(\alpha; t_{ji})], \\
D_2(\alpha, \beta; t_{ji}) &= D_1(t_{ji}) + \beta \omega_3(\alpha; t_{ji}) \omega_4(\alpha; t_{ji}) [1 + \omega_5(\alpha, \beta; t_{ji})], \\
D_3(\alpha, \beta; t_{ji}) &= \omega_1(\alpha; t_{ji}) [1 + \omega_5(\alpha, \beta; t_{ji})]
\end{aligned}$$

and $\omega_1(\alpha; t_{ji}), \omega_3(\alpha; t_{ji}), \omega_4(\alpha; t_{ji}), \omega_5(\alpha, \beta; t_{ji}), \frac{\partial \omega_1(\alpha; t_{ji})}{\partial \alpha}, \frac{\partial \omega_2(\alpha, \beta; t_{ji})}{\partial \alpha}$ and $\frac{\partial \omega_2(\alpha, \beta; t_{ji})}{\partial \beta}$ are as given in (5), (7) and (8).

Appendix B

In a three parameter case $U = U(\alpha, \beta, \lambda)$, Lindley's approximation form reduces to

$$\begin{aligned}
E(U(\alpha, \beta, \lambda) | \mathbf{t}) &= U + [U_1 a_1 + U_2 a_2 + U_3 a_3 + a_4 + a_5] \\
&+ \frac{1}{2} [\psi_1 (U_1 \sigma_{11} + U_2 \sigma_{12} + U_3 \sigma_{13}) + \psi_2 (U_1 \sigma_{21} + U_2 \sigma_{22} + U_3 \sigma_{23}) \\
&+ \psi_3 (U_1 \sigma_{31} + U_2 \sigma_{32} + U_3 \sigma_{33})]. \tag{B.1}
\end{aligned}$$

where $U_i = \frac{\partial U}{\partial \xi_i}$, σ_{ij} is the element (i, j) in the variance-covariance matrix $(-L_{ij})$, $i, j = 1, 2, 3$, and

$$\begin{aligned}
a_i &= \rho_1 \sigma_{i1} + \rho_2 \sigma_{i2} + \rho_3 \sigma_{i3}, \quad i = 1, 2, 3 \\
a_4 &= U_{12} \sigma_{12} + U_{13} \sigma_{13} + U_{23} \sigma_{23}, \\
a_5 &= \frac{1}{2} (U_{11} \sigma_{11} + U_{22} \sigma_{22} + U_{33} \sigma_{33}), \\
\psi_1 &= \sigma_{11} L_{111} + 2(\sigma_{12} L_{121} + \sigma_{13} L_{131} + \sigma_{23} L_{231}) + \sigma_{22} L_{221} + \sigma_{33} L_{331}, \\
\psi_2 &= \sigma_{11} L_{112} + 2(\sigma_{12} L_{122} + \sigma_{13} L_{132} + \sigma_{23} L_{232}) + \sigma_{22} L_{222} + \sigma_{33} L_{332}, \\
\psi_3 &= \sigma_{11} L_{113} + 2(\sigma_{12} L_{123} + \sigma_{13} L_{133} + \sigma_{23} L_{233}) + \sigma_{22} L_{223} + \sigma_{33} L_{333}, \quad \text{where } \rho_i = \frac{\partial \rho}{\partial \xi_i}, U_{ij} = \\
&\frac{\partial^2 U}{\partial \xi_i \partial \xi_j}, L_{ijk} = \frac{\partial^3 L}{\partial \xi_i \partial \xi_j \partial \xi_k}.
\end{aligned}$$

To apply Lindley's approximation form (B.1), we first observe

$$\begin{aligned}\rho(\alpha, \beta, \lambda) &= \ln \pi(\alpha, \beta, \lambda) \\ &\propto -\mu \ln \alpha + (\mu - 1) \ln \beta - \ln \lambda - \left(\frac{\alpha^2 + \beta\nu}{\alpha\nu}\right).\end{aligned}$$

therefore,

$$\rho_1 = \frac{\partial \rho}{\partial \alpha} = -\frac{\mu}{\alpha} - \frac{1}{\nu} + \frac{\beta}{\alpha^2}, \quad \rho_2 = \frac{\partial \rho}{\partial \beta} = \frac{\mu - 1}{\beta} - \frac{1}{\alpha}, \quad \rho_3 = \frac{\partial \rho}{\partial \lambda} = -\frac{1}{\lambda}.$$

and

$$\begin{aligned}L_{111} &= \frac{\partial^3 L}{\partial \alpha^3} = \frac{2(m_1+m_2)}{\alpha^3} - \sum_{j=1}^2 \sum_{i=1}^{m_j} \left[\frac{\partial^2 \omega_3(\alpha; t_{ji})}{\partial \alpha^2} - (\beta - 1) \frac{\partial^3 \omega_1(\alpha; t_{ji})}{\partial \alpha^3} \right. \\ &\quad \left. - [r_{ji} \lambda^{j-1} - 1] \frac{\partial^3 \omega_2(\alpha, \beta; t_{ji})}{\partial \alpha^3} \right], \\ L_{112} &= L_{121} = \frac{\partial^3 L}{\partial \alpha^2 \partial \beta} = \sum_{j=1}^2 \sum_{i=1}^{m_j} \left[\frac{\partial^2 \omega_1(\alpha; t_{ji})}{\partial \alpha^2} + [r_{ji} \lambda^{j-1} - 1] \frac{\partial^3 \omega_2(\alpha, \beta; t_{ji})}{\partial \alpha^2 \partial \beta} \right], \\ L_{113} &= L_{131} = \frac{\partial^3 L}{\partial \alpha^2 \partial \lambda} = \sum_{i=1}^{m_2} r_{2i} \frac{\partial^2 \omega_2(\alpha, \beta; t_{2i})}{\partial \alpha^2}, \\ L_{123} &= L_{132} = L_{231} = \frac{\partial^3 L}{\partial \alpha \partial \beta \partial \lambda} = \sum_{i=1}^{m_2} r_{2i} \frac{\partial^2 \omega_2(\alpha, \beta; t_{2i})}{\partial \alpha \partial \beta}, \\ L_{221} &= L_{122} = \frac{\partial^3 L}{\partial \beta^2 \partial \alpha} = \sum_{j=1}^2 \sum_{i=1}^{m_j} [r_{ji} \lambda^{j-1} - 1] \frac{\partial^3 \omega_2(\alpha, \beta; t_{ji})}{\partial \beta^2 \partial \alpha}, \\ L_{222} &= \frac{\partial^3 L}{\partial \beta^3} = \frac{2(m_1+m_2)}{\beta^3} + \sum_{j=1}^2 \sum_{i=1}^{m_j} [r_{ji} \lambda^{j-1} - 1] \frac{\partial^3 \omega_2(\alpha, \beta; t_{ji})}{\partial \beta^3}, \\ L_{223} &= L_{232} = \frac{\partial^3 L}{\partial \beta^2 \partial \lambda} = \sum_{i=1}^{m_2} r_{2i} \frac{\partial^2 \omega_2(\alpha, \beta; t_{2i})}{\partial \beta^2}, \\ L_{331} &= L_{133} = \frac{\partial^3 L}{\partial \lambda^2 \partial \alpha} = 0, \\ L_{332} &= L_{233} = \frac{\partial^3 L}{\partial \lambda^2 \partial \beta} = 0, \\ L_{333} &= \frac{\partial^3 L}{\partial \lambda^3} = \frac{2m_2}{\lambda^3}.\end{aligned}$$

where

$$\begin{aligned}
\frac{\partial^2 \omega_3(\alpha; t_{ji})}{\partial \alpha^2} &= \frac{\partial \omega_3(\alpha; t_{ji})}{\partial \alpha} \ln t_{ji} \\
\frac{\partial^3 \omega_1(\alpha; t_{ji})}{\partial \alpha^3} &= \left[\frac{\partial^2 \omega_1(\alpha; t_{ji})}{\partial \alpha^2} - \left(\frac{\partial \omega_1(\alpha; t_{ji})}{\partial \alpha} \right)^2 \right] D_1(\alpha; t_{ji}) \\
&\quad - \frac{\partial \omega_1(\alpha; t_{ji})}{\partial \alpha} \frac{\partial \omega_3(\alpha; t_{ji})}{\partial \alpha}, \\
\frac{\partial^3 \omega_2(\alpha, \beta; t_{ji})}{\partial \alpha^3} &= \left[\frac{\partial^2 \omega_2(\alpha, \beta; t_{ji})}{\partial \alpha^2} - \left(\frac{\partial \omega_2(\alpha, \beta; t_{ji})}{\partial \alpha} \right)^2 \right] D_2(\alpha, \beta; t_{ji}) \\
&\quad - \frac{\partial \omega_2(\alpha, \beta; t_{ji})}{\partial \alpha} \frac{\partial \omega_3(\alpha; t_{ji})}{\partial \alpha} + (\beta - 1) D_1(\alpha; t_{ji}) \\
&\quad \times \frac{\partial \omega_1(\alpha; t_{ji})}{\partial \alpha} \frac{\partial \omega_2(\alpha, \beta; t_{ji})}{\partial \alpha}, \\
\frac{\partial^3 \omega_2(\alpha, \beta; t_{ji})}{\partial \alpha^2 \partial \beta} &= \left[D_2(\alpha, \beta; t_{ji}) - \frac{\partial \omega_2(\alpha, \beta; t_{ji})}{\partial \alpha} \right] \frac{\partial^2 \omega_2(\alpha, \beta; t_{ji})}{\partial \alpha \partial \beta} \\
&\quad + \frac{\partial \omega_1(\alpha; t_{ji})}{\partial \alpha} \frac{\partial \omega_2(\alpha, \beta; t_{ji})}{\partial \alpha}, \\
\frac{\partial^3 \omega_2(\alpha, \beta; t_{ji})}{\partial \beta^2 \partial \alpha} &= D_3(\alpha, \beta; t_{ji}) \left[\frac{\partial^2 \omega_2(\alpha, \beta; t_{ji})}{\partial \beta \partial \alpha} + \omega_5(\alpha, \beta; t_{ji}) \right. \\
&\quad \left. \times \left[\omega_1(\alpha; t_{ji}) \frac{\partial \omega_2(\alpha, \beta; t_{ji})}{\partial \alpha} - \frac{\partial \omega_1(\alpha; t_{ji})}{\partial \alpha} \right] \right], \\
\frac{\partial^3 \omega_2(\alpha, \beta; t_{ji})}{\partial \beta^3} &= D_3(\alpha, \beta; t_{ji}) \left[\omega_1(\alpha; t_{ji}) \omega_5(\alpha, \beta; t_{ji}) \right. \\
&\quad \left. + D_3(\alpha; t_{ji}) \right] \frac{\partial \omega_2(\alpha, \beta; t_{ji})}{\partial \beta}
\end{aligned}$$

Furthermore,

$$\begin{aligned}
\psi_1 &= \left[\frac{2(m_1 + m_2)}{\alpha^3} - \sum_{j=1}^2 \sum_{i=1}^{m_j} \left[\frac{\partial^2 \omega_3(\alpha; t_{ji})}{\partial \alpha^2} - (\beta - 1) \frac{\partial^3 \omega_1(\alpha; t_{ji})}{\partial \alpha^3} \right. \right. \\
&\quad \left. \left. - [r_{ji} \lambda^{j-1} - 1] \frac{\partial^3 \omega_2(\alpha, \beta; t_{ji})}{\partial \alpha^3} \right] \right] \sigma_{11} + 2 \left[\sum_{j=1}^2 \sum_{i=1}^{m_j} \left[\frac{\partial^2 \omega_1(\alpha; t_{ji})}{\partial \alpha^2} \right. \right. \\
&\quad \left. \left. + [r_{ji} \lambda^{j-1} - 1] \frac{\partial^3 \omega_2(\alpha, \beta; t_{ji})}{\partial \alpha^2 \partial \beta} \right] \right] \sigma_{12} + 2 \left[\sum_{i=1}^{m_2} r_{2i} \frac{\partial^2 \omega_2(\alpha, \beta; t_{2i})}{\partial \alpha^2} \right] \sigma_{13} \\
&\quad + 2 \left[\sum_{i=1}^{m_2} r_{2i} \frac{\partial^2 \omega_2(\alpha, \beta; t_{2i})}{\partial \alpha \partial \beta} \right] \sigma_{23} + \left[\sum_{j=1}^2 \sum_{i=1}^{m_j} [r_{ji} \lambda^{j-1} - 1] \frac{\partial^3 \omega_2(\alpha, \beta; t_{ji})}{\partial \beta^2 \partial \alpha} \right] \sigma_{22},
\end{aligned}$$

$$\begin{aligned}
\psi_2 &= \left[\sum_{j=1}^2 \sum_{i=1}^{m_j} \left[\frac{\partial^2 \omega_1(\alpha; t_{ji})}{\partial \alpha^2} + [r_{ji} \lambda^{j-1} - 1] \frac{\partial^3 \omega_2(\alpha, \beta; t_{ji})}{\partial \alpha^2 \partial \beta} \right] \right] \sigma_{11} \\
&+ 2 \left[\sum_{j=1}^2 \sum_{i=1}^{m_j} [r_{ji} \lambda^{j-1} - 1] \frac{\partial^3 \omega_2(\alpha, \beta; t_{ji})}{\partial \beta^2 \partial \alpha} \right] \sigma_{12} \\
&+ 2 \left[\sum_{i=1}^{m_2} r_{2i} \frac{\partial^2 \omega_2(\alpha, \beta; t_{2i})}{\partial \alpha \partial \beta} \right] \sigma_{13} + 2 \left[\sum_{i=1}^{m_2} r_{2i} \frac{\partial^2 \omega_2(\alpha, \beta; t_{2i})}{\partial \beta^2} \right] \sigma_{23} \\
&+ \left[\frac{2(m_1 + m_2)}{\beta^3} + \sum_{j=1}^2 \sum_{i=1}^{m_j} [r_{ji} \lambda^{j-1} - 1] \frac{\partial^3 \omega_2(\alpha, \beta; t_{ji})}{\partial \beta^3} \right] \sigma_{22}, \\
\psi_3 &= \left[\sum_{i=1}^{m_2} r_{2i} \frac{\partial^2 \omega_2(\alpha, \beta; t_{2i})}{\partial \alpha^2} \right] \sigma_{11} + 2 \left[\sum_{i=1}^{m_2} r_{2i} \frac{\partial^2 \omega_2(\alpha, \beta; t_{2i})}{\partial \alpha \partial \beta} \right] \sigma_{12} \\
&+ \left[\sum_{i=1}^{m_2} r_{2i} \frac{\partial^2 \omega_2(\alpha, \beta; t_{2i})}{\partial \beta^2} \right] \sigma_{22} + \frac{2m_2}{\lambda^3} \sigma_{33}.
\end{aligned}$$

Acknowledgments

The authors thank the referees for their helpful remarks and suggestions that improved the original manuscript.

References

- [1] Abd-Elfattah, A. M., Hassan, A. S. and Nassr, S. G. (2008). Estimation in step-stress partially accelerated life tests for the Burr Type-XII distribution using Type-I censoring. *Statistical Methodology*, Vol. 5, 502–514.
- [2] Abdel-Hamid, A. H. (2009). Constant-partially accelerated life tests for Burr Type-XII distribution with progressive Type-II censoring. *Comput. Statist. Data Anal.*, Vol. 53, 2511- 2523.
- [3] AL-Hussaini, E. K. and Abdel-Hamid, A. H. (2004). Bayesian estimation of the parameters, reliability and hazard rate functions of mixtures under accelerated life tests. *Commun. Stat. - Simul. Comput.* Vol. 33 (4), 963-982.
- [4] AL-Hussaini, E. K. and Abdel-Hamid, A. H. (2006). Accelerated life tests under finite mixture models. *J. Stat. Comput. Simul.* Vol. 76 (8), 673- 690.

- [5] Balakrishnan, N. and Aggarwale, R. (2000). Progressive censoring: Theory, Methods and Applications. Birkhäuser, Boston.
- [6] Chib, S. and Greenberg, E. (1995). Understanding the Metropolis-Hastings algorithm. *Am. Statistn*, Vol. 49, 327-335.
- [7] Choudhury, A. (2005). A simple derivation of moments of the exponentiated Weibull distribution. *Metrika*, Vol. 62, 17–22.
- [8] Glasserman, P. (2003). Monte Carlo Methods in Financial Engineering. Springer.
- [9] Guure, C. B., Ibrahim, N. A., Adam, M. B., Bosomprah, S. and Mohammed, A. (2013). Bayesian parameter and reliability estimate of Weibull failure time distribution. *Bull. Malays. Math. Sci. Soc.* (2), (to appear)
- [10] Jaheen, Z. F. (2005). Estimation based on generalized order statistics from the Burr model. *Com.m. Statist. Theory Meth.*, Vol. 34, 785–794.
- [11] Jiang, R. and Murthy, D. N. P. (1999). The exponentiated Weibull family: a graphical approach. *IEEE Trans. Reliab.* Vol. 48(1), 68–72.
- [12] Lindley, D. V. (1980). Approximate Bayesian method. *Trabajos de Estadística*, Vol. 31, 223–237.
- [13] Liu, J. (2001). Monte Carlo Strategies in Scientific Computing. Springer.
- [14] Mudholkar, G. S., Srivastava, D. K. (1993). Exponentiated Weibull family for analyzing bathtub failure-rate data. *IEEE Trans. Reliab.* Vol. 42, 99–302.
- [15] Nassar, M. M. and Eissa, F. H. (2003). On the exponentiated Weibull distribution. *Commun. Statist. Theory Meth.*, Vol. 32, 1317–1336.
- [16] Nassar, M. M. and Eissa, F. H. (2004). Bayesian estimation for the exponentiated Weibull model. *Commun. Statist. Theory Meth.*, Vol. 33, 2343–2236.
- [17] Nelson, W. (1990). Accelerated Testing: Statistical Models, Test Plans and Data Analysis. Wiley, New York.
- [18] Pandey, B. N. (1997). Estimator of the scale parameter of the exponential distribution using LINEX loss function. *Commun. Statist. Theory Meth.* Vol. 26(9), 2191–2202.

- [19] Robert, C. and Casella, G. (1999). Monte Carlo Statistical Methods. Springer.
- [20] Soliman, A. A. (2001). LINEX and quadratic approximate Bayes estimators applied to the Pareto model. *Commun. Statist. Simulat. Comput.* Vol. 30(1), 63–77.