

THE ZARISKI TOPOLOGY ON THE SECOND SPECTRUM OF A MODULE (II)

H. ANSARI-TOROGHY, S. KEYVANI AND F. FARSHADIFAR

ABSTRACT. Let R be a commutative ring and M an R -module. Let $Spec^s(M)$ be the collection of all second submodules of M . In this article, we consider a new topology on $Spec^s(M)$, called the second classical Zariski topology, and investigate the interplay between the module theoretic properties of M and the topological properties of $Spec^s(M)$. Moreover, we study $Spec^s(M)$ from point of view of spectral space.

1. INTRODUCTION

Throughout this paper, R will denote a commutative ring with identity. If N is a subset of an R -module M we write $N \leq M$ to indicate that N is a submodule of M .

Let M be an R -module. A proper submodule N of M is said to be *prime* if for any $r \in R$ and $m \in M$ with $rm \in N$, we have $m \in N$ or $r \in (N :_R M)$. The *prime spectrum* of M denoted by $Spec(M)$ is the set of all prime submodules of M .

A non-zero submodule N of M is said to be *second* if for each $a \in R$, the homomorphism $N \xrightarrow{a} N$ is either surjective or zero [24]. More information about this class of modules can be found in [5], [6], [7], [14], and [15].

The concept of prime submodule has led to the development of topologies on the spectrum of modules. A brief history of this development can be seen in [20, Page 808]. More information concerning the spectrum of rings, posets, and modules can be found in [1], [2], [10], [11], [12], [13], [17], [18], and [23].

Let $Spec^s(M)$ be the set of all second submodules of M . For any submodule N of M , $V^{s*}(N)$ is defined to be the set of all second submodules of M contained in N . Of course, $V^{s*}(0)$ is just the empty set and $V^{s*}(M)$ is $Spec^s(M)$. It is easy to see that for any family of submodules N_i ($i \in I$) of M , $\bigcap_{i \in I} V^{s*}(N_i) = V^{s*}(\bigcap_{i \in I} N_i)$. Thus if $\zeta^{s*}(M)$ denotes the collection of all subsets $V^{s*}(N)$ of $Spec^s(M)$, then $\zeta^{s*}(M)$ contains the empty set and $Spec^s(M)$, and $\zeta^{s*}(M)$ is closed under arbitrary intersections. In general $\zeta^{s*}(M)$ is not closed under finite unions. A module M is called a *cotop module* if $\zeta^{s*}(M)$ is closed under finite unions. In this case, $\zeta^{s*}(M)$ is called the *quasi Zariski topology* [7].

Now let N be a submodule of M . We define $W^s(N) = Spec^s(M) - V^{s*}(N)$ and put $\Omega^s(M) = \{W^s(N) : N \leq M\}$. Let $\eta^s(M)$ be the topology on $Spec^s(M)$ by the sub-basis $\Omega^s(M)$. In fact $\eta^s(M)$ is the collection U of all unions of finite intersections of elements of $\Omega^s(M)$ [21]. We call this topology the *second classical Zariski topology* of M . It is clear that if M is a cotop module, then its related topology, as

2000 *Mathematics Subject Classification.* 13C13, 13C99.

Key words and phrases. Second submodule, maximal second submodule, second spectrum, spectral space, and Zariski topology.

it was mentioned in the above paragraph, coincide with the second classical Zariski topology. In this paper, we obtain some new results analogous to those for classical Zariski topology considered in [12] and [13]. In Section 2 of this paper, among other results, we investigate the relationship between the module theoretic properties of M and the topological properties of $\text{Spec}^s(M)$ (see Proposition 2.2, Corollary 2.6, and Theorems 2.7, 2.9, 2.11, 2.12). Moreover, Theorems 2.12 and 2.19 provide some useful characterizations for those modules whose second classical Zariski topologies are cofinite topologies.

Following M. Hochster [16], we say that a topological space W is a spectral space if W is homeomorphic to $\text{Spec}(S)$, with the Zariski topology, for some ring S . Spectral spaces have been characterized by M. Hochster as quasi-compact T_0 -spaces W having a quasi-compact open base closed under finite intersection and each irreducible closed subset of W has a generic point [16]. In Section 3, we follow the Hochster's characterization and consider $\text{Spec}^s(M)$ from point view of spectral spaces. We prove that if M has dcc on socle submodules, then for each $n \in \mathbb{N}$, and submodule N_i ($1 \leq i \leq n$) of M , $W^s(N_1) \cap W^s(N_2) \cap \dots \cap W^s(N_n)$. In particular $\text{Spec}^s(M)$ is quasi compact with second classical Zariski topology (see Theorem 3.15). It is shown that if M is a finite R -module, then $\text{Spec}^s(M)$ is a spectral space (see Theorem 3.10). Also, it is proved that if M is an R -module such that M has dcc on socle submodules, then $\text{Spec}^s(M)$ is a spectral space (see Theorem 3.17). Moreover, we show that if R is a commutative Noetherian ring and M is a comultiplication R -module with finite length, then $\text{Spec}^s(M)$ is spectral (see Proposition 3.9).

In the rest of this paper, $X^s(M)$ will denote $\text{Spec}^s(M)$.

2. TOPOLOGY ON $\text{Spec}^s(M)$

We will consider the cases that an R -module M satisfies the following condition:

(**) For any submodules $N_1, N_2 \leq M$, $V^{s*}(N_1) = V^{s*}(N_2) \Rightarrow N_1 = N_2$.

Example 2.1. Every vector space satisfies the (**) condition.

We recall that for an R -module M , the *second socle* of M is defined to be the sum of all second submodules of M and denoted by $\text{soc}(M)$ (or $\text{sec}(M)$). If M has no second submodule, then $\text{soc}(M)$ is defined to be 0. Also, a submodule N of M is said to be a *socle submodule* of M if $\text{soc}(N) = N$ [5].

Proposition 2.2. Let M be a nonzero R -module. Then the following statements are equivalent.

- (a) M satisfies the (**) condition.
- (b) Every nonzero submodule of M is a socle submodule of M .

Proof. (a) \Rightarrow (b). Let S_1 be a nonzero submodule of M . We claim that $V^{s*}(S_1) \neq \emptyset$. Otherwise, $V^{s*}(S_1) = \emptyset = V^{s*}(0)$ implies that $S_1 = 0$, a contradiction. Now let $S_2 = \sum_{S \in V^{s*}(S_1)} S$. Clearly, $V^{s*}(S_1) = V^{s*}(S_2)$. So by hypothesis, $S_1 = S_2$. Hence S_1 is a socle submodule of M .

(b) \Rightarrow (a). This follows from the fact that every submodule N of M is a sum of second submodules if and only if $N = \sum_{S \in V^{s*}(N)} S$. \square

Corollary 2.3. Every semisimple R -module M satisfies the (**) condition.

Proof. This follows from the fact that every minimal submodule of M is a second submodule of M by [24, 1.6] \square

Let M be an R -module. A proper submodule P of M is said to be a *semiprime submodule* if $I^2N \subseteq P$, where $N \leq M$ and I is an ideal of R , then $IN \subseteq P$. M is said to be *fully semiprime* if each proper submodule of M is semiprime.

A nonzero submodule N of M is said to be *semisecund* if $rN = r^2N$ for each $r \in R$ [4].

Remark 2.4. Let M be an R -module and N be a submodule of M . Let denote the set of all prime submodules of M by $Spec_R(M)$. Define $V(N) = \{P \in Spec_R(M) : P \supseteq N\}$. An R -module M is said to satisfy the $(*)$ condition provided that if N_1, N_2 are submodules of M with $V(N_1) = V(N_2)$, then $N_1 = N_2$ [12].

Definition 2.5. We call an R -module M *fully semisecund* if each nonzero submodule of M is semisecund.

Corollary 2.6. Let M be an R -module. Then the following statements are equivalent.

- (a) M satisfies the $(*)$ condition.
- (b) M is a fully semiprime module.
- (c) M is a cosemisimple module.
- (d) M satisfies the $(**)$ condition.
- (e) M is a fully semisecund module.

Proof. (a) \Leftrightarrow (b) \Leftrightarrow (c). By [12, 2.6].

(d) \Leftrightarrow (e). By Proposition 2.2.

(b) \Leftrightarrow (e). By [4, 4.8]. \square

The *second submodule dimension* of an R -module M , denoted by $S.dim M$, is defined to be the supremum of the length of chains of second submodules of M if $X^s(M) \neq \emptyset$ and -1 otherwise [8].

Let X be a topological space and let x and y be points in X . We say that x and y can be *separated* if each lies in an open set which does not contain the other point. X is a T_1 -space if any two distinct points in X can be separated. A topological space X is a T_1 -space if and only if all points of X are closed in X .

Theorem 2.7. Let M be an R -module. Then $X^s(M)$ is a T_1 -space if and only if $S.dim(M) \leq 0$.

Proof. First assume that $X^s(M)$ is a T_1 -space. If $X^s(M) = \emptyset$, then $dim(M) = -1$. Also, if $X^s(M)$ has one element, clearly $S.dim M = 0$. So we can assume that $Spec^s(M)$ has more than two elements. We show that every element of $X^s(M)$ is minimal. To show this, let $S_1 \subseteq S_2$, where $S_1, S_2 \in X^s(M)$. Now $\{S_2\} = \bigcap_{i \in I} (\bigcup_{j=1}^{n_i} V^{s*}(N_{ij}))$, where $N_{ij} \leq M$ and I is an index set. So for each $i \in I$, $S_2 \in \bigcup_{j=1}^{n_i} V^{s*}(N_{ij})$ so that $S_2 \in V^{s*}(N_{it})$, $1 \leq t \leq n_i$. Thus $S_1 \in V^{s*}(N_{it})$ for $1 \leq t \leq n_i$. This implies that $S_1 \in \bigcup_{j=1}^{n_i} V^{s*}(N_{ij})$ for each i . Therefore, $S_1 \in \bigcap_{i \in I} (\bigcup_{j=1}^{n_i} V^{s*}(N_{ij})) = \{S_2\}$ as desired.

Conversely, suppose that $S.dim(M) \leq 0$. If $S.dim(M) = -1$, then $X^s(M) = \emptyset$ and hence it is a T_1 -space. Now let $S.dim(M) = 0$. Then $X^s(M) \neq \emptyset$ and for every second submodule S of M , $V^{s*}(S) = \{S\}$. Hence $X^s(M)$ is a T_1 -space. \square

Proposition 2.8. For every finitely cogenerated R -module M , the following are equivalent.

- (a) M is a semisimple module with $S.\dim(M) = 0$.
- (b) $X^s(M)$ is a T_1 -space and M satisfies the $(**)$ condition.

Proof. (a) \Rightarrow (b). By Corollary 2.3 and Theorem 2.7.

(b) \Rightarrow (a). Since $X^s(M)$ is a T_1 -space, $S.\dim(M) \leq 0$ by Theorem 2.7. As M is finitely cogenerated, every second submodule of M is a minimal submodule of M . Now the claim follows from Proposition 2.2 \square

An R -module M is said to be a *comultiplication module* if for every submodule N of M there exists an ideal I of R such that $N = (0 :_M I)$, equivalently, for each submodule N of M , we have $N = (0 :_M \text{Ann}_R(N))$ [3]. Further M is said to be a *weak comultiplication module* if M does not have any second submodule or for every second submodule S of M , $S = (0 :_M I)$ for some I is an ideal of R [6]

Theorem 2.9. Let M be a finite length module over a commutative Noetherian ring R such that $X^s(M)$ is a T_1 -space. Then M is a comultiplication module.

Proof. By [6, 3.6], it is enough to show that M is a weak comultiplication R -module. To see this, let S be a second submodule of M . Since S is finitely cogenerated and $S.\dim(M) = 0$ by Theorem 2.7, S is a minimal submodule of M . Thus $\text{Ann}_R(S) = P$ is a maximal ideal of R . Since $S \subseteq (0 :_M P)$ and $(0 :_M P)$ is a second submodule of M by [24, 1.4], $S = (0 :_M P)$, as required. \square

Theorem 2.10. Let M be an R -module. If either R is an Artinian ring or M is a Noetherian module, then M has a minimal submodule if and only if M has a second submodule. In addition if M has a second submodule, then every second submodule of M is a semisimple submodule of M .

Proof. First assume that R is an Artinian ring. Then every prime ideal of R is maximal. Let S be a second submodule of M . Then $\text{Ann}_R(S)$ is a maximal ideal of R . Thus S is a semisimple $R/\text{Ann}_R(S)$ -module. Hence S has a minimal submodule and so M has minimal submodule. Now let M be a Noetherian R -module and S be a second submodule of M . Since S is finitely generated, one can see that S is a semisimple R -module. Therefore, S has a minimal submodule and so M has a minimal submodule, as required. \square

Theorem 2.11. Let M be an R -module. If either R is an Artinian ring or M is a Noetherian R -module, then $X^s(M)$ is a T_1 -space if and only if either $X^s(M) = \emptyset$ or $X^s(M) = \text{Min}(M)$, where $\text{Min}(M)$ denotes the set of all minimal submodules of M .

Proof. By Theorem 2.7, $X^s(M)$ is a T_1 -space if and only if $S.\dim(M) \leq 0$. First suppose that $S.\dim(M) \leq 0$. If $S.\dim(M) = -1$, then $X^s(M) = \emptyset$. If $S.\dim(M) = 0$, then $X^s(M) \neq \emptyset$. Let S be a second submodule of M . Then by Theorem 2.10, S has a minimal submodule. Since $S.\dim(M) = 0$, S is a minimal submodule of M . Hence, $X^s(M) \subseteq \text{Min}(M)$. The reverse inclusion is clear. Now suppose that either $X^s(M) = \emptyset$ or $X^s(M) = \text{Min}(M)$. In the first case, $S.\dim(M) = -1$. In the second case, $S.\dim(M) = 0$ and hence $X^s(M)$ is a T_1 -space by Theorem 2.7 \square

The cofinite topology is a topology which can be defined on every set X . It has precisely the empty set and all cofinite subsets of X as open sets. As a consequence, in the cofinite topology, the only closed subsets are finite sets, or the whole of X .

Theorem 2.12. *Let M be an R -module. Then the following are equivalent.*

- (a) $X^s(M)$ is the cofinite topology.
- (b) $S.\dim(M) \leq 0$ and for every submodule N of M either $V^{s*}(N) = X^s(M)$ or $V^{s*}(N)$ is finite.

Proof. (a) \Rightarrow (b). Assume that $X^s(M)$ is the cofinite topology. Since every cofinite topology satisfies the T_1 axiom, $S.\dim(M) \leq 0$ by Theorem 2.7. Now assume that there exists a submodule N of M such that $|V^{s*}(N)| = \infty$ and $V^{s*}(N) \neq X^s(M)$. Then $W^s(N) = X^s(M) - V^{s*}(N)$ is open in $X^s(M)$ with infinite complement, a contradiction.

(b) \Rightarrow (a). Suppose that $S.\dim(M) \leq 0$ and for every submodule N of M , $V^{s*}(N) = X^s(M)$ or $|V^{s*}(N)| < \infty$. Then the complement of every open set in $X^s(M)$ is of the form $\cap_{i \in I} (\cup_{j=1}^n V^{s*}(N_{ij}))$ which is a finite set or $X^s(M)$ obviously. \square

Corollary 2.13. *Let M be an R -module such that $X^s(M)$ is finite. Then the following statements are equivalent.*

- (a) $X^s(M)$ is a Hausdorff space.
- (b) $X^s(M)$ is a T_1 -space.
- (c) $X^s(M)$ is the cofinite topology.
- (d) $X^s(M)$ is discrete.
- (e) $S.\dim(M) \leq 0$.

Lemma 2.14. *Let M be a finite length weak comultiplication module. Then $X^s(M)$ is a cofinite topology.*

Proof. The result follows from Corollary 2.13 (e) \Rightarrow (c) because M has a finite number of second submodules and every second submodule of M is minimal by [6, 3.4]. \square

Corollary 2.15. *Let R be a Noetherian ring and M be a finitely generated cocyclic R -module. Then M is Artinian and $X^s(M)$ is a cofinite topology.*

Proof. By [22], M is Artinian. Also, M is a comultiplication R -module [9, 2.5]. Now the result follows from the above Lemma. \square

The following example shows that the converse of the above corollary is not true in general.

Example 2.16. Consider $M = \mathbb{Z}_6$ as a \mathbb{Z} -module. Then M is an Artinian \mathbb{Z} -module and $X^s(M)$ is a cofinite topology but M is not a cocyclic \mathbb{Z} -module.

Theorem 2.17. *Let M be an R -module with $|X^s(M)| \geq 2$. If $X^s(M)$ is a Hausdorff space, then $S.\dim(M) = 0$ and there exist submodules N_1, N_2, \dots, N_n of M such that $V^{s*}(N_i) \neq X^s(M)$, for all i , and $X^s(M) = \cup_{i=1}^n V^{s*}(N_i)$.*

Proof. The proof is similar to that of [12, 2.26]. \square

Maximal second submodules are defined in a natural way. By Zorn's Lemma one can easily see that each second submodule of a module M is contained in a maximal second submodule of M [5]. In [5] and [6], it is shown that Artinian modules and Noetherian modules contain only finitely many maximal second submodules.

Corollary 2.18. *Let M be an Artinian R -module. Then the following statements are equivalent.*

- (a) $X^s(M)$ is a Hausdorff space.
- (b) $X^s(M)$ is a T_1 -space.
- (c) $X^s(M)$ is the cofinite topology.
- (d) $X^s(M)$ is discrete.
- (e) $X^s(M) = \text{Min}(M)$.

Proof. First we note that since M is Artinian, $X^s(M)$ is not empty.

(a) \Rightarrow (b). This is clear.

(b) \Rightarrow (c). By Theorem 2.7, $S.\dim(M) \leq 0$. Thus every second submodule of M is a maximal second submodule of M . As M is Artinian, it contains only finitely many maximal second submodules by [5, 2.6]. Therefore, $X^s(M)$ is finite. Hence, $X^s(M)$ is a cofinite topology by Corollary 2.13.

(c) \Rightarrow (d). By Theorem 2.12, $S.\dim(M) \leq 0$ so that, as we saw in the proof of (b) \Rightarrow (c), $X^s(M)$ is finite. Now the result follows from the corollary 2.13.

(d) \Rightarrow (e). Since $X^s(M)$ is a T_1 -space, by Theorem 2.7, $S.\dim(M) = 0$. Since M is Artinian, every second submodule of M contains a minimal submodule of M . Therefore, every second submodule of M is minimal so that $X^s(M) \subseteq \text{Min}(M)$. The reverse inclusion follows from the fact that every minimal submodule of M is second by [24, 1.6].

(e) \Rightarrow (a). Since M is Artinian, M contains only a finite number of maximal second submodule by [5, 2.6]. As $X^s(M) = \text{Min}(M)$, every maximal second submodule of M is minimal. Therefore, $S.\dim(M) \leq 0$ and $X^s(M)$ is finite. Now the result follows from Corollary 2.13. \square

Theorem 2.19. *Let M be a Noetherian R -module. Then the following statements are equivalent.*

- (a) $X^s(M)$ is a Hausdorff space.
- (b) $X^s(M)$ is a T_1 -space.
- (c) $X^s(M)$ is the cofinite topology.
- (d) $X^s(M)$ is discrete.
- (e) Either $X^s(M) = \emptyset$ or $X^s(M) = \text{Min}(M)$.

Proof. (a) \Rightarrow (b). This is clear.

(b) \Rightarrow (c). Let $X^s(M)$ be a T_1 -space. By Theorem 2.11, either $X^s(M) = \emptyset$ or $X^s(M) = \text{Min}(M)$. Let $X^s(M) \neq \emptyset$. Then by using [5, 2.2], every second submodule of M is a maximal second submodule of M . Since M is Noetherian, it has a finite number of maximal second submodules by [6, 2.4]. Thus, $X^s(M)$ is finite and so by Corollary 2.13, $X^s(M)$ is the cofinite topology.

(c) \Rightarrow (d). Assume that $X^s(M)$ is the cofinite topology. Then by Theorem 2.12, $S.\dim(M) \leq 0$. Now as we see in the proof of (b) \Rightarrow (c), $X^s(M)$ is finite. Therefore, $X^s(M)$ is discrete by Corollary 2.13.

(d) \Rightarrow (e). This follows from Theorem 2.11

(e) \Rightarrow (a). If $X^s(M) = \text{Min}(M)$, then by using [5, 2.2], every second submodule of M is a maximal second submodule of M . As M is Noetherian, M has a finite number of maximal second submodules. Therefore, $X^s(M)$ is finite. Now by Corollary 2.13, $X^s(M)$ is a Hausdorff space. \square

Lemma 2.20. *Let M be a second module. Then $X^s(M)$ is T_1 -space if and only if M is the only second submodule of M .*

Proof. This follows from Theorem 2.7. \square

Lemma 2.21. Let $f : \acute{M} \rightarrow M$ be an R -module monomorphism, and let N be a submodule of M such that $N \subseteq f(\acute{M})$. Then $V^{s*}(N) \rightarrow V^{s*}(f^{-1}(N))$, given by $S \rightarrow f^{-1}(S)$ is a bijection. If $V^{s*}(N) = \emptyset$, then so is $V^{s*}(f^{-1}(N))$.

Proof. It is straightforward. \square

Theorem 2.22. Let $f : \acute{M} \rightarrow M$ be an R -module homomorphism. Define $\varphi : \Omega^s(M) \rightarrow \Omega^s(\acute{M})$ by $\varphi(\cap_{i \in I} (\cup_{j=1}^{n_i} V^{s*}(N_{ij}))) = \cap_{i \in I} (\cup_{j=1}^{n_i} V^{s*}(f^{-1}(N_{ij})))$, where I is an index set, $n_i \in \mathbb{N}$ and $N_{ij} \leq M$. Then φ is a well-defined map.

Proof. Suppose that $\cap_{i \in I} (\cup_{j=1}^{n_i} V^{s*}(N_{ij})) = \cap_{t \in T} (\cup_{j=1}^{n_t} V^{s*}(K_{tj}))$, where $N_{ij}, K_{tj} \leq M$ and I, T are index sets. We show that

$$\cap_{i \in I} (\cup_{j=1}^{n_i} V^{s*}(f^{-1}(N_{ij}))) = \cap_{t \in T} (\cup_{j=1}^{n_t} V^{s*}(f^{-1}(K_{tj}))) \quad (1)$$

Let $S \in \cap_{i \in I} (\cup_{j=1}^{n_i} V^{s*}(f^{-1}(N_{ij})))$. Then for each $i \in I$, there exists j_i ($1 \leq j_i \leq n_i$) such that $S \in V^{s*}(f^{-1}(N_{ij_i}))$. If $S \subseteq \text{Ker}(f)$, then for each $t \in T$, and each j ($1 \leq j \leq n_t$) we have $S \subseteq f^{-1}(K_{tj})$. It follows that $S \in \cap_{t \in T} (\cup_{j=1}^{n_t} V^{s*}(f^{-1}(K_{tj})))$. Now let $S \not\subseteq \text{Ker}(f)$. Then $f(S)$ is a second submodule of M . Hence for each $i \in I$, $f(S) \in V^{s*}(N_{ij})$. Thus $f(S) \in \cap_{i \in I} (\cup_{j=1}^{n_i} V^{s*}(N_{ij}))$. Therefore, $f(S) \in \cap_{t \in T} (\cup_{j=1}^{n_t} V^{s*}(K_{tj}))$, and hence for each $t \in T$, there exists j_t ($1 \leq j_t \leq n_t$) such that $f(S) \in V^{s*}(K_{tj_t})$. It follows that for each $t \in T$, $S \subseteq f^{-1}(K_{tj_t})$ so that $S \in V^{s*}(f^{-1}(K_{tj_t}))$. Consequently, we have $S \in \cap_{t \in T} (\cup_{j=1}^{n_t} V^{s*}(f^{-1}(K_{tj})))$. Therefore,

$$\cap_{i \in I} (\cup_{j=1}^{n_i} V^{s*}(f^{-1}(N_{ij}))) \subseteq \cap_{t \in T} (\cup_{j=1}^{n_t} V^{s*}(f^{-1}(K_{tj}))).$$

By a similar argument we see that

$$\cap_{t \in T} (\cup_{j=1}^{n_t} V^{s*}(f^{-1}(K_{tj}))) \subseteq \cap_{i \in I} (\cup_{j=1}^{n_i} V^{s*}(f^{-1}(N_{ij}))).$$

Thus (1) holds. \square

Proposition 2.23. Let $f : \acute{M} \rightarrow M$ be a monomorphism such that \acute{M} is not secondless. Define $\nu : X^s(\acute{M}) \rightarrow X^s(M)$ by $\nu(S) = f(S) \in X^s(M)$ for each $S \in X^s(\acute{M})$. Then ν is a continuous map.

Proof. Clearly, ν is well-defined. Let $V = \cap_{i \in I} (\cup_{j=1}^{n_i} V^{s*}(N_{ij}))$ be a closed set in $X^s(M)$. We show that $\nu^{-1}(V) = \cap_{i \in I} (\cup_{j=1}^{n_i} V^{s*}(f^{-1}(N_{ij})))$. Let $S \in \nu^{-1}(V)$. Then $\nu(S) \in V$, so $f(S) \in \cap_{i \in I} (\cup_{j=1}^{n_i} V^{s*}(N_{ij}))$. Therefore, for each $i \in I$, there exists j_i such that $f(S) \in V^{s*}(N_{ij_i})$. But $\phi : V^{s*}(N_{ij_i} \cap \text{Im}(f)) \rightarrow V^{s*}(f^{-1}(N_{ij_i}))$, given by $\acute{S} \rightarrow f^{-1}(\acute{S})$, is a bijective map by Lemma 2.21. Hence we have $\phi(f(S)) = f^{-1}f(S) = S + \text{Ker}(f) = S \in V^{s*}(f^{-1}(N_{ij_i}))$ by Lemma 2.21. It follows that $\nu^{-1}(V) \subseteq \cap_{i \in I} (\cup_{j=1}^{n_i} V^{s*}(f^{-1}(N_{ij})))$. The reverse inclusion is proved similarly and the proof is completed. \square

Remark 2.24. Clearly, for an R -module M , $X^s(M) = X^s(\text{soc}(M))$. This fact shows that the study of Zariski topology on the second spectrum of M can be easily reduced to that of socle modules.

Lemma 2.25. Let M be an R -module and $S \in X^s(M)$. Let $V^{s*}(S)$ be endowed with the induced topology of $X^s(M)$. Then $V^{s*}(S) = X^s(S)$, where S is a second submodule of M .

Proof. Straightforward \square

Proposition 2.26. Let M be an R -module, $I_M := \text{Ann}_R(\text{Soc}_R(M))$ and $\bar{R} = R/I_M$. Then the natural map $\psi_M : X^s(M) \rightarrow \text{Spec}(R/I_M)$ given by $S \rightarrow \overline{\text{Ann}_R(S)} = \text{Ann}_R(S)/I_M$ is a continuous for the Zariski topology on $X^s(M)$ (see [7, 3.1] and [7, 3.4]) and Zariski topology in rings. Moreover, $\psi_M = \psi_{\text{soc}(M)}$.

Proof. Let A be a closed subset of $\text{Spec}(\bar{R})$. Then $A = V(\bar{I})$ for some ideal \bar{I} of \bar{R} . We claim that $\psi_M^{-1}(V(\bar{I})) = V^{s*}((0 :_M I))$. So let $S \in \psi_M^{-1}(V(\bar{I}))$. Then $\psi(S) \in V(\bar{I})$. Hence $\bar{I} \subseteq \overline{\text{Ann}_R(S)}$ and so $I \subseteq \text{Ann}_R(S)$. Thus $S \subseteq (0 :_M I)$ so that $S \in V^{s*}((0 :_M I))$. To see the reverse inclusion, let $S \in V^{s*}((0 :_M I))$. Then $S \subseteq (0 :_M I)$. Hence $I \subseteq \text{Ann}_R(S)$. Therefore, $\bar{I} \subseteq \overline{\text{Ann}_R(S)}$. Hence $\overline{\text{Ann}_R(S)} \in V(\bar{I})$. This implies that $\psi(S) \in V(\bar{I})$ and so $S \in \psi^{-1}(V(\bar{I}))$. Therefore, $\psi_M^{-1}(V(\bar{I})) = V^{s*}((0 :_M I))$ and hence ψ_M is continuous. The second assertion follows from the Remark 2.24 . \square

3. MODULES WHOSE SECOND CLASSICAL ZARISKI TOPOLOGIES ARE SPECTRAL SPACES

Let Z be a subset of a topological space W . Then the notation $cl(Z)$ will denote the closure of Z in W .

A topological space X is called irreducible if $X \neq \emptyset$ and every finite intersection of non-empty open subset of X is non-empty. A non-empty subset Y of a topology X is called *irreducible set* if the subspace Y of X is irreducible, equivalently if Y_1 and Y_2 are closed subset of X and satisfy $Y \subseteq Y_1 \cup Y_2$, then $Y \subseteq Y_1$ or $Y \subseteq Y_2$.

Let Y be a closed subset of a topological space. An element $y \in Y$ is called a *generic point* of Y if $Y = cl(\{y\})$. A generic point of an irreducible closed subset of a topological space is unique if the topological space is a T_0 space.

A spectral space is a topological space homeomorphic to the prime spectrum of a commutative ring equipped with the Zariski topology. This concept plays an important role in studying of algebraic properties of an R -module M when we have a related topology. For an example, when $X^s(M)$ is homeomorphic to $\text{Spec}(S)$, where S is a commutative ring, we can transfer some of known topological properties of $\text{Spec}(S)$ to $X^s(M)$ and then by using these properties explore some of algebraic properties of M .

Spectral spaces have been characterized by Hochster [16, p.52, Prop.4] as the topological spaces W which satisfy the following conditions:

- (a) W is a T_0 -space;
- (b) W is quasi-compact;
- (c) the quasi-compact open subsets of W are closed under finite intersection and form an open base;
- (d) each irreducible closed subset of W has a generic point.

Let M be an R -module and N be a submodule of M . In [12], among other nice results, Proposition 3.1 states that if Y is a nonempty subset of $\text{Spec}_R(M)$, then $cl(Y) = \cup_{P \in Y} V(P)$, where $V(N) = \{P \in \text{Spec}_R(M) \mid P \supseteq N\}$. Unfortunately, this result is not true in general because if we take $M = \mathbb{Z}$, where \mathbb{Z} is the ring of integers, and $Y = \text{Max}(\mathbb{Z})$, then we have $cl(Y) = \text{Max}(\mathbb{Z})$, while $cl(Y) = V(\cap_{P \in Y} P) = V(0) = \mathbb{Z}$ by [19, 5.1]. However, this theorem is true when Y is a finite set which has been used by the authors during their results in [12] and [13].

Let M be an R -module and Y a subset of $X^s(M)$. We will denote $\sum_{S \in Y} S$ by $T(Y)$.

Proposition 3.1. Let M be an R -module.

- (a) If Y is a finite subset of $X^s(M)$, then $cl(Y) = \cup_{S \in Y} V^{s*}(S)$.
- (b) if Y is a closed subset of $X^s(M)$, we have $Y = \cup_{S \in Y} V^{s*}(S)$.
- (c) If M is a cotop R -module and if Y is a subset of $X^s(M)$, then $cl(Y) = V^{s*}(T(Y)) = V^{s*}(\sum_{S \in Y} S)$.

Proof. (a) Clearly, $Y \subseteq \cup_{S \in Y} V^{s*}(S)$. Let K be a closed subset of $X^s(M)$ containing Y . Thus $K = \cap_{i \in I} (\cup_{j=1}^{n_i} V^{s*}(N_{ij}))$, for some $N_{ij} \subseteq M$, $i \in I$, and $n_i \in \mathbb{N}$. Let $S_1 \in \cup_{S \in Y} V^{s*}(S)$. Then there exists $S \in Y$ such that $S_1 \in V^{s*}(S)$ so that $S_1 \subseteq S$. But $S \in Y$ implies that for each $i \in I$, there exists j , $1 \leq j \leq n_i$, such that $S \subseteq N_{ij}$. Thus we have $S_1 \subseteq S \subseteq N_{ij}$. Therefore $S_1 \in K$ so that $\cup_{S \in Y} V^{s*}(S) \subseteq K$. In other words, $\cup_{S \in Y} V^{s*}(S)$ is the smallest closed subset of $X^s(M)$ containing Y , i.e., $cl(Y) = \cup_{S \in Y} V^{s*}(S)$.

(b) It is enough to show $\cup_{S \in Y} V^{s*}(S) \subseteq Y$ because the other side is clear. To see this, we note that for every element S of Y , $V^{s*}(S) = cl(\{S\}) \subseteq cl(Y) = Y$ by part (a). Hence $\cup_{S \in Y} V^{s*}(S) \subseteq Y$, as required.

(c) First we note that since M is a cotop module, each closed set is of the form of $V^{s*}(N)$ for some $N \leq M$. Clearly $Y \subseteq V^{s*}(T(Y))$. Now let $V^{s*}(N)$ be any closed subset of $X^s(M)$ containing Y . Then for each $S \in Y$, we have $S \subseteq N$ so that $T(Y) \subseteq N$. So if $S \in V^{s*}(T(Y))$, then $S \subseteq T(Y) \subseteq N$. Hence $S \in V^{s*}(N)$, i.e., $V^{s*}(T(Y)) \subseteq V^{s*}(N)$. This shows that $cl(Y) = V^{s*}(T(Y))$. This completes the proof. \square

Corollary 3.2. Let M be an R -module. Then we have the following.

- (a) $cl(\{S\}) = V^{s*}(S)$, for all $S \in Spec^s(M)$.
- (b) $S_1 \in cl(\{S\}) \Leftrightarrow S_1 \subseteq S \Leftrightarrow V^{s*}(S_1) \subseteq V^{s*}(S)$.
- (c) The set $\{S\}$ is closed in $X^s(M)$ if and only if S is a minimal second submodule of M .

Proof. Use Proposition 3.1(a). \square

Lemma 3.3. Let M be an R -module. Then for each $S \in X^s(M)$, $V^{s*}(S)$ is irreducible. In particular, $X^s(M)$ is irreducible.

Proof. The proof is straightforward. \square

We need the following evident Lemma.

Lemma 3.4. Let S be a submodule of an R -module M . Then the following are equivalent.

- (a) S is a second submodule of M .
- (b) For each $r \in R$ and submodule K of M , $rS \subseteq K$ implies that $rS = 0$ or $S \subseteq K$.

Proof. The proof is straightforward. \square

Theorem 3.5. Let M be an R -module and $Y \subseteq X^s(M)$.

- (a) If Y is irreducible, then $T(Y)$ is a second submodule.
- (b) If $T(Y)$ is a second submodule and $T(Y) \in cl(Y)$, then Y is irreducible.

Proof. (a) Assume that Y is an irreducible subset of $X^s(M)$. Then obviously $T(Y)$ is a nonzero submodule of M and $Y \subseteq V^{s*}(T(Y))$. Now let $rT(Y) \subseteq K$, whence $r \in R$ and K is a submodule of M . It is easy to see that $Y \subseteq V^{s*}((K :_M r)) \subseteq$

$V^{s*}(K) \cup V^{s*}((0 :_M r))$. Since Y is irreducible, we have $Y \subseteq V^{s*}((0 :_M r))$ or $Y \subseteq V^{s*}(K)$. If $Y \subseteq V^{s*}((0 :_M r))$, then $rS = 0$ for all $S \in Y$. Thus $rT(Y) = 0$. If $Y \subseteq V^{s*}(K)$, then $S \subseteq K$ for each $S \in Y$, so $T(Y) \subseteq K$. Hence by Lemma 3.4, $T(Y)$ is a second submodule of M .

(b) Assume that $S := T(Y)$ is a second submodule of M and $S \in cl(Y)$. It is easy to see that $cl(Y) = V^{s*}(S)$. Now let $Y \subseteq Y_1 \cup Y_2$, where Y_1, Y_2 are closed sets. Then we have $V^{s*}(S) = cl(Y) \subseteq Y_1 \cup Y_2$. Since $V^{s*}(S)$ is irreducible, $V^{s*}(S) \subseteq Y_1$ or $V^{s*}(S) \subseteq Y_2$. Hence $Y \subseteq Y_1$ or $Y \subseteq Y_2$. So Y is irreducible. \square

Corollary 3.6. Let M be an R -module and let N be a submodule of M . Then $V^{s*}(N)$ of $Spec^s(M)$ is irreducible if and only if $soc(N)$ is a second submodule of M . Consequently, $Spec^s(M)$ is irreducible if and only if $soc(M)$ is a second module.

Proof. (\Rightarrow). Let $Y := V^{s*}(N)$ be an irreducible subset of $X^s(M)$. Then we have $T(Y) = soc(N)$ so that $soc(N)$ is a second submodule of M by Theorem 3.5(a).

(\Leftarrow). Suppose $soc(N)$ is a second submodule of M . Then by Theorem 3.1(b), $Y := V^{s*}(N) = \cup_{S \in Y} V^{s*}(S)$ so that $soc(N) \in cl(Y)$. Hence $V^{s*}(N)$ is irreducible by Theorem 3.5(b). \square

We remark that any closed subset of a spectral space is spectral for the induced topology, and we note that a generic point of an irreducible closed subset Y of a topological space is unique if the topological space is a T_0 -space. The following proposition shows that for any R -module M , $X^s(M)$ is always a T_0 -space.

Lemma 3.7. Let M be an R -module. Then the following hold.

- (a) $X^s(M)$ is a T_0 -space.
- (b) Let $S \in X^s(M)$. Then S is a generic point of the irreducible closed subset $V^{s*}(S)$.

Proof. (a) This follows from Corollary 3.2 and the fact that a topological space is a T_0 -space if and only if the closures of distinct points are distinct.

(b) By Corollary 3.2. \square

By [7, 3.3], if M is a comultiplication R -module M , then the second classical Zariski topology of M and the Zariski topology of M considered in [7] coincide (note that every comultiplication module is a cotop module).

Proposition 3.8. Let R be a commutative Noetherian ring and let M be a cotop R -module with finite length. Then M is a comultiplication R -module.

Proof. This follows from [7, 2.6 (e)] and [7, 2.11]. \square

Proposition 3.9. Let M be a comultiplication R -module with finite length. Then $X^s(M)$ is a spectral space (with the second classical Zariski topology).

Proof. This follows from Lemma 3.7, [7, 3.10], and [7, 6.2] and the fact that since M is a cotop module, its assigned topology coincide with the the second classical topology. In this case, every closed set can be written as $V^{s*}(N)$ for some submodule N of M . \square

Theorem 3.10. Let M be an R -module with finite second spectrum. Then $X^s(M)$ is a spectral space (with the second Zariski topology). Consequently, for each finite module M , $Spec^s(M)$ is a spectral space.

Proof. Since $X^s(M)$ is a nonempty finite set, every subset of $X^s(M)$ is quasi-compact. So the quasi-compact open sets of $X^s(M)$ are closed under finite intersection. Further $\beta = \{W^s(N_1) \cap W^s(N_2) \cap \dots \cap W^s(N_n), N_i \leq M, n \in \mathbb{N}\}$ is a basis for $X^s(M)$ with the property that each basis element, in particular $W^s(M) = X^s(M)$, is quasi-compact. Also if $Y = \{y_1, y_2, \dots, y_n\}$ is an irreducible subset of $X^s(M)$, then we have $cl(Y) = \cup_{i=1}^n cl(\{y_i\})$. Since Y is irreducible, $cl(Y) = cl(\{y_i\})$ for some $1 \leq i \leq n$. Moreover, $X^s(M)$ is a T_0 -space by Lemma 3.7. Therefore $X^s(M)$ is a spectral space by Hochster's characterization. \square

Definition 3.11. Let M be an R -module and let $H(M)$ be the family of all subsets of $X^s(M)$ of the form $V^{s*}(N) \cap W^s(K)$, where $N, K \leq M$. $H(M)$ contains both $X^s(M)$ and \emptyset because $X^s(M) = V^{s*}(M) \cap W^s(0)$ and $\emptyset = V^{s*}(M) \cap W^s(M)$. Let $Z(M)$ be the collection of all unions of finite intersections of elements of $H(M)$. Then $Z(M)$ is a topology on $X^s(M)$ and called the *finer patch topology* or *constructible topology*. In fact, $H(M)$ is a sub-basis for the finer patch topology of M .

Theorem 3.12. *Let M be an R -module such that M satisfies descending chain condition of socle submodules. Then $X^s(M)$ with the finer patch topology is a compact space.*

Proof. Let A be a family of finer patch-open sets covering $X^s(M)$, and suppose that no finite subfamily of A covers $X^s(M)$. Since $V^{s*}(soc(M)) = V^{s*}(M) = X^s(M)$, we may use the *dcc* on socle submodules to choose a submodule N minimal with respect to the property that no finite subfamily of A covers $V^{s*}(N)$ (note that we may assume $N = soc(N)$ because $V^{s*}(N) = V^{s*}(soc(N))$). We claim that N is a second submodule of M , for if not, then there exist a submodule L of M and $r \in R$ such that $rN \subseteq L$, $rN \neq 0$, and $N \not\subseteq L$. Thus $N \cap L \subset N$ and $Soc(N \cap (0 :_M r)) \subseteq N \cap (0 :_M r) \subset N$. Hence without loss of generality, there must be a finite subfamily \acute{A} of A that covers both $V^{s*}(N \cap (0 :_M r))$ and $V^{s*}(N \cap L)$. Let $S \in V^{s*}(N)$. Since $rN \subseteq L$, we have $rS \subseteq L$. Since S is second, $S \subseteq L$ or $rS = 0$. Thus either $S \in V^{s*}(N \cap L)$ or $S \in V^{s*}(N \cap (0 :_M r))$. Therefore,

$$V^{s*}(N) \subseteq V^{s*}(N \cap (0 :_M r)) \cup V^{s*}(N \cap L).$$

Thus $V^{s*}(N)$ is covered with the finite subfamily \acute{A} , a contradiction. Hence N is a second submodule of M . Now choose $U \in A$ such that $N \in U$. Thus N must have a patch-neighborhood $\cap_{i=1}^n (W^s(K_i) \cap V^{s*}(N_i))$, for some $K_i, N_i \leq M$, $n \in \mathbb{N}$, such that

$$\cap_{i=1}^n [W^s(K_i) \cap V^{s*}(N_i)] \subseteq U.$$

We claim that for each i ($1 \leq i \leq n$),

$$N \in W^s(K_i \cap N) \cap V^{s*}(N) \subseteq W^s(K_i) \cap V^{s*}(N_i).$$

To see this, assume that $S \in W^s(K_i \cap N) \cap V^{s*}(N)$ so that $S \not\subseteq K_i \cap N$ and $S \subseteq N$. Thus $S \not\subseteq K_i$, i.e., $S \in W^s(K_i)$. On the other hand, $N \in V^{s*}(N_i)$ and $S \subseteq N$. Thus $S \in V^{s*}(N_i)$. Hence we have

$$N \in \cap_{i=1}^n [W^s(K_i \cap N) \cap V^{s*}(N)] \subseteq \cap_{i=1}^n [W^s(K_i) \cap V^{s*}(N_i)] \subseteq U.$$

Thus $[\cap_{i=1}^n W^s(\acute{K}_i)] \cap V^{s*}(N)$, where $\acute{K}_i := K_i \cap N \subset N$ is a neighborhood of N with $[\cap_{i=1}^n W^s(\acute{K}_i)] \cap V^{s*}(N) \subseteq U$. Since for each i ($1 \leq i \leq n$), $\acute{K}_i \subset N$, $V^{s*}(\acute{K}_i)$

can be covered by some finite subfamily \acute{A}_i of A . But

$$N^{s*}(N) - [\cup_{i=1}^n V^{s*}(\acute{K}_i)] = V^{s*}(N) - [\cap_{i=1}^n W^s(\acute{K}_i)]^c = [\cap_{i=1}^n [W^s(\acute{K}_i)] \cap V^{s*}(N)] \subseteq U.$$

Hence $V^{s*}(N)$ can be covered by $\acute{A}_1 \cup \acute{A}_2 \cup \dots \cup \acute{A}_n \cup \{U\}$, contrary to our choice of N . Thus there exists a finite subfamily of A which covers $X^s(M)$. Therefore, $X^s(M)$ is compact in the finer patch topology of M . \square

Proposition 3.13. Let M be R -module such that M has dcc on socle submodules. Then every irreducible closed subset of $X^s(M)$ (with second classical Zariski topology) has a generic point.

Proof. Let Y be an irreducible closed subset of $X^s(M)$. First we note that if N is a submodule of M , then $V^{s*}(N)$ and $W^s(N)$ are both open and closed in finer patch topology. Hence $V^{s*}(S)$, where $S \in Y$, and Y are also an open and a closed set in finer patch topology respectively. Since $X^s(M)$ is a compact space in finer patch topology by Theorem 3.12 and Y is closed in $X^s(M)$, we have Y is a compact space in finer patch topology. Now $Y = \cup_{S \in Y} V^{s*}(S)$ by Proposition 3.1(b) and each $V^{s*}(N)$ is open in finer patch topology. Hence there exists a finite set $Y_1 \subseteq Y$ such that $Y = \cup_{S \in Y_1} V^{s*}(S)$. Since Y is irreducible, $Y = V^{s*}(S) = cl(\{S\})$ for some $S \in Y$. Hence S is a generic point for Y , as desired. \square

Lemma 3.14. Assume ζ_1 and ζ_2 are two topologies on $X^s(M)$ such that $\zeta_1 \leq \zeta_2$. If $X^s(M)$ is quasi-compact (i.e., every open cover of it has a finite subcover) in ζ_2 , then $X^s(M)$ is also quasi-compact in ζ_1 .

Theorem 3.15. Let M be an R -module such that M has dcc on socle submodules. Then for each $n \in \mathbb{N}$, and submodules N_i ($1 \leq i \leq n$) of M , $W^s(N_1) \cap W^s(N_2) \cap \dots \cap W^s(N_n)$, is a quasi compact subset of $X^s(M)$ with the second classical Zariski topology.

Proof. Clearly, for each $n \in \mathbb{N}$ and each submodule N_i ($1 \leq i \leq n$) of M , $W^s(N_1) \cap W^s(N_2) \cap \dots \cap W^s(N_n)$ is a closed set in $X^s(M)$ with finer patch topology so that it compact in $X^s(M)$ with finer patch topology. Hence it is quasi-compact in $X^s(M)$ with the second classical Zariski topology by Lemma 3.14, as desired. \square

Corollary 3.16. Let M be an R -module such that M has dcc on socle submodules. Then quasi-compact open sets of $X^s(M)$ are closed under finite intersections.

Proof. It suffices to show that the intersection $U = U_1 \cap U_2$ of two quasi-compact open sets U_1 and U_2 of $X^s(M)$ is a quasi-compact set. Each U_i , $i = 1, 2$, is a finite union of members of the open base $\beta = \{W^s(N_1) \cap W^s(N_2) \cap \dots \cap W^s(N_n), N_i \leq M, n \in \mathbb{N}\}$, hence so is $U = \cup_{i=1}^m (\cap_{j=1}^{n_i} W^s(N_j))$. Let Ω be any open cover of U . Then Ω also covers each $\cap_{j=1}^{n_i} W^s(N_j)$ which is quasi-compact by Theorem 3.15. Hence each $\cap_{j=1}^{n_i} W^s(N_j)$ has a finite subcover of Ω and so does U . \square

Theorem 3.17. Let M be an R -module such that M has dcc on socle submodules. Then $X^s(M)$ with the second classical Zariski topology is a spectral space.

Proof. We have $X^s(M)$ is a T_0 space by Lemma 3.7. Further $\beta = \{W^s(N_1) \cap W^s(N_2) \cap \dots \cap W^s(N_n), N_i \leq M, n \in \mathbb{N}\}$ is a basis for $X^s(M)$ with the property that each basis element, in particular $W^s(M) = X^s(M)$, is quasi-compact by Theorem 3.15. Moreover, by Corollary 3.16, the quasi-compact open sets are closed

under any finite intersections. Finally, every irreducible closed set has a generic point by Proposition 3.13. Therefore, $X^s(M)$ is a spectral space by Hochster's characterization. \square

Acknowledgments. We would like to thank Dr. R. Oulyaee-Sarmazdeh for his helpful suggestions. Also we are grateful to the referee for careful reading of our manuscript and useful comments.

REFERENCES

1. J. Y. Abuhlail, *Zariski topologies for coprime and second submodules*, to appear in Algebra Colloquium.
2. J. Y. Abuhlail, *A Dual Zariski Topology for Modules*, Topology and its Applications, **158** (3) (2011) 457467.
3. H. Ansari-Toroghy and F. Farshadifar, *The dual notion of multiplication modules*, Taiwanese Journal of Mathematics **11** (4) (2007), 1189–1201.
4. H. Ansari-Toroghy and F. Farshadifar, *The dual notions of some generalizations of prime submodules*, Comm. Algebra **39** (7)(2011), 2396-2416.
5. H. Ansari-Toroghy and F. Farshadifar, *On the dual notion of prime submodules*, Algebra Colloq. **19** (Spec 1) (2012), 1109-1116.
6. H. Ansari-Toroghy and F. Farshadifar, *On the dual notion of prime submodules (II)*, Mediterr. J. Math. **9** (2) (2012), 329-338.
7. H. Ansari-Toroghy and F. Farshadifar, *The Zariski topology on the second spectrum of a module*, to appear in Algebra Colloq.
8. H. Ansari-Toroghy and F. Farshadifar, *On the dual notion of prime radicals of submodules*, Asian Eur. J. Math. **6** (2) (2013), 1350024 (11 pages).
9. H. Ansari-Toroghy and F. Farshadifar, *On the dual notion of multiplication modules*, AJSE-Mathematics, **36** (6) (2011), 925-932.
10. H. Ansari-Toroghy and S. Keyvani, *Strongly top modules*, Bull. Malays. Math. Sci. Soc. (2), **37**(1) (2014), 73-82.
11. H. Ansari-Toroghy and S. Keyvani, *On the maximal spectrum of a module and Zariski topology*, Bull. Malays. Math. Sci. Soc. (2), to appear.
12. M. Behboodi and M. R. Haddadi, *Classical Zariski Topology of Modules and Spectral Spaces I*, Inter. Electronic J. Alg. **4** (2008), 104-130.
13. M. Behboodi and M. R. Haddadi, *Classical Zariski Topology of Modules and Spectral Spaces II*, Inter. Electronic J. Alg. **4** (2008), 131-148.
14. S. Ceken, M. Alkan, and P.F. Smith, *Second modules over noncommutative rings*, Comm. Algebra, **41**(1) (2013), 83-98.
15. S. Ceken, M. Alkan, and P.F. Smith, *The dual notion of the prime radical of a module*, J. Algebra, **392** (2013), 255-265.
16. M. Hochster, *Prime ideal structure in commutative rings*, Trans. Amer. Math. Soc. **142** (1969), 43-60.
17. M. Hochster, *The minimal prime spectrum of a commutative Ring*, Canad. J. Math. **23**(1971), 749-758.
18. H.C. Li and K.P. Shum, *On a problem of spectral posets*. J. Appl. Algebra Discrete Struct. **1** (3) (2003), 203-209.
19. C.P. Lu, *The Zariski topology on the prime spectrum of a module*, Houston J. Math. **25** (3) (1999), 417–432.
20. C.P. Lu, *Modules with Noetherian spectrum*, Comm. Algebra **38** (2010), 807-828.
21. J. R. Munkres, *Topology. A First Course*. Prentice Hall, 1975.
22. P. Vámos, *The dual of the notion of "finitely generated"*, J. London Math. Soc **43** (1965), 643-646.
23. N. Van Sanh, L.P. Thas, N.F.A. Al-Mepahi, and K. P.Shum, *Zariski Topology of prime spectrum of a module*, Proceedings of the International conference 2010 held in Gajha Mada Univesity, Novenber, 2011, Yojakarta, Indonesia (Abvance in algebraic structure), World Scientific, Inc., 461-477.
24. S. Yassemi, *The dual notion of prime submodules*, Arch. Math (Brno) **37** (2001), 273–278.

DEPARTMENT OF PURE MATHEMATICS, FACULTY OF MATHEMATICAL SCIENCES, UNIVERSITY OF
GUILAN, P. O. BOX 41335-19141 RASHT, IRAN.

E-mail address: `ansari@guilan.ac.ir`

DEPARTMENT OF PURE MATHEMATICS, FACULTY OF MATHEMATICAL SCIENCES, UNIVERSITY OF
GUILAN, P. O. BOX 41335-19141 RASHT, IRAN.

E-mail address: `siamak.Keyvani@guilan.ac.ir`

UNIVERSITY OF FARHANGIAN, TEHRAN, IRAN.

E-mail address: `f.farshadifar@gmail.com`