

**CONVOLUTION, COEFFICIENT AND RADIUS
PROBLEMS OF CERTAIN UNIVALENT
FUNCTIONS**

MAISARAH BT. HAJI MOHD

UNIVERSITI SAINS MALAYSIA

2009

**CONVOLUTION, COEFFICIENT AND RADIUS
PROBLEMS OF CERTAIN UNIVALENT
FUNCTIONS**

by

MAISARAH BT. HAJI MOHD

**Thesis submitted in fulfilment
of the requirements for the Degree of
Master of Science in Mathematics**

July 2009

ACKNOWLEDGEMENT

IN THE NAME OF ALLAH S.W.T (THE AL-MIGHTY) THE GRACIOUS, THE MOST MERCIFUL.

First and foremost, I am very grateful to Allah S.W.T for giving me the strength through out my journey to complete this thesis.

I would like to express my gratitude to my supervisor, Dr. Lee See Keong, my co-supervisor, Professor Dato' Rosihan M. Ali, from the School of Mathematical Sciences, Universiti Sains Malaysia and my field supervisor, Dr.V. Ravichandran, reader at the Mathematical Department of Delhi University for their valuable guidance, assistance, encouragement and support throughout my research. Also my greatest appreciation to the whole GFT group in USM, especially, Professor K. G. Subramaniam, Dr. Adolf Stephen, Abeer Badghaish, Chandrashekar and Shamani Supramaniam. I cannot fully express my appreciation for their generosity, enthusiasms and tiredless guidance.

My sincere appreciation to the Dean, Assoc. Professor Ahmad Izani Md. Ismail and the entire staffs of the School of Mathematical Sciences, USM.

I am also very thankful to my family and friends for their understanding, helpfulness, continuous support and encouragement all the way through my studies.

CONTENTS

| | |
|--|-----|
| ACKNOWLEDGEMENT | ii |
| CONTENTS | iii |
| SYMBOLS | v |
| ABSTRAK | vii |
| ABSTRACT | ix |
| CHAPTER 1. INTRODUCTION | 1 |
| CHAPTER 2. CERTAIN SUBCLASSES OF MEROMORPHIC FUNCTIONS ASSOCIATED WITH CONVOLUTION AND DIFFERENTIAL SUBORDINATION | 4 |
| 2.1. MOTIVATION AND PRELIMINARIES | 4 |
| 2.2. DEFINITIONS | 8 |
| 2.3. INCLUSION AND CONVOLUTION THEOREM | 9 |
| CHAPTER 3. A GENERALIZED CLASS OF UNIVALENT FUNCTIONS WITH NEGATIVE COEFFICIENTS | 16 |
| 3.1. MOTIVATION AND PRELIMINARIES | 16 |
| 3.2. COEFFICIENT ESTIMATE | 19 |
| 3.3. GROWTH THEOREM | 21 |
| 3.4. COVERING THEOREM | 23 |
| 3.5. DISTORTION THEOREM | 24 |
| 3.6. CLOSURE THEOREM | 26 |
| 3.7. RADIUS PROBLEM | 30 |

| | |
|--|----|
| CHAPTER 4. RADIUS PROBLEMS FOR SOME CLASSES OF ANALYTIC FUNCTIONS | 33 |
| 4.1. MOTIVATION AND PRELIMINARIES | 33 |
| 4.2. RADIUS OF STARLIKENESS OF ORDER α | 34 |
| 4.3. RADIUS OF STRONG STARLIKENESS | 40 |
| 4.4. RADIUS OF PARABOLIC STARLIKENESS | 42 |
| CONCLUSION | 46 |
| REFERENCES | 47 |

SYMBOLS

| Symbol | Description |
|--------------------|--|
| \mathcal{A} | Class of analytic functions of the form $f(z) = z + \sum_{n=2}^{\infty} a_n z^n \quad (z \in U)$ |
| \arg | Argument |
| \mathcal{C} | Complex plane |
| $f * g$ | Convolution or Hadamard product of functions f and g |
| $\mathcal{H}(U)$ | Class of analytic functions in U |
| \Im | Imaginary part of a complex number |
| \prec | Subordinate to |
| K | Class of convex functions in U |
| $K(\alpha)$ | Class of convex functions of order α in U |
| $K(\phi)$ | $\{f \in \mathcal{A} : 1 + \frac{zf''(z)}{f'(z)} \prec \phi(z)\}$ |
| $k(z)$ | Koebe function |
| \mathcal{M} | Class of meromorphic functions of the form $f(z) = \frac{1}{z} + \sum_{n=0}^{\infty} a_n z^n \quad (z \in U^*)$ |
| $M(\alpha)$ | $\{f \in \mathcal{A} : \frac{zf'(z)}{f(z)} \prec \frac{1+(1-2\alpha)z}{1-z} \quad (-1 \leq B < A \leq 1, \alpha > 1)\}$ |
| \mathcal{R} | Set of all real numbers |
| \Re | Real part of a complex number |
| R_α | Class of prestarlike functions of order α in U |
| \mathcal{S} | Class of all univalent functions in U |
| S^* | Class of starlike functions in U |
| $S^*(\alpha)$ | Class of starlike functions of order α in U |
| $S^*(\phi)$ | $\{f \in \mathcal{A} : \frac{zf'(z)}{f(z)} \prec \phi(z)\}$ |
| $S^*[A, B]$ | $\{f \in \mathcal{A} : \frac{zf'(z)}{f(z)} \prec \frac{1+Az}{1+Bz}\}$ |
| $SP(\alpha, A, B)$ | $\{f \in \mathcal{A} : e^{i\alpha} \frac{zf'(z)}{f(z)} \prec \cos \alpha \frac{1+Az}{1+Bz} + i \sin \alpha$ $(-1 \leq B < A \leq 1, 0 \leq \alpha < 1)\}$ |

| | |
|---------------|---|
| S_p | Class of parabolic starlike functions in U |
| T | Subclass of \mathcal{A} consisting of functions of the form $f(z) = z - \sum_{n=2}^{\infty} a_n z^n \quad (z \in U)$ |
| U | Open unit disk $\{z \in \mathcal{C} : z < 1\}$ |
| U^* | Punctured unit disk $U \setminus \{0\}$ |
| \mathcal{Z} | Set of all integers |

KONVOLUSI, PEKALI DAN MASALAH JEJARI UNTUK FUNGSI UNIVALEN TERTENTU

ABSTRAK

Suatu fungsi f yang tertakrif dalam cakera unit $U := \{z \in \mathcal{C} : |z| < 1\}$ dalam satah kompleks \mathcal{C} dikatakan univalen jika fungsi tersebut memetakan titik berlainan dalam U ke titik berlainan dalam \mathcal{C} . Andaikan \mathcal{A} kelas fungsi analisis ternormalkan yang tertakrif dalam U dan mempunyai siri Taylor dalam bentuk

$$(0.0.1) \quad f(z) = z + \sum_{n=2}^{\infty} a_n z^n.$$

Suatu fungsi f dikatakan subordinasi kepada suatu fungsi univalen F jika $f(0) = F(0)$ dan $f(U) \subset F(U)$. Hasil darab Hadamard atau konvolusi dua fungsi analisis, f berbentuk yang seperti (0.0.1) dan $g(z) = z + \sum_{n=2}^{\infty} b_n z^n$, ditakrif sebagai

$$(f * g)(z) = \sum_{n=1}^{\infty} a_n b_n z^n.$$

Andaikan \mathcal{M} kelas fungsi meromorfi, h berbentuk

$$(0.0.2) \quad h(z) = \frac{1}{z} + \sum_{n=0}^{\infty} a_n z^n,$$

yang analisis dan univalen dalam $U^* = \{z : 0 < |z| < 1\}$. Konvolusi dua fungsi meromorfi h dan k , dengan h diberi dalam bentuk (0.0.2) dan $k(z) = \frac{1}{z} + \sum_{n=0}^{\infty} b_n z^n$, ditakrif sebagai

$$(h * k)(z) = \frac{1}{z} + \sum_{n=0}^{\infty} a_n b_n z^n.$$

Dengan menggunakan ciri-ciri konvolusi dan teori subordinasi, beberapa subkelas fungsi meromorfi diperkenalkan. Dengan mensubordinasikan fungsi di dalam kelas ini dengan suatu fungsi cembung ternormalkan yang mempunyai nilai nyata positif, subkelas-subkelas ini merangkumi subkelas klasik meromorfi bak-bintang, cembung, hampir-cembung dan kuasi-cembung. Hubungan kelas dan ciri-ciri konvolusi subkelas-subkelas ini juga dikaji.

Andaikan T subkelas \mathcal{A} yang mengandung fungsi t dalam bentuk

$$t(z) = z - \sum_{n=2}^{\infty} |a_n| z^n.$$

Silverman [44] telah menyiasat subkelas T yang mengandung fungsi bak-bintang peringkat α dan cembung peringkat α ($0 \leq \alpha < 1$). Dengan motivasi ini, pelbagai subkelas T telah dikaji. Kelas $T[\{b_k\}_{m+1}^{\infty}, \beta, m]$ yang ditakrifkan secara am akan dikaji dan anggaran pekali, teorem pertumbuhan dan beberapa keputusan untuk kelas ini diperoleh. Keputusan ini merangkumi beberapa keputusan awal sebagai kes khas.

Untuk pemalar kompleks A dan B , andaikan $S^*[A, B]$ kelas yang mengandung fungsi analisis ternormalkan yang mematuhi $\frac{zf'(z)}{f(z)} \prec \frac{1+Az}{1+Bz}$. Jejari bak-bintang peringkat α , jejari bak-bintang kuat dan jejari bak-bintang parabola diperoleh untuk kelas $S^*[A, B]$. Keputusan ini juga merangkumi beberapa keputusan awal.

CONVOLUTION, COEFFICIENT AND RADIUS PROBLEMS OF CERTAIN UNIVALENT FUNCTIONS

ABSTRACT

A function f defined on the open unit disc $U := \{z \in \mathcal{C} : |z| < 1\}$ of the complex plane \mathcal{C} is univalent if it maps different points of U to different points in \mathcal{C} . Let \mathcal{A} denote the class of analytic functions defined on U which is normalized and has the Taylor series of the form

$$(0.0.3) \quad f(z) = z + \sum_{n=2}^{\infty} a_n z^n.$$

The function f is subordinate to a univalent function F if $f(0) = F(0)$ and $f(U) \subset F(U)$. Hadamard product or convolution of two analytic functions f , given by (0.0.3) and $g(z) = z + \sum_{n=2}^{\infty} b_n z^n$ is given by

$$(f * g)(z) = \sum_{n=1}^{\infty} a_n b_n z^n.$$

Let \mathcal{M} denote the class of meromorphic functions h of the form

$$(0.0.4) \quad h(z) = \frac{1}{z} + \sum_{n=0}^{\infty} a_n z^n,$$

that are analytic and univalent in the punctured unit disk $U^* = \{z : 0 < |z| < 1\}$. The convolution of two meromorphic functions h and k , where h is given by (0.0.4) and $k(z) = \frac{1}{z} + \sum_{n=0}^{\infty} b_n z^n$, is given by

$$(h * k)(z) = \frac{1}{z} + \sum_{n=0}^{\infty} a_n b_n z^n.$$

By making use of the properties of convolution and theory of subordination, several subclasses of meromorphic functions are introduced. Subjecting each convoluted-derived function in the class to be subordinated to a given normalized convex function with positive real part, these subclasses extend the classical subclasses of meromorphic starlikeness, convexity, close-to-convexity, and quasi-convexity. Class relations, as well as inclusion and convolution properties of these subclasses are investigated.

Let T denote the subclass of \mathcal{A} consisting of functions t of the form

$$t(z) = z - \sum_{n=2}^{\infty} |a_n| z^n.$$

Silverman [44] investigated the subclasses of T consisting of functions which are starlike of order α and convex of order α ($0 \leq \alpha < 1$). Motivated by his work, many other subclasses of T were studied in the literature. The class $T[\{b_k\}_{m+1}^{\infty}, \beta, m]$ which is defined in a general manner is studied and the coefficient estimate, growth theorem and other results for this class are obtained. Our results contain several earlier results as special cases.

For complex constants A and B , let $S^*[A, B]$ be the class consisting of normalized analytic functions f satisfying $\frac{zf'(z)}{f(z)} \prec \frac{1+Az}{1+Bz}$. The radius of starlikeness of order α , radius of strong-starlikeness, and radius of parabolic-starlikeness are obtained for $S^*[A, B]$. Several known results are shown to be simple consequences of results derived here.

CHAPTER 1

INTRODUCTION

The theory of univalent functions is a remarkable area of study. This field which is more often associated with 'geometry' and 'analysis' has raised the interest of many since the beginning of 20th century to recent times. The name univalent functions or schlicht (the German word for simple) functions is given to functions defined on the open unit disc $U := \{z \in \mathcal{C} : |z| < 1\}$ of the complex plane \mathcal{C} that are characterized by the fact that such a function provides one-to-one mapping onto its image.

Let $\mathcal{H}(U)$ be the class of all analytic functions on U and \mathcal{A} denote the class of analytic functions defined on U which is normalized by the condition $f(0) = 0$, $f'(0) = 1$ and has the Taylor series of the form

$$f(z) = z + \sum_{n=2}^{\infty} a_n z^n.$$

Geometrically, the function f is univalent if $f(z_1) = f(z_2)$ implies $z_1 = z_2$ in U and is locally univalent at $z_0 \in U$ if it is univalent in some neighborhood of z_0 . The subclass of \mathcal{A} consisting of univalent functions is denoted by \mathcal{S} .

The Koebe function $k(z) = z/(1 - z)^2$ is a univalent function and it plays a very significant role in the study of the class \mathcal{S} . In fact, the Koebe function and its rotations $e^{-i\alpha}k(e^{i\alpha}z)$, $\alpha \in \mathcal{R}$ are the only extremal functions for various problems in the class \mathcal{S} . For example, the famous findings of Bieberbach. In 1916, Bieberbach proved that if $f \in \mathcal{S}$, then the second coefficient $|a_2| \leq 2$ with equality if and only if f is a rotation of the Koebe function. He also conjectured that $|a_n| \leq n$, ($n = 2, 3, \dots$) which is generally valid and this was proved by de Branges [6] in 1985.

Several special subclasses of analytic univalent functions play prominent role in the study of this area. Notable among them are the classes of starlike and convex functions.

Let w_0 be an interior point of a set \mathcal{D} in the complex plane. The set \mathcal{D} is starlike with respect to w_0 if the line segment joining w_0 to every other point in \mathcal{D} lies in the interior of \mathcal{D} . If a function $f \in \mathcal{A}$ maps U onto a starlike domain, then f is a starlike function. The class of starlike functions with respect to origin is denoted by S^* . Analytically,

$$S^* := \left\{ f \in \mathcal{A} : \Re \left(\frac{zf'(z)}{f(z)} \right) > 0 \right\}.$$

A set \mathcal{D} in the complex plane is convex if for every pair of points w_1 and w_2 in the interior of \mathcal{D} , the line segment joining w_1 and w_2 lies in the interior of \mathcal{D} . If a function $f \in \mathcal{A}$ maps U onto a convex domain, then f is a convex function. Let K denote the class of all convex functions in \mathcal{A} . An analytic description of the class K is given by

$$K := \left\{ f \in \mathcal{A} : \Re \left(1 + \frac{zf''(z)}{f'(z)} \right) > 0 \right\}.$$

The well known connection between these two classes was first observed by Alexander in 1915. The Alexander theorem [2] states that for an analytic function f , $f(z) \in K$ if and only if $zf'(z) \in S^*$.

Ma and Minda gave a unified presentation of these classes by using the method of subordination. For two functions f and g analytic in U , the function f is subordinate to g , written

$$f(z) \prec g(z) \quad (z \in U),$$

if there exists a function w , analytic in U with $w(0) = 0$ and $|w(z)| < 1$ such that $f(z) = g(w(z))$. In particular, if the function g is univalent in U , then $f(z) \prec g(z)$ is equivalent to $f(0) = g(0)$ and $f(U) \subset g(U)$.

With $g(z) = (1+z)/(1-z)$, a function $f \in \mathcal{A}$ is starlike if $zf'(z)/f(z)$ is subordinate to g and is convex if $1+zf''(z)/f'(z)$ is subordinate to g . Ma and Minda [17] introduced the classes

$$S^*(\phi) = \left\{ f \in \mathcal{A} \mid \frac{zf'(z)}{f(z)} \prec \phi(z) \right\}$$

and

$$K(\phi) = \left\{ f \in \mathcal{A} \mid 1 + \frac{zf''(z)}{f'(z)} \prec \phi(z) \right\},$$

where ϕ is an analytic function with positive real part, $\phi(0) = 1$ and ϕ maps the unit disk U onto a region starlike with respect to 1.

The convolution or Hadamard product is another interesting exploration of these classes. The convolution of two analytic functions $f(z) = z + \sum_{n=2}^{\infty} a_n z^n$ and $g(z) = z + \sum_{n=2}^{\infty} b_n z^n$ is given by

$$(f * g)(z) = \sum_{n=1}^{\infty} a_n b_n z^n.$$

Polya-Schoenberg [24] conjectured that the class of convex functions is preserved under convolution with convex functions. In 1973, Ruscheweyh and Sheil-Small [36] proved the Polya-Schoenberg conjecture. In fact, they proved that the classes of starlike functions and convex functions are closed under convolution with convex functions.

Detailed treatment of univalent functions are available in books by Pommerenke [25], Duren [7] and Goodman [12].

CHAPTER 2

CERTAIN SUBCLASSES OF MEROMORPHIC FUNCTIONS ASSOCIATED WITH CONVOLUTION AND DIFFERENTIAL SUBORDINATION

2.1. MOTIVATION AND PRELIMINARIES

The convolution or the Hadamard product of two analytic functions $f(z) = \sum_{n=1}^{\infty} a_n z^n$ and $g(z) = \sum_{n=1}^{\infty} b_n z^n$ is given by

$$(f * g)(z) = \sum_{n=1}^{\infty} a_n b_n z^n.$$

The geometric series $\sum_{n=1}^{\infty} z^n = z/(1-z)$ acts as the identity element under convolution. The convolution of f with the geometric series $\sum_{n=1}^{\infty} n z^n = z/(1-z)^2$ is given by $\sum_{n=1}^{\infty} n a_n z^n$ which is equivalent to $z f'(z)$. In terms of convolution, $f = f * (z/(1-z))$ and $z f' = f * (z/(1-z)^2)$. The well known Alexander's theorem states that a function f is convex if and only if $z f'$ is starlike. Since $z f' = f * (z/(1-z)^2)$, it follows that f is convex if and only if $f * (z/(1-z)^2)$ is starlike. Also, a function f is starlike if $f * (z/(1-z))$ is starlike. These ideas led to the study of the class of all functions f such that $f * g$ is starlike for some fixed function g in \mathcal{A} . In this direction, Shanmugam [41] introduced and investigated various subclasses of analytic functions by using the convex hull method [5, 23, 36] and the method of differential subordination. Ravichandran [26] introduced certain classes of analytic functions with respect to n -ply symmetric points, conjugate points and symmetric conjugate points and also discussed their convolution properties. Some other related studies were also made in [3, 22], and more recently by Shamani *et al.* [40].

Let \mathcal{M} denote the class of meromorphic functions f of the form

$$(2.1.1) \quad f(z) = \frac{1}{z} + \sum_{n=0}^{\infty} a_n z^n,$$

that are analytic and univalent in the punctured unit disk $U^* = \{z : 0 < |z| < 1\}$.

The convolution of two meromorphic functions f and g , where f is given by (2.1.1) and

$$(2.1.2) \quad g(z) = \frac{1}{z} + \sum_{n=0}^{\infty} b_n z^n,$$

is given by

$$(f * g)(z) = \frac{1}{z} + \sum_{n=0}^{\infty} a_n b_n z^n.$$

For $0 \leq \alpha < 1$, we recall that the classes of meromorphic starlike, meromorphic convex, meromorphic close-to-convex, meromorphic γ -convex (Mocanu sense) and meromorphic quasi-convex functions of order α , denoted by \mathcal{M}^s , \mathcal{M}^k , \mathcal{M}^c , \mathcal{M}_γ^k and \mathcal{M}^q respectively, are defined by

$$(2.1.3) \quad \begin{aligned} \mathcal{M}^s &= \left\{ f \in \mathcal{M} \mid -\Re \frac{zf'(z)}{f(z)} > \alpha \right\}, \\ \mathcal{M}^k &= \left\{ f \in \mathcal{M} \mid -\Re \left(1 + \frac{zf''(z)}{f'(z)} \right) > \alpha \right\}, \\ \mathcal{M}^c &= \left\{ f \in \mathcal{M} \mid -\Re \frac{zf'(z)}{g(z)} > \alpha, g(z) \in \mathcal{M}^s \right\}, \\ \mathcal{M}_\gamma^k &= \left\{ f \in \mathcal{M} \mid -\Re \left[(1 - \gamma) \frac{zf'(z)}{f(z)} + \gamma \left(1 + \frac{zf''(z)}{f'(z)} \right) \right] > \alpha \right\}, \\ \mathcal{M}^q &= \left\{ f \in \mathcal{M} \mid -\Re \frac{[zf'(z)]'}{g'(z)} > \alpha, g(z) \in \mathcal{M}^k \right\}. \end{aligned}$$

Motivated by the investigation of Shanmugam [41], Ravichandran [26], and Ali *et al.* [3], several subclasses of meromorphic functions defined by means of convolution with a given fixed meromorphic function are introduced in Section 2.2. These new subclasses extend the classical classes of meromorphic starlike, convex, close-to-convex, γ -convex, and quasi-convex functions given in (2.1.3). Section 2.3 is devoted

to the investigation of the class relations as well as inclusion and convolution properties of these newly defined classes.

We shall need the following definition and results to prove our main results.

Let $S^*(\alpha)$ denote the class of starlike functions of order α . The class R_α of prestarlike functions of order α is defined by

$$R_\alpha = \left\{ f \in \mathcal{A} \mid f(z) * \frac{z}{(1-z)^{2-2\alpha}} \in S^*(\alpha) \right\}$$

for $\alpha < 1$, and

$$R_1 = \left\{ f \in \mathcal{A} \mid \Re \frac{f(z)}{z} > \frac{1}{2} \right\}.$$

THEOREM 2.1.1. [35, Theorem 2.4] *Let $\alpha \leq 1$, $f \in R_\alpha$ and $g \in S^*(\alpha)$. Then, for any analytic function $H \in \mathcal{H}(U)$,*

$$\frac{f * Hg}{f * g}(U) \subset \overline{co}(H(U))$$

where $\overline{co}(H(U))$ denotes the closed convex hull of $H(U)$.

THEOREM 2.1.2. [8] *Let h be convex in U and $\beta, \gamma \in \mathcal{C}$ with $\Re(\beta h(z) + \gamma) > 0$. If p is analytic in U with $p(0) = h(0)$, then*

$$p(z) + \frac{zp'(z)}{\beta p(z) + \gamma} \prec h(z) \quad \text{implies} \quad p(z) \prec h(z).$$

We will also be using the following convolution properties.

(i) For two meromorphic functions f and g of the forms $f(z) = \frac{1}{z} + \sum_{n=0}^{\infty} a_n z^n$ and $g(z) = \frac{1}{z} + \sum_{n=0}^{\infty} b_n z^n$, we have

$$(f * g)(z) = (g * f)(z)$$

PROOF. For f and g of the form $f(z) = \frac{1}{z} + \sum_{n=0}^{\infty} a_n z^n$ and $g(z) = \frac{1}{z} + \sum_{n=0}^{\infty} b_n z^n$, we have

$$\begin{aligned} (f * g)(z) &= \frac{1}{z} + \sum_{n=0}^{\infty} a_n b_n z^n \\ &= \frac{1}{z} + \sum_{n=0}^{\infty} b_n a_n z^n \\ &= (g * f)(z). \end{aligned} \quad \square$$

(ii) For two meromorphic functions f and g of the forms $f(z) = \frac{1}{z} + \sum_{n=0}^{\infty} a_n z^n$ and $g(z) = \frac{1}{z} + \sum_{n=0}^{\infty} b_n z^n$, we have

$$-z(g * f)'(z) = (g * -zf')(z).$$

PROOF. For f of the form $f(z) = \frac{1}{z} + \sum_{n=0}^{\infty} a_n z^n$, we have

$$-zf'(z) = \frac{1}{z} - \sum_{n=0}^{\infty} n a_n z^n$$

and hence

$$\begin{aligned} (g * -zf')(z) &= \frac{1}{z} - \sum_{n=0}^{\infty} n a_n b_n z^n \\ &= -z \left(-\frac{1}{z^2} + \sum_{n=0}^{\infty} n a_n b_n z^{n-1} \right) \\ &= -z(g * f)'(z). \end{aligned} \quad \square$$

(iii) For two meromorphic functions f and g of the forms $f(z) = \frac{1}{z} + \sum_{n=0}^{\infty} a_n z^n$ and $g(z) = \frac{1}{z} + \sum_{n=0}^{\infty} b_n z^n$, we have

$$z^2(g * f)(z) = (z^2 g * z^2 f)(z).$$

PROOF. For

$$(g * f)(z) = \frac{1}{z} + \sum_{n=0}^{\infty} a_n b_n z^n,$$

we observe that

$$\begin{aligned}
z^2(g * f)(z) &= z^2 \left(\frac{1}{z} + \sum_{n=0}^{\infty} a_n b_n z^n \right) \\
&= z + z^2 \sum_{n=0}^{\infty} a_n b_n z^n \\
&= z^2 g(z) * z^2 f(z). \quad \square
\end{aligned}$$

2.2. DEFINITIONS

In this section, various subclasses of \mathcal{M} are defined by means of convolution and subordination. Let g be a fixed function in \mathcal{M} , and h be a convex univalent function with positive real part in U and $h(0) = 1$.

DEFINITION 2.2.1. The class $\mathcal{M}_g^s(h)$ consists of functions $f \in \mathcal{M}$ satisfying $(g * f)(z) \neq 0$ in U^* and the subordination

$$-\frac{z(g * f)'(z)}{(g * f)(z)} \prec h(z).$$

REMARK 2.2.1. If $g(z) = \frac{1}{z} + \frac{1}{1-z} = \frac{1}{z} + \sum_{n=0}^{\infty} z^n$, then $\mathcal{M}_g^s(h)$ coincides with $\mathcal{M}^s(h)$, where

$$\mathcal{M}^s(h) = \left\{ f \in \mathcal{M} \mid -\frac{zf'(z)}{f(z)} \prec h(z) \right\}.$$

DEFINITION 2.2.2. The class $\mathcal{M}_g^k(h)$ consists of functions $f \in \mathcal{M}$ satisfying $(g * f)'(z) \neq 0$ in U^* and the subordination

$$-\left\{ 1 + \frac{z(g * f)''(z)}{(g * f)'(z)} \right\} \prec h(z).$$

DEFINITION 2.2.3. The class $\mathcal{M}_g^c(h)$ consists of functions $f \in \mathcal{M}$ such that $(g * \psi)(z) \neq 0$ in U^* for some $\psi \in \mathcal{M}_g^s(h)$ and satisfying the subordination

$$-\frac{z(g * f)'(z)}{(g * \psi)(z)} \prec h(z).$$

DEFINITION 2.2.4. For γ real, the class $\mathcal{M}_{g,\gamma}^k(h)$ consists of functions $f \in \mathcal{M}$ satisfying $(g * f)(z) \neq 0$, $(g * f)'(z) \neq 0$ in U^* and the subordination

$$-\left\{ \gamma \left(1 + \frac{z(g * f)''(z)}{(g * f)'(z)} \right) + (1 - \gamma) \left(\frac{z(g * f)'(z)}{(g * f)(z)} \right) \right\} \prec h(z).$$

REMARK 2.2.2. For $\gamma = 0$, the class $\mathcal{M}_{g,\gamma}^k(h)$ coincides with the class $\mathcal{M}_g^s(h)$ and for $\gamma = 1$, it reduces to the class $\mathcal{M}_g^k(h)$.

DEFINITION 2.2.5. The class $\mathcal{M}_g^q(h)$ consists of functions $f \in \mathcal{M}$ such that $(g * \varphi)'(z) \neq 0$ in U^* for some $\varphi \in \mathcal{M}_g^k(h)$ and satisfying the subordination

$$\frac{[-z(g * f)'(z)]'}{(g * \varphi)'(z)} \prec h(z).$$

2.3. INCLUSION AND CONVOLUTION THEOREM

This section is devoted to the investigation of class relations as well as inclusion and convolution properties of the new subclasses given in Section 2.2.

We will begin with the theorem which is analogue to the well known Alexander's theorem.

THEOREM 2.3.1. *The function f is in $\mathcal{M}_g^k(h)$ if and only if $-zf'$ is in $\mathcal{M}_g^s(h)$.*

PROOF. Since

$$\begin{aligned} -\left(1 + \frac{z(g * f)''(z)}{(g * f)'(z)} \right) &= -\frac{(g * f)'(z) + z(g * f)''(z)}{(g * f)'(z)} \\ &= -\frac{(z(g * f)'(z))'}{(g * f)'(z)} \\ &= \frac{-z}{-z} \cdot \frac{((g * -zf')(z))'}{(g * f)'(z)} \\ &= -\frac{z(g * -zf')'(z)}{(g * -zf')(z)}, \end{aligned}$$

it follows that $f \in \mathcal{M}_g^k(h)$ if and only if $-zf' \in \mathcal{M}_g^s(h)$. □

THEOREM 2.3.2. *Let h be a convex univalent function satisfying $\Re h(z) < 2 - \alpha$, $0 \leq \alpha < 1$, and $\phi \in \mathcal{M}$ with $z^2\phi \in R_\alpha$. If $f \in \mathcal{M}_g^s(h)$, then $\phi * f \in \mathcal{M}_g^s(h)$.*

PROOF. Since $f \in \mathcal{M}_g^s(h)$, it follows that

$$-\Re \left\{ \frac{z(g * f)'(z)}{(g * f)(z)} \right\} < 2 - \alpha$$

or

$$(2.3.1) \quad \Re \left\{ \frac{z(g * f)'(z) + 2(g * f)(z)}{(g * f)(z)} \right\} > \alpha.$$

The inequality (2.3.1) yields

$$\Re \left\{ \frac{z^2 z(g * f)'(z) + 2z^2(g * f)(z)}{z^2(g * f)(z)} \right\} > \alpha$$

and thus

$$(2.3.2) \quad \Re \left\{ \frac{z(z^2(g * f))'(z)}{z^2(g * f)(z)} \right\} > \alpha.$$

Let

$$P(z) = -\frac{z(g * f)'(z)}{(g * f)(z)}.$$

We have

$$\begin{aligned} -\frac{z(\phi * g * f)'(z)}{(\phi * g * f)(z)} &= \frac{\phi(z) * -z(g * f)'(z)}{\phi(z) * (g * f)(z)} \\ &= \frac{\phi(z) * (g * f)(z)P(z)}{\phi(z) * (g * f)(z)} \cdot \frac{z^2}{z^2} \\ &= \frac{z^2 \phi(z) * z^2(g * f)(z)P(z)}{z^2 \phi(z) * z^2(g * f)(z)}. \end{aligned}$$

Inequality (2.3.2) shows that $z^2(g * f) \in S^*(\alpha)$. Therefore Theorem 2.1.1 yields

$$-\frac{z(\phi * g * f)'(z)}{(\phi * g * f)(z)} = \frac{z^2 \phi(z) * z^2(g * f)(z)P(z)}{z^2 \phi(z) * z^2(g * f)(z)} \in \overline{co}(P(U)),$$

and since $P(z) \prec h(z)$, it follows that

$$-\frac{z(\phi * g * f)'(z)}{(\phi * g * f)(z)} \prec h(z).$$

Hence $\phi * f \in \mathcal{M}_g^s(h)$. □

COROLLARY 2.3.1. $\mathcal{M}_g^s(h) \subset \mathcal{M}_{\phi * g}^s(h)$ under the conditions of Theorem 2.3.2.

PROOF. Let $f \in \mathcal{M}_g^s(h)$, then by Theorem 2.3.2 we have $\phi * f \in \mathcal{M}_g^s(h)$ or

$$-\frac{z(\phi * g * f)'(z)}{(\phi * g * f)(z)} \prec h(z),$$

which equivalently yields $f \in \mathcal{M}_{\phi * g}^s(h)$. \square

In particular, when $g(z) = \frac{1}{z} + \frac{1}{1-z}$, the following corollary is obtained.

COROLLARY 2.3.2. *Let h and ϕ satisfy the conditions of Theorem 2.3.2. If $f \in \mathcal{M}^s(h)$, then $f \in \mathcal{M}_\phi^s(h)$.*

THEOREM 2.3.3. *Let h and ϕ satisfy the conditions of Theorem 2.3.2. If $f \in \mathcal{M}_g^k(h)$, then $\phi * f \in \mathcal{M}_g^k(h)$. Equivalently $\mathcal{M}_g^k(h) \subset \mathcal{M}_{\phi * g}^k(h)$.*

PROOF. If $f \in \mathcal{M}_g^k(h)$, then it follows from Theorem 2.3.1 that $-zf' \in \mathcal{M}_g^s(h)$. Theorem 2.3.2 shows that $-z(\phi * f)' = \phi * -zf' \in \mathcal{M}_g^s(h)$. Hence $\phi * f \in \mathcal{M}_g^k(h)$. \square

THEOREM 2.3.4. *Let h and ϕ satisfy the conditions of Theorem 2.3.2. If $f \in \mathcal{M}_g^c(h)$ with respect to $\psi \in \mathcal{M}_g^s(h)$, then $\phi * f \in \mathcal{M}_g^c(h)$ with respect to $\phi * \psi \in \mathcal{M}_g^s(h)$.*

PROOF. Since $\psi \in \mathcal{M}_g^s(h)$, Theorem 2.3.2 shows that $\phi * \psi \in \mathcal{M}_g^s(h)$ and inequality (2.3.2) yields $z^2(g * \psi) \in S^*(\alpha)$.

Let the function G be defined by

$$G(z) = -\frac{z(g * f)'(z)}{(g * \psi)(z)}.$$

Observe that

$$\begin{aligned} -\frac{z(\phi * g * f)'(z)}{(\phi * g * \psi)(z)} &= \frac{\phi(z) * -z(g * f)'(z)}{\phi(z) * (g * \psi)(z)} \\ &= \frac{\phi(z) * (g * \psi)(z)G(z)}{\phi(z) * (g * \psi)(z)} \cdot \frac{z^2}{z^2} \\ &= \frac{z^2\phi(z) * z^2(g * \psi)(z)G(z)}{z^2\phi(z) * z^2(g * \psi)(z)}. \end{aligned}$$

Since $z^2\phi \in R_\alpha$ and $z^2(g * \psi) \in S^*(\alpha)$, it follows from Theorem 2.1.1 that

$$(2.3.3) \quad -\frac{z(\phi * g * f)'(z)}{(\phi * g * \psi)(z)} = \frac{z^2\phi(z) * z^2(g * \psi)(z)G(z)}{z^2\phi(z) * z^2(g * \psi)(z)} \prec h(z).$$

Thus $\phi * f \in \mathcal{M}_g^c(h)$ with respect to $\phi * \psi \in \mathcal{M}_g^s(h)$. \square

COROLLARY 2.3.3. $\mathcal{M}_g^c(h) \subset \mathcal{M}_{\phi * g}^c(h)$ under the conditions of Theorem 2.3.2.

PROOF. If $f \in \mathcal{M}_g^c(h)$ with respect to $\psi \in \mathcal{M}_g^s(h)$, then Theorem 2.3.4 shows that $\phi * f \in \mathcal{M}_g^c(h)$ with respect to $\phi * \psi \in \mathcal{M}_g^s(h)$ which is equivalent to

$$-\frac{z(\phi * g * f)'(z)}{(\phi * g * \psi)(z)} \prec h(z)$$

or $f \in \mathcal{M}_{\phi * g}^c(h)$. Hence $\mathcal{M}_g^c(h) \subset \mathcal{M}_{\phi * g}^c(h)$. \square

THEOREM 2.3.5. Let $\Re(\gamma h(z)) < 0$. Then

- (i) $\mathcal{M}_{g,\gamma}^k(h) \subset \mathcal{M}_g^s(h)$,
- (ii) $\mathcal{M}_{g,\gamma}^k(h) \subset \mathcal{M}_{g,\beta}^k(h)$ for $\gamma < \beta \leq 0$.

PROOF. Define the function $J_g(\gamma; f)$ by

$$J_g(\gamma; f)(z) = -\left\{ \gamma \left(1 + \frac{z(g * f)''(z)}{(g * f)'(z)} \right) + (1 - \gamma) \left(\frac{z(g * f)'(z)}{(g * f)(z)} \right) \right\}.$$

For $f \in \mathcal{M}_{g,\gamma}^k(h)$, it follows that $J_g(\gamma; f)(z) \prec h(z)$. Let the function P be defined by

$$(2.3.4) \quad P(z) = -\frac{z(g * f)'(z)}{(g * f)(z)}.$$

The logarithmic derivative of $P(z)$ yields

$$(2.3.5) \quad \frac{P'(z)}{P(z)} = \frac{1}{z} + \frac{(g * f)''(z)}{(g * f)'(z)} - \frac{(g * f)'(z)}{(g * f)(z)},$$

and multiplication with $-\gamma z$ to (2.3.5) gives

$$(2.3.6) \quad -\gamma \frac{zP'(z)}{P(z)} = -\gamma - \gamma \frac{z(g * f)''(z)}{(g * f)'(z)} + \gamma \frac{z(g * f)'(z)}{(g * f)(z)}.$$

Adding $P(z)$ to (2.3.6) yields

$$\begin{aligned}
P(z) - \gamma \frac{zP'(z)}{P(z)} &= -\gamma - \gamma \frac{z(g * f)''(z)}{(g * f)'(z)} + \gamma \frac{z(g * f)'(z)}{(g * f)(z)} - \frac{z(g * f)'(z)}{(g * f)(z)} \\
&= - \left\{ \gamma \left(1 + \frac{z(g * f)''(z)}{(g * f)'(z)} \right) + (1 - \gamma) \left(\frac{z(g * f)'(z)}{(g * f)(z)} \right) \right\} \\
(2.3.7) \qquad &= J_g(\gamma; f)(z)
\end{aligned}$$

and hence

$$P(z) - \gamma \frac{zP'(z)}{P(z)} \prec h(z).$$

(i) Since $\Re(\gamma h(z)) < 0$ and

$$P(z) - \gamma \frac{zP'(z)}{P(z)} \prec h(z),$$

Theorem 2.1.2 yields $P(z) \prec h(z)$. Hence $f \in \mathcal{M}_g^s(h)$ and this concludes that $\mathcal{M}_{g,\gamma}^k(h) \subset \mathcal{M}_g^s(h)$.

(ii) The logarithmic derivative of $P(z)$ and multiplication of z yields

$$(2.3.8) \qquad \frac{zP'(z)}{P(z)} = 1 + \frac{z(g * f)''(z)}{(g * f)'(z)} + P(z).$$

From (2.3.7), it follows that

$$(2.3.9) \qquad \frac{zP'(z)}{P(z)} = \frac{P(z) - J_g(\gamma; f)}{\gamma}.$$

Let

$$(2.3.10) \qquad J_g(\beta; f)(z) = - \left\{ \beta \left(1 + \frac{z(g * f)''(z)}{(g * f)'(z)} \right) + (1 - \beta) \left(\frac{z(g * f)'(z)}{(g * f)(z)} \right) \right\}.$$

Substituting (2.3.8) and (2.3.9) in (2.3.10) yield

$$(2.3.11) \qquad J_g(\beta; f)(z) = \left(1 - \frac{\beta}{\gamma} \right) P(z) + \frac{\beta}{\gamma} J_g(\gamma; f)(z).$$

We know that $J_g(\gamma; f)(z) \prec h(z)$ and $P(z) \prec h(z)$ from (i) and since $0 < \frac{\beta}{\gamma} < 1$ and $h(U)$ is convex, we deduce that $J_g(\beta; f)(z) \in h(U)$. Therefore, $J_g(\beta; f)(z) \prec h(z)$ and hence $\mathcal{M}_{g,\gamma}^k(h) \subset \mathcal{M}_{g,\beta}^k(h)$ for $\gamma < \beta \leq 0$ □

COROLLARY 2.3.4. *The class $\mathcal{M}_g^k(h)$ is a subset of the class $\mathcal{M}_g^q(h)$.*

PROOF. Let $f \in \mathcal{M}_g^k(h)$ and by taking $f = \varphi$, it follows from the definition of the class $\mathcal{M}_g^q(h)$ that $\mathcal{M}_g^k(h) \subset \mathcal{M}_g^q(h)$. \square

THEOREM 2.3.6. *The function f is in $\mathcal{M}_g^q(h)$ if and only if $-zf'$ is in $\mathcal{M}_g^c(h)$.*

PROOF. If $f \in \mathcal{M}_g^q(h)$, then there exists $\varphi \in \mathcal{M}_g^k(h)$ such that

$$\frac{[-z(g * f)'(z)]'}{(g * \varphi)'(z)} \prec h(z).$$

Note that

$$\frac{[-z(g * f)'(z)]'}{(g * \varphi)'(z)} = \frac{[(g * -zf')(z)]'}{(g * \varphi)'(z)} \cdot \frac{-z}{-z} = \frac{-z(g * -zf')'(z)}{(g * -z\varphi)'(z)}.$$

Hence

$$\frac{-z(g * -zf')'(z)}{(g * -z\varphi)'(z)} \prec h(z).$$

Since $\varphi \in \mathcal{M}_g^k(h)$, by Theorem 2.3.1, $-z\varphi' \in \mathcal{M}_g^s(h)$. Thus by definition 2.2.3, we have $-zf' \in \mathcal{M}_g^c(h)$.

Conversely, if $-zf' \in \mathcal{M}_g^c(h)$, then

$$-\frac{z(g * -zf')'(z)}{(g * \varphi_1)(z)} \prec h(z)$$

for some $\varphi_1 \in \mathcal{M}_g^s(h)$. Let $\varphi \in \mathcal{M}_g^k(h)$ be such that $-z\varphi' = \varphi_1 \in \mathcal{M}_g^s(h)$. The proof is completed by observing that

$$\frac{[-z(g * f)'(z)]'}{(g * \varphi)'(z)} = -\frac{z(g * -zf')'(z)}{(g * -z\varphi)'(z)} \prec h(z). \quad \square$$

COROLLARY 2.3.5. *Let h and ϕ satisfy the conditions of Theorem 2.3.2. If $f \in \mathcal{M}_g^q(h)$, then $\phi * f \in \mathcal{M}_g^q(h)$.*

PROOF. If $f \in \mathcal{M}_g^q(h)$, Theorem 2.3.6 gives $-zf' \in \mathcal{M}_g^c(h)$. Theorem 2.3.4 next gives $\phi * -zf' = -z(\phi * f)' \in \mathcal{M}_g^c(h)$. Thus, Theorem 2.3.6 yields $\phi * f \in \mathcal{M}_g^q(h)$. \square

COROLLARY 2.3.6. $\mathcal{M}_g^q(h) \subset \mathcal{M}_{\phi * g}^q(h)$ under the conditions of Theorem 2.3.2.

PROOF. If $f \in \mathcal{M}_g^q(h)$, it follows from Corollary 2.3.5 that $\phi * f \in \mathcal{M}_g^q(h)$. The subordination

$$\frac{[-z(\phi * g * f)'(z)]'}{(\phi * g * \varphi)'(z)} \prec h(z)$$

gives $f \in \mathcal{M}_{\phi * g}^q(h)$. Therefore $\mathcal{M}_g^q(h) \subset \mathcal{M}_{\phi * g}^q(h)$. □

CHAPTER 3

A GENERALIZED CLASS OF UNIVALENT FUNCTIONS WITH NEGATIVE COEFFICIENTS

3.1. MOTIVATION AND PRELIMINARIES

Let T denote the subclass of \mathcal{A} consisting of functions f of the form

$$f(z) = z - \sum_{n=2}^{\infty} |a_n| z^n.$$

A function $f \in T$ is called a function with negative coefficients. In [44], Silverman investigated the subclasses of T which were denoted by $TS^*(\alpha)$ and $TK(\alpha)$ respectively consisting of functions which are starlike of order α and convex of order α ($0 \leq \alpha < 1$). He proved the following:

THEOREM 3.1.1. [44] *Let $f(z) = z - \sum_{n=2}^{\infty} |a_n| z^n$. Then $f \in TS^*(\alpha)$ if and only if $\sum_{n=2}^{\infty} (n - \alpha) |a_n| \leq 1 - \alpha$.*

COROLLARY 3.1.1. [44] *Let $f(z) = z - \sum_{n=2}^{\infty} |a_n| z^n$. Then $f \in TK(\alpha)$ if and only if $\sum_{n=2}^{\infty} n(n - \alpha) |a_n| \leq 1 - \alpha$.*

The work of Silverman has brought special interest in the exploration of the functions with negative coefficients. Motivated by his work, many other subclasses of T were studied in the literature. For example, the class $\mathcal{U}(k, \tau, \alpha)$ defined below, was studied by Shanmugam [43].

DEFINITION 3.1.1. [43] For $0 \leq \tau \leq 1$, $0 \leq \alpha < 1$ and $k \geq 0$, let $\mathcal{U}(k, \tau, \alpha)$, consist of functions $f \in T$ satisfying the condition

$$\Re \left\{ \frac{\tau z^3 f'''(z) + (1 + 2\tau) z^2 f''(z) + z f'(z)}{\tau z^2 f''(z) + z f'(z)} \right\}$$

$$> k \left| \frac{\tau z^3 f'''(z) + (1 + 2\tau)z^2 f''(z) + z f'(z)}{\tau z^2 f''(z) + z f'(z)} - 1 \right| + \alpha.$$

The class $\mathcal{U}(k, \tau, \alpha)$ contains well known classes as special cases. In particular, $\mathcal{U}(k, 0, 0)$ is the class of k -uniformly convex function introduced and studied by Kanas and Wisniowska [15] and $\mathcal{U}(0, 0, \alpha)$ coincides with the class $TK(\alpha)$ studied in [44]. In [43], Shanmugam proved the coefficients bounds, extreme points as well as radius of starlikeness and convexity theorem for functions in $\mathcal{U}(k, \tau, \alpha)$.

Kadioglu in [13] extended the results by Silverman by defining the class $T_s(\alpha)$ with the use of Sălăgean derivative operator and proved some properties of the functions in this class. The Sălăgean derivatives operator was introduced in [38], where for $f(z) \in \mathcal{A}$,

$$D^0 f(z) = f(z), \quad D^1 f(z) = z f'(z)$$

and

$$D^s f(z) = D(D^{s-1} f(z)) \quad (s = 1, 2, 3, \dots).$$

Observe that

$$D^0 f(z) = f(z) = z - \sum_{n=2}^{\infty} |a_n| z^n,$$

$$D^1 f(z) = z f'(z) = z \left(1 - \sum_{n=2}^{\infty} n |a_n| z^{n-1} \right) = z - \sum_{n=2}^{\infty} n |a_n| z^n$$

and

$$D^2 f(z) = D(D^1 f(z)) = z(z f'(z))' = z - \sum_{n=2}^{\infty} n^2 |a_n| z^n$$

⋮

Hence

$$D^s f(z) = D(D^{s-1} f(z)) = z - \sum_{n=2}^{\infty} n^s |a_n| z^n.$$

Kadioglu defined the class $T_s(\alpha)$ as the following:

DEFINITION 3.1.2. [13]

$$T_s(\alpha) = \left\{ f \in T : \Re \left\{ \frac{D^{s+1} f(z)}{D^s f(z)} \right\} > \alpha \right\}$$

Note that $T_0(\alpha) = TS^*(\alpha)$ and $T_1(\alpha) = TK(\alpha)$. He proved the following:

THEOREM 3.1.2. [13] *A function $f(z) = z - \sum_{n=2}^{\infty} |a_n|z^n$ is in $T_s(\alpha)$ if and only if*

$$\sum_{n=2}^{\infty} (n^{s+1} - n^s \alpha) |a_n| \leq 1 - \alpha.$$

Ahuja [1] defined the class $T_\lambda(\alpha)$ with the use of Ruscheweyh derivative operator. The Ruscheweyh derivative operator D^λ is defined using the Hadamard product or convolution by

$$D^\lambda f = \frac{z}{(1-z)^{\lambda+1}} * f \quad \text{for } \lambda \geq -1.$$

DEFINITION 3.1.3. [1] A function $f \in T$ is said to be in the class $T_\lambda(\alpha)$ if it satisfies

$$\Re \left\{ z \frac{(D^\lambda f(z))'}{D^\lambda f(z)} \right\} > \alpha$$

for $\lambda > -1$ and $\alpha < 1$.

By letting $\lambda = 0$ and $\lambda = 1$, the class $T_\lambda(\alpha)$ will reduce to the class $TS^*(\alpha)$ and $TK(\alpha)$ respectively. The following theorem was proved in [1].

THEOREM 3.1.3. *A function $f(z) = z - \sum_{n=2}^{\infty} a_n z^n$ is in $T_\lambda(\alpha)$ if and only if*

$$\sum_{n=2}^{\infty} (n - \alpha) B_n(\lambda) a_n \leq 1 - \alpha$$

where

$$B_n(\lambda) = \frac{(\lambda + 1)_{n-1}}{(n-1)!} = \frac{(\lambda + 1)(\lambda + 2) \dots (\lambda + n - 1)}{(n-1)!}.$$

The above defined classes as well as numerous other classes of functions can be investigated in a unified manner. For this purpose, we study the following class of $T[\{b_k\}_{m+1}^{\infty}, \beta, m]$.

DEFINITION 3.1.4. Let $b_{m+1} > 0$, $b_{m+1} \leq b_k$ for $k \geq m + 1$. Also let $\beta \geq 0$ and $m \geq 1$ be an integer. The class $T[\{b_k\}_{m+1}^\infty, \beta, m]$ is defined by

$$T[\{b_k\}_{m+1}^\infty, \beta, m] = \left\{ f = z - \sum_{k=m+1}^{\infty} a_k z^k \mid a_k \geq 0 \text{ for } k \geq m + 1, \sum_{k=m+1}^{\infty} b_k a_k \leq \beta \right\}.$$

For convenience, we will denote $T[\{b_k\}_{m+1}^\infty, \beta, m]$ as $T[b_k, \beta, m]$ and adopt this notation hereafter.

For $m = 1$, the class $T[b_k, \beta, m]$ coincides with the class introduced by Frasin [9]. In his paper, Frasin investigated the partial sums of functions belonging to this class. There are many subclasses of T studied by various authors of which several can be represented as $T[b_k, \beta, m]$ with suitable choices of b_k, β and m (see example below) (also see [4, 14, 18, 19, 20, 34, 37, 39]).

EXAMPLE 3.1.1.

- (1) $T[\frac{k-\alpha}{1-\alpha} \frac{(\lambda+1)_{k-1}}{(k-1)!}(\lambda), 1, 1] = T_\lambda(\alpha)$, [1]
- (2) $T[n^{s+1} - \alpha n^s, 1 - \alpha, 1] = T_s(\alpha)$, [13]
- (3) $T[n((n - \alpha)(\tau n - \tau + 1) + k(\tau + n - 1)), 1 - \alpha, 1] = U(k, \tau, \alpha)$, [43]
- (4) $T[n - \alpha, 1 - \alpha, 1] = TS^*(\alpha)$, [44]
- (5) $T[n(n - \alpha), 1 - \alpha, 1] = TK(\alpha)$. [44]

We will be using Example 3.1.1 later to prove the corollaries in this chapter.

In this chapter we obtain the coefficient estimates, growth theorem, distortion theorem, covering theorem and closure theorem for the class $T[b_k, \beta, m]$. We also consider the extreme points and investigate radius problem for this class.

3.2. COEFFICIENT ESTIMATE

We begin with the theorem which gives us the estimate for the coefficient of functions in the class $T[b_k, \beta, m]$.

THEOREM 3.2.1. *If $f \in T[b_k, \beta, m]$, then*

$$a_j \leq \frac{\beta}{b_j}, \quad (j = m + 1, m + 2, \dots)$$

with the equality only for functions of the form $f_j(z) = z - \frac{\beta}{b_j}z^j$.

PROOF. Let $f \in T[b_k, \beta, m]$, then by definition, we have

$$\sum_{k=m+1}^{\infty} b_k a_k \leq \beta.$$

Hence, it follows that

$$b_j a_j \leq \sum_{k=m+1}^{\infty} b_k a_k \leq \beta \quad (j = m + 1, m + 2, \dots)$$

or

$$a_j \leq \frac{\beta}{b_j}.$$

It is clear that for the function of the form

$$f_j(z) = z - \frac{\beta}{b_j}z^j \in T[b_k, \beta, m]$$

we have

$$a_j = \frac{\beta}{b_j}. \quad \square$$

COROLLARY 3.2.1. [44] *If $f \in TS^*(\alpha)$, then*

$$a_n \leq \frac{1 - \alpha}{n - \alpha},$$

with equality for function of the form

$$f_n(z) = z - \frac{1 - \alpha}{n - \alpha}z^n.$$

COROLLARY 3.2.2. [43] *If $f \in \mathcal{U}(k, \tau, \alpha)$, then*

$$a_n \leq \frac{1 - \alpha}{n[(n - \alpha)(\tau n - \tau + 1) + k(\tau + n - 1)]},$$

with equality for the function of the form

$$f(z) = z - \frac{1 - \alpha}{n[(n - \alpha)(\tau n - \tau + 1) + k(\tau + n - 1)]}z^n.$$

COROLLARY 3.2.3. [1] *If $f \in T_\lambda(\alpha)$, then*

$$a_k \leq \frac{1 - \alpha}{(k - \alpha)B_k(\lambda)},$$

with equality for function of the form

$$f(z) = z - \frac{1 - \alpha}{(k - \alpha)B_k(\lambda)} z^k.$$

3.3. GROWTH THEOREM

We now prove the growth theorem for the functions in the class $T[b_k, \beta, m]$.

THEOREM 3.3.1. *If $f \in T[b_k, \beta, m]$, then*

$$r - \frac{\beta}{b_{m+1}} r^{m+1} \leq |f(z)| \leq r + \frac{\beta}{b_{m+1}} r^{m+1}, \quad |z| = r < 1.$$

with equality for

$$(3.3.1) \quad f(z) = z - \frac{\beta}{b_{m+1}} z^{m+1}$$

at $z = r$ for the lower bound and $z = r e^{\frac{i\pi(2p+1)}{m}}$, ($p \in \mathbb{Z}^+$) for the upper bound.

PROOF. For $f \in T[b_k, \beta, m]$, we have

$$\sum_{k=m+1}^{\infty} b_k a_k \leq \beta$$

and since $b_{m+1} \leq b_k$ for $k \geq m + 1$, we have

$$b_{m+1} \sum_{k=m+1}^{\infty} a_k \leq \sum_{k=m+1}^{\infty} b_k a_k \leq \beta$$

or

$$(3.3.2) \quad \sum_{k=m+1}^{\infty} a_k \leq \frac{\beta}{b_{m+1}}.$$

Let $|z| = r$. Since $f = z - \sum_{k=m+1}^{\infty} a_k z^k \in T[b_k, \beta, m]$, we have

$$\begin{aligned} |f(z)| &\leq r + \sum_{k=m+1}^{\infty} a_k r^k \\ &\leq r + r^{m+1} \sum_{k=m+1}^{\infty} a_k. \end{aligned}$$

By using (3.3.2), we obtain

$$|f(z)| \leq r + \frac{\beta}{b_{m+1}} r^{m+1}$$

and similarly

$$|f(z)| \geq r - \frac{\beta}{b_{m+1}} r^{m+1}. \quad \square$$

COROLLARY 3.3.1. [43] *If $f \in \mathcal{U}(k, \tau, \alpha)$, then*

$$r - \frac{1 - \alpha}{2(1 + \tau)(2 + k - \alpha)} r^2 \leq |f(z)| \leq r + \frac{1 - \alpha}{2(1 + \tau)(2 + k - \alpha)} r^2 \quad (|z| = r)$$

with equality for

$$f(z) = z - \frac{1 - \alpha}{2(1 + \tau)(2 + k - \alpha)} z^2.$$

COROLLARY 3.3.2. [13] *If $f \in T_s(\alpha)$, then*

$$r - \frac{1 - \alpha}{2^{s+1} - 2^s \alpha} r^2 \leq |f(z)| \leq r + \frac{1 - \alpha}{2^{s+1} - 2^s \alpha} r^2 \quad (|z| = r)$$

with equality for

$$f(z) = z - \frac{1 - \alpha}{2^{s+1} - 2^s \alpha} z^2.$$

COROLLARY 3.3.3. [1] *If $f \in T_\lambda(\alpha)$, then*

$$r - \frac{1 - \alpha}{(2 - \alpha)(\lambda + 1)} r^2 \leq |f(z)| \leq r + \frac{1 - \alpha}{(2 - \alpha)(\lambda + 1)} r^2 \quad (|z| = r)$$

with equality for

$$f(z) = z - \frac{1 - \alpha}{(2 - \alpha)(\lambda + 1)} z^2.$$

3.4. COVERING THEOREM

THEOREM 3.4.1. *The disk $|z| < 1$ is mapped onto a domain that contains the disk*

$$|w| < 1 - \frac{\beta}{b_{m+1}}$$

by any $f \in T[b_k, \beta, m]$. The result is sharp for the function f given in (3.3.1).

PROOF. The proof follows by letting $r \rightarrow 1$ in Theorem 3.3.1. □

COROLLARY 3.4.1. [44] *The disk $|z| < 1$ is mapped onto a domain that contains the disk*

$$|w| < 1 - \frac{1}{2 - \alpha}$$

by any $f \in TS^(\alpha)$. The theorem is sharp with extremal function $f(z) = z - \frac{1-\alpha}{2-\alpha}z^2$.*

COROLLARY 3.4.2. [44] *The disk $|z| < 1$ is mapped onto a domain that contains the disk*

$$|w| < 1 - \frac{3 - \alpha}{2(2 - \alpha)}$$

by any $f \in TK(\alpha)$. The theorem is sharp with extremal function $f(z) = z - \frac{1-\alpha}{2(2-\alpha)}z^2$.

COROLLARY 3.4.3. [13] *The disk $|z| < 1$ is mapped onto a domain that contains the disk*

$$|w| < 1 - \frac{1 - \alpha}{2^{s+1} - 2^s \alpha}$$

by any $f \in T_s(\alpha)$. The theorem is sharp with extremal function $f(z) = z - \frac{1-\alpha}{2^{s+1}-2^s\alpha}z^2$.

COROLLARY 3.4.4. [1] *The disk $|z| < 1$ is mapped onto a domain that contains the disk*

$$|w| < 1 - \frac{1 - \alpha}{(2 - \alpha)(\lambda + 1)}$$

by any $f \in T_\lambda(\alpha)$. The theorem is sharp with extremal function $f(z) = z - \frac{1-\alpha}{(2-\alpha)(\lambda+1)}z^2$.

3.5. DISTORTION THEOREM

The distortion theorem for the functions in the class $T[b_k, \beta, m]$ is given in the following theorem.

THEOREM 3.5.1. *If $f \in T[b_k, \beta, m]$, then*

$$1 - \frac{\beta(m+1)}{b_{m+1}}r^m \leq |f'(z)| \leq 1 + \frac{\beta(m+1)}{b_{m+1}}r^m, \quad |z| = r < 1$$

where $\frac{b_{m+1}}{m+1} \leq \frac{b_k}{k}$. The result is sharp for the function f given in (3.3.1).

PROOF. For $f \in T[b_k, \beta, m]$, we have

$$\sum_{k=m+1}^{\infty} \frac{b_k}{k}ka_k = \sum_{k=m+1}^{\infty} b_ka_k \leq \beta.$$

Since $\frac{b_{m+1}}{m+1} \leq \frac{b_k}{k}$ for $k \geq m+1$, we obtain

$$(3.5.1) \quad \sum_{k=m+1}^{\infty} \frac{b_{m+1}}{m+1}ka_k \leq \sum_{k=m+1}^{\infty} \frac{b_k}{k}ka_k \leq \beta$$

or equivalently

$$\frac{b_{m+1}}{m+1} \sum_{k=m+1}^{\infty} ka_k \leq \sum_{k=m+1}^{\infty} b_ka_k \leq \beta.$$

Thus we have

$$(3.5.1) \quad \sum_{k=m+1}^{\infty} ka_k \leq \frac{\beta(m+1)}{b_{m+1}}.$$

Since $f(z) = z - \sum_{k=m+1}^{\infty} a_kz^k$, we have

$$f'(z) = 1 - \sum_{k=m+1}^{\infty} ka_kz^{k-1}.$$

Let $|z| = r$, then

$$\begin{aligned} |f'(z)| &\leq 1 + \sum_{k=m+1}^{\infty} ka_kr^{k-1} \\ &\leq 1 + r^m \sum_{k=m+1}^{\infty} ka_k \\ &\leq 1 + \frac{\beta(m+1)}{b_{m+1}}r^m \end{aligned}$$

and similarly

$$|f'(z)| \geq 1 - \frac{\beta(m+1)}{b_{m+1}} r^m.$$

The result is sharp for the function given in (3.3.1). □

COROLLARY 3.5.1. [44] *If $f \in TS^*(\alpha)$ then*

$$1 - \frac{2(1-\alpha)}{2-\alpha} r \leq |f'(z)| \leq 1 + \frac{2(1-\alpha)}{2-\alpha} r, \quad (|z| = r)$$

with equality for

$$f(z) = z - \frac{(1-\alpha)}{2-\alpha} z^2.$$

COROLLARY 3.5.2. [44] *If $f \in TK(\alpha)$ then*

$$1 - \frac{1-\alpha}{2-\alpha} r \leq |f'(z)| \leq 1 + \frac{1-\alpha}{2-\alpha} r, \quad (|z| = r)$$

with equality for

$$f(z) = z - \frac{1-\alpha}{2(2-\alpha)} z^2.$$

COROLLARY 3.5.3. [43] *If $f \in \mathcal{U}(k, \tau, \alpha)$, then*

$$1 - \frac{1-\alpha}{(1+\tau)(2+k-\alpha)} r \leq |f'(z)| \leq 1 + \frac{1-\alpha}{(1+\tau)(2+k-\alpha)} r \quad (|z| = r)$$

with equality for

$$f(z) = z - \frac{1-\alpha}{2(1+\tau)(2+k-\alpha)} z^2.$$

COROLLARY 3.5.4. [13] *If $f \in T_s(\alpha)$, then*

$$1 - \frac{1-\alpha}{2^s - 2^{s-1}\alpha} r \leq |f'(z)| \leq 1 + \frac{1-\alpha}{2^s - 2^{s-1}\alpha} r \quad (|z| = r)$$

with equality for

$$f(z) = z - \frac{1-\alpha}{2^{s+1} - 2^s\alpha} z^2.$$

3.6. CLOSURE THEOREM

Let the functions $F_l(z)$ be given by

$$(3.6.1) \quad F_l(z) = z - \sum_{k=m+1}^{\infty} f_{k,l} z^k, \quad (l = 1, 2, \dots, j).$$

We shall now prove the following closure theorems for the class $T[b_k, \beta, m]$.

THEOREM 3.6.1. *Let the function F_l , defined by (3.6.1), be in the class $T[b_k, \beta, m]$ for every $l = 1, 2, \dots, j$. Then the function $f(z)$ defined by*

$$f(z) = z - \sum_{k=m+1}^{\infty} a_k z^k$$

belongs to the class $T[b_k, \beta, m]$ where $a_k = \frac{1}{j} \sum_{l=1}^j f_{k,l}$ ($k = m+1, m+2, \dots$).

PROOF. Since $F_l \in T[b_k, \beta, m]$, it follows that

$$\sum_{k=m+1}^{\infty} b_k f_{k,l} \leq \beta$$

for every $l = 1, 2, \dots, j$. Hence

$$\begin{aligned} \sum_{k=m+1}^{\infty} b_k a_k &= \sum_{k=m+1}^{\infty} b_k \left(\frac{1}{j} \sum_{l=1}^j f_{k,l} \right) \\ &= \frac{1}{j} \sum_{l=1}^j \left(\sum_{k=m+1}^{\infty} b_k f_{k,l} \right) \\ &\leq \beta. \end{aligned}$$

Therefore it follows that $f(z) \in T[b_k, \beta, m]$ □

THEOREM 3.6.2. *The class $T[b_k, \beta, m]$ is closed under convex linear combination.*

PROOF. Let $F_l(z)$ for $l = 1, 2$ be given by $F_l(z) = z - \sum_{k=m+1}^{\infty} f_{k,l} z^k \in T[b_k, \beta, m]$ and define the function H as

$$H(z) = \gamma F_1(z) + (1 - \gamma) F_2(z) \quad (0 \leq \gamma \leq 1)$$

Since $0 \leq \gamma \leq 1$, it follows that

$$\begin{aligned}
H(z) &= \gamma \left(z - \sum_{k=m+1}^{\infty} f_{k,1} z^k \right) + (1 - \gamma) \left(z - \sum_{k=m+1}^{\infty} f_{k,2} z^k \right) \\
&= \gamma z + z - \gamma z - \sum_{k=m+1}^{\infty} [\gamma f_{k,1} z^k + (1 - \gamma) f_{k,2} z^k] \\
&= z - \sum_{k=m+1}^{\infty} [\gamma f_{k,1} + (1 - \gamma) f_{k,2}] z^k.
\end{aligned}$$

Observe that

$$\begin{aligned}
&\sum_{k=m+1}^{\infty} b_k [\gamma f_{k,1} + (1 - \gamma) f_{k,2}] \\
&= \gamma \sum_{k=m+1}^{\infty} b_k f_{k,1} + (1 - \gamma) \sum_{k=m+1}^{\infty} b_k f_{k,2} \\
&\leq \gamma \beta + (1 - \gamma) \beta \\
&= \beta.
\end{aligned}$$

This shows that $H(z) \in T[b_k, \beta, m]$ and the theorem is proved. □

THEOREM 3.6.3. *Let*

$$F_k(z) = \begin{cases} z, & k=m; \\ z - \frac{\beta}{b_k} z^k, & k=m+1, m+2, \dots \end{cases}$$

Then $f(z) \in T[b_k, \beta, m]$ if and only if $f(z)$ can be expressed in the form

$$f(z) = \sum_{k=m}^{\infty} \delta_k F_k(z),$$

where $\delta_k \geq 0$ and $\sum_{k=m}^{\infty} \delta_k = 1$.

PROOF. Suppose

$$f(z) = \sum_{k=m}^{\infty} \delta_k F_k(z),$$

then we have

$$\begin{aligned}
f(z) &= \delta_m F_m + \sum_{k=m+1}^{\infty} \delta_k F_k(z) \\
&= \delta_m z + \sum_{k=m+1}^{\infty} \delta_k \left(z - \frac{\beta}{b_k} z^k \right) \\
&= \sum_{k=m}^{\infty} \delta_k z - \sum_{k=m+1}^{\infty} \delta_k \frac{\beta}{b_k} z^k \\
(3.6.2) \quad &= z - \sum_{k=m+1}^{\infty} \delta_k \frac{\beta}{b_k} z^k.
\end{aligned}$$

For the function given in (3.6.2), we see that

$$\begin{aligned}
\sum_{k=m+1}^{\infty} \delta_k \frac{\beta}{b_k} b_k &= \sum_{k=m+1}^{\infty} \delta_k \beta \\
&= \beta \left(\sum_{k=m}^{\infty} \delta_k - \delta_m \right) \\
&= \beta(1 - \delta_m).
\end{aligned}$$

Since $\delta_m > 0$, we obtain

$$\sum_{k=m+1}^{\infty} \delta_k \frac{\beta}{b_k} b_k \leq \beta.$$

Thus, $f(z) \in T[b_k, \beta, m]$.

Conversely, suppose $f(z) \in T[b_k, \beta, m]$, then

$$a_k \leq \frac{\beta}{b_k}, \quad (k = m + 1, m + 2, \dots).$$

Therefore we set

$$\delta_k = \frac{b_k}{\beta} a_k, \quad (k = m + 1, m + 2, \dots)$$

and

$$\delta_m = 1 - \sum_{k=m+1}^{\infty} \delta_k.$$

Then we have

$$\begin{aligned}
f(z) &= z - \sum_{k=m+1}^{\infty} \delta_k \frac{\beta}{b_k} z^k \\
&= \sum_{k=m}^{\infty} \delta_k z - \sum_{k=m+1}^{\infty} \delta_k \frac{\beta}{b_k} z^k \\
&= \delta_m z + \sum_{k=m+1}^{\infty} \delta_k \left(z - \frac{\beta}{b_k} z^k \right) \\
&= \delta_m F_m(z) + \sum_{k=m+1}^{\infty} \delta_k F_k(z) \\
&= \sum_{k=m}^{\infty} \delta_k F_k(z).
\end{aligned}$$

This complete the proof. □

COROLLARY 3.6.1. [44] *Let $f_1(z) = z$ and*

$$f_n(z) = z - \frac{1 - \alpha}{n - \alpha} z^n$$

for $n = 2, 3, \dots$. Then $f(z) \in TS^(\alpha)$ if and only if $f(z)$ can be expressed in the form*

$$f(z) = \sum_{n=1}^{\infty} \delta_n f_n(z),$$

where $\delta_n \geq 0$ and $\sum_{n=1}^{\infty} \delta_n = 1$.

COROLLARY 3.6.2. [13] *Let $f_1(z) = z$ and*

$$f_n(z) = z - \frac{1 - \alpha}{n^{s+1} - n^s \alpha} z^n$$

for $n = 2, 3, \dots$. Then $f(z) \in T_s(\alpha)$ if and only if $f(z)$ can be expressed in the form

$$f(z) = \sum_{n=1}^{\infty} \delta_n f_n(z),$$

where $\delta_n > 0$ and $\sum_{n=1}^{\infty} \delta_n = 1$.

COROLLARY 3.6.3. [1] *Let $f_1(z) = z$ and*

$$f_n(z) = z - \frac{1 - \alpha}{(n - \alpha)B_n(\lambda)} z^n$$

for $n = 2, 3, \dots$. Then $f(z) \in T_\lambda(\alpha)$ if and only if $f(z)$ can be expressed in the form

$$f(z) = \sum_{n=1}^{\infty} \delta_n f_n(z),$$

where $\delta_n \geq 0$ and $\sum_{n=1}^{\infty} \delta_n = 1$.

3.7. RADIUS PROBLEM

In this section, we determine the radius of starlikeness of order γ and the radius of convexity of order γ for functions in the class $T[b_k, \beta, m]$. Let us recall that a function f is said to be starlike of order γ if $\Re \frac{zf'(z)}{f(z)} \geq \gamma$ and convex of order γ if $\Re(1 + \frac{zf''(z)}{f'(z)}) \geq \gamma$.

THEOREM 3.7.1. *If $f \in T[b_k, \beta, m]$, then f is starlike of order γ ($0 \leq \gamma < 1$) in $|z| < R$, where*

$$R = \inf_{k \geq m+1} \left[\frac{b_k(1-\gamma)}{\beta(k-\gamma)} \right]^{\frac{1}{k-1}}.$$

PROOF. For $f \in T[b_k, \beta, m]$ we have

$$\begin{aligned} \left| \frac{zf'(z)}{f(z)} - 1 \right| &= \left| \frac{zf'(z) - f(z)}{f(z)} \right| \\ &= \left| \frac{\sum_{k=m+1}^{\infty} (1-k)a_k z^k}{z - \sum_{k=m+1}^{\infty} a_k z^k} \right| \\ &\leq \frac{\sum_{k=m+1}^{\infty} (k-1)a_k |z|^{k-1}}{1 - \sum_{k=m+1}^{\infty} a_k |z|^{k-1}} \end{aligned}$$

Thus f is starlike of order γ if

$$\frac{\sum_{k=m+1}^{\infty} (k-1)a_k |z|^{k-1}}{1 - \sum_{k=m+1}^{\infty} a_k |z|^{k-1}} \leq 1 - \gamma$$

or

$$\begin{aligned} \sum_{k=m+1}^{\infty} (k-1)a_k |z|^{k-1} &\leq (1-\gamma) \left(1 - \sum_{k=m+1}^{\infty} a_k |z|^{k-1} \right) \\ &= 1 - \gamma - \sum_{k=m+1}^{\infty} (1-\gamma)a_k |z|^{k-1}. \end{aligned}$$

Then

$$\sum_{k=m+1}^{\infty} [(k-1) + (1-\gamma)] a_k |z|^{k-1} \leq 1 - \gamma.$$

Thus we obtain

$$(3.7.1) \quad \sum_{k=m+1}^{\infty} \frac{(k-\gamma)}{1-\gamma} a_k |z|^{k-1} \leq 1.$$

The inequality (3.7.1) holds if

$$\sum_{k=m+1}^{\infty} \frac{(k-\gamma)}{1-\gamma} a_k |z|^{k-1} \leq \sum_{k=m+1}^{\infty} \frac{b_k}{\beta} a_k$$

or if

$$|z|^{k-1} \leq \frac{b_k}{\beta} \frac{1-\gamma}{(k-\gamma)} \quad (k = m+1, m+2, \dots)$$

and this proves the result. \square

THEOREM 3.7.2. *If $f \in T[b_k, \beta, m]$, then f is convex of order γ ($0 \leq \gamma < 1$) in $|z| < R$, where*

$$R = \inf_{k \geq m+1} \left[\frac{b_k(1-\gamma)}{\beta k(k-\gamma)} \right]^{\frac{1}{k-1}}.$$

PROOF. For $f \in T[b_k, \beta, m]$ we have

$$\begin{aligned} \left| \frac{z f''(z)}{f'(z)} \right| &= \left| \frac{-\sum_{k=m+1}^{\infty} k(k-1) a_k z^{k-1}}{1 - \sum_{k=m+1}^{\infty} k a_k z^{k-1}} \right| \\ &\leq \frac{\sum_{k=m+1}^{\infty} k(k-1) a_k |z|^{k-1}}{1 - \sum_{k=m+1}^{\infty} k a_k |z|^{k-1}} \end{aligned}$$

Thus f is convex of order γ if

$$\frac{\sum_{k=m+1}^{\infty} k(k-1) a_k |z|^{k-1}}{1 - \sum_{k=m+1}^{\infty} k a_k |z|^{k-1}} \leq 1 - \gamma$$

or

$$\begin{aligned} \sum_{k=m+1}^{\infty} k(k-1) a_k |z|^{k-1} &\leq (1-\gamma) \left(1 - \sum_{k=m+1}^{\infty} k a_k |z|^{k-1} \right) \\ &\leq 1 - \gamma - \sum_{k=m+1}^{\infty} k(1-\gamma) a_k |z|^{k-1}. \end{aligned}$$

Then

$$\sum_{k=m+1}^{\infty} k[(k-1) + (1-\gamma)]a_k|z|^{k-1} \leq 1 - \gamma.$$

Hence we obtain

$$(3.7.2) \quad \sum_{k=m+1}^{\infty} \frac{k(k-\gamma)}{1-\gamma} a_k |z|^{k-1} \leq 1.$$

The inequality (3.7.2) follows if

$$\sum_{k=m+1}^{\infty} \frac{k(k-\gamma)}{1-\gamma} a_k |z|^{k-1} \leq \sum_{k=m+1}^{\infty} \frac{b_k}{\beta} a_k$$

or if

$$|z|^{k-1} \leq \frac{b_k}{\beta} \frac{1-\gamma}{k(k-\gamma)} \quad (k = m+1, m+2, \dots)$$

and this completes the proof. □

CHAPTER 4

RADIUS PROBLEMS FOR SOME CLASSES OF ANALYTIC FUNCTIONS

4.1. MOTIVATION AND PRELIMINARIES

Lets recall the class $S^*(\phi)$ which was introduced by Ma and Minda [17]. The class $S^*(\phi)$ denote the class of functions f in \mathcal{S} for which

$$\frac{zf'(z)}{f(z)} \prec \phi(z), \quad z \in U,$$

where $\phi(z)$ is an analytic function with positive real part in U , $\phi(0) = 1$, $\phi'(0) > 0$, and ϕ maps U onto a region starlike with respect to 1 and symmetric with respect to the real axis. The class $S^*(\alpha)$ consisting of starlike functions of order α , $0 \leq \alpha < 1$ and the class $S^*[A, B]$ of Janowski starlike functions are special cases of $S^*(\phi)$ when $\phi(z) := (1+(1-2\alpha)z)/(1-z)$ and $\phi(z) := (1+Az)/(1+Bz)$ ($-1 \leq B < A \leq 1$) respectively.

The function $f(z) \in \mathcal{S}$ is uniformly convex [11] if for every circular arc γ contained in U with center $\zeta \in U$, the image arc $f(\gamma)$ is convex. Denote the class of all uniformly convex functions by UCV . Ma and Minda [16] and Ronning [33], independently showed that the function $f(z)$ is uniformly convex if and only if

$$\Re \left\{ 1 + \frac{zf''(z)}{f'(z)} \right\} > \left| \frac{zf''(z)}{f'(z)} \right| \quad (z \in U).$$

A corresponding class S_p consisting of parabolic starlike functions f , where $f(z) = zg'(z)$ for g in UCV , was introduced in [33]. A function f is in S_p if and only if

$$\Re \left\{ \frac{zf'(z)}{f(z)} \right\} > \left| \frac{zf'(z)}{f(z)} - 1 \right| \quad (z \in U).$$

A survey of these functions can be found in [30], while radius problems associated with the classes UCV and S_p can be found in [10, 28, 32, 42, 29].

In this chapter, several radius problems associated with the class $S^*[A, B]$ defined by

$$S^*[A, B] := \left\{ f \in \mathcal{A} : \frac{zf'(z)}{f(z)} \prec \frac{1 + Az}{1 + Bz}, \quad (A, B \in \mathcal{C}, A \neq B, |B| \leq 1; z \in U) \right\}$$

are investigated. Several known results relating to radii problems are shown to be simple consequences of the results obtained here. Unless explicitly stated otherwise, it is assumed that the complex constants A and B satisfy $A \neq B$ and $|B| \leq 1$.

For $A = 1 - 2\alpha$, $\alpha > 1$ and $B = -1$, denote the class $S^*[1 - 2\alpha, -1]$ by $M(\alpha)$. Equivalently, $M(\alpha)$ can be expressed in the form

$$M(\alpha) := \left\{ f \in \mathcal{A} : \Re \left(\frac{zf'(z)}{f(z)} \right) < \alpha, \quad (z \in U) \right\}.$$

The class $M(\alpha)$ was investigated by Uralegaddi *et al.* [47], while a subclass of $M(\alpha)$ was investigated by Owa and Srivastava [21]. Since functions in the class $M(\alpha)$ and in general $S^*[A, B]$ need not be starlike, in this paper, we find the radius of starlikeness for functions in these classes, as well as the radius of strong-starlikeness and parabolic-starlikeness.

4.2. RADIUS OF STARLIKENESS OF ORDER α

The Lindelöf principle or subordination principle states that if $f \prec g$, then $f(U_r) \subset g(U_r)$, where U_r is the disk $|z| \leq r < 1$. This principle is used to prove the following lemma.

LEMMA 4.2.1. *Let $p(z)$ be analytic in U . Then*

$$p(z) \prec \frac{1 + Az}{1 + Bz}$$

if and only if

$$(4.2.1) \quad \left| p(z) - \frac{1 - A\bar{B}r^2}{1 - |B|^2r^2} \right| \leq \frac{|B - A|r}{1 - |B|^2r^2} \quad (|z| \leq r < 1).$$

The result is sharp for $|z| = r$.

PROOF. Let the function w be defined by

$$w(z) = \frac{1 + Az}{1 + Bz}.$$

Then we have

$$w(z) + Bzw(z) = 1 + Az.$$

Further

$$|w(z) - 1| = |z(A - Bw(z))|$$

implies

$$|w(z) - 1| \leq r|A - Bw(z)|$$

where $|z| \leq r < 1$. Then

$$|w(z) - 1|^2 \leq r^2|A - Bw(z)|^2$$

which is equivalent to

$$|w(z)|^2 - 2\Re w(z) + 1 \leq r^2[|A|^2 + |B|^2|w(z)|^2 - 2\Re A\bar{B}w(z)].$$

Hence

$$|w(z)|^2(1 - |B|^2r^2) - 2\Re[w(z)(1 - A\bar{B}r^2)] \leq |A|^2r^2 - 1$$

or

$$|w(z)|^2 - 2\Re \left[w(z) \left(\frac{1 - A\bar{B}r^2}{1 - |B|^2r^2} \right) \right] \leq \frac{|A|^2r^2 - 1}{1 - |B|^2r^2}.$$

Then we have

$$|w(z)|^2 - 2\Re \left[w(z) \left(\frac{1 - A\bar{B}r^2}{1 - |B|^2r^2} \right) \right] + \left| \frac{1 - A\bar{B}r^2}{1 - |B|^2r^2} \right|^2 \leq \frac{|A|^2r^2 - 1}{1 - |B|^2r^2} + \left| \frac{1 - A\bar{B}r^2}{1 - |B|^2r^2} \right|^2,$$

which gives

$$\begin{aligned} \left| w(z) - \frac{1 - A\bar{B}r^2}{1 - |B|^2r^2} \right|^2 &\leq \frac{(|A|^2r^2 - 1)(1 - |B|^2r^2) + |1 - A\bar{B}r^2|^2}{(1 - |B|^2r^2)^2} \\ &\leq \frac{|A|^2r^2 + |B|^2r^2 - 2\Re A\bar{B}r^2}{(1 - |B|^2r^2)^2} \\ &\leq \frac{|B - A|^2r^2}{(1 - |B|^2r^2)^2}. \end{aligned}$$

Therefore,

$$\left| w(z) - \frac{1 - A\bar{B}r^2}{1 - |B|^2r^2} \right| \leq \frac{|B - A|r}{1 - |B|^2r^2}.$$

Hence by using the subordination principle we have

$$p(z) \prec \frac{1 + Az}{1 + Bz}$$

if and only if

$$\left| p(z) - \frac{1 - A\bar{B}r^2}{1 - |B|^2r^2} \right| \leq \frac{|B - A|r}{1 - |B|^2r^2} \quad (|z| \leq r < 1). \quad \square$$

THEOREM 4.2.1. *Let $0 \leq \alpha < 1$. If $f \in S^*[A, B]$, then the function f is starlike of order α in $|z| \leq R(\alpha)$ where*

$$R(\alpha) := \min \left\{ \frac{2(1 - \alpha)}{|B - A| + |(2\alpha - 1)B - A|}, 1 \right\}.$$

The result is sharp for the function $f(z) \in \mathcal{A}$ given by

$$(4.2.2) \quad f(z) = \begin{cases} z(1 + Bz)^{(A-B)/B} & \text{if } B \neq 0 \\ z \exp Az & \text{if } B = 0. \end{cases}$$

PROOF. Since $f(z) \in S^*[A, B]$, it follows that

$$\frac{zf'(z)}{f(z)} \prec \frac{1 + Az}{1 + Bz}$$

and with $p(z) = \frac{zf'(z)}{f(z)}$, Lemma 4.2.1 gives

$$(4.2.3) \quad \Re \frac{zf'(z)}{f(z)} \geq \Re \left(\frac{1 - A\bar{B}r^2}{1 - |B|^2r^2} \right) - \frac{|B - A|r}{1 - |B|^2r^2}.$$

Also if $f(z)$ is starlike of order α , then

$$(4.2.4) \quad \Re \frac{zf'(z)}{f(z)} > \alpha.$$

From (4.2.3) and (4.2.4), we see that

$$\Re \frac{zf'(z)}{f(z)} \geq \Re \left(\frac{1 - A\bar{B}r^2}{1 - |B|^2r^2} \right) - \frac{|B - A|r}{1 - |B|^2r^2} \geq \alpha$$

is true if

$$1 - \Re A\bar{B}r^2 - |B - A|r \geq \alpha(1 - |B|^2r^2)$$

or

$$(\Re A\bar{B} - \alpha|B|^2)r^2 + |B - A|r + \alpha - 1 \leq 0.$$

Therefore,

$$r = R_\alpha = \frac{-|B - A| \pm \sqrt{|B - A|^2 - 4(\Re A\bar{B} - \alpha|B|^2)(\alpha - 1)}}{2(\Re A\bar{B} - \alpha|B|^2)}.$$

By taking the conjugate and simplifying it, we get

$$R_\alpha = \frac{2(1 - \alpha)}{|B - A| + |(2\alpha - 1)B - A|}.$$

Therefore the $S^*(\alpha)$ -radius for $f \in S^*[A, B]$ is $R(\alpha) = \min \{R_\alpha, 1\}$.

To get the extremal function, let

$$\begin{aligned} \frac{zf'(z)}{f(z)} &= \frac{1 + Az}{1 + Bz} \\ &= 1 + \frac{(A - B)z}{1 + Bz}. \end{aligned}$$

For $B \neq 0$, we have

$$\frac{f'(z)}{f(z)} - \frac{1}{z} = \frac{A - B}{1 + Bz}$$

or

$$\log \frac{f}{z} = \frac{A - B}{B} \log(1 + Bz).$$

Therefore,

$$f(z) = z(1 + Bz)^{\frac{A-B}{B}}.$$

For $B = 0$,

$$\frac{f'(z)}{f(z)} - \frac{1}{z} = 1 + A.$$

Integrating both sides, we get

$$\log \frac{f}{z} = Az$$

and hence

$$f(z) = z \exp(Az).$$

The result is sharp for the functions obtained here. \square

The case when A and B are real numbers yields the following corollary:

COROLLARY 4.2.1. [27] *Let $A, B \in \mathcal{R}$, $A < B$, $|B| \leq 1$ and $0 \leq \alpha < 1$. If $f \in S^*[A, B]$, then the function f belongs to $S^*(\alpha)$ in $|z| \leq R(\alpha)$ where*

$$R(\alpha) = \begin{cases} 1 & (\frac{1+A}{1+B} \geq \alpha) \\ \frac{1-\alpha}{\alpha B - A} & (\frac{1+A}{1+B} \leq \alpha). \end{cases}$$

THEOREM 4.2.2. *Let $\alpha > 1$. If $f \in S^*[A, B]$, then the function f is in $M(\alpha)$ for $|z| \leq R_M(\alpha)$ where*

$$R_M(\alpha) := \min \left\{ \frac{2(\alpha - 1)}{|B - A| + |(2\alpha - 1)B - A|}, 1 \right\}.$$

The result is sharp for the function $f(z) \in \mathcal{A}$ given by (4.2.2).

PROOF. Since $f \in S^*[A, B]$, Lemma 4.2.1 gives

$$(4.2.5) \quad \Re \frac{zf'(z)}{f(z)} \leq \Re \left(\frac{1 - A\bar{B}r^2}{1 - |B|^2r^2} \right) + \frac{|B - A|r}{1 - |B|^2r^2}$$

and $f(z) \in M(\alpha)$ if

$$(4.2.6) \quad \Re \frac{zf'(z)}{f(z)} < \alpha.$$

From (4.2.5) and (4.2.6), we see that

$$\Re \frac{zf'(z)}{f(z)} \leq \Re \left(\frac{1 - A\bar{B}r^2}{1 - |B|^2r^2} \right) + \frac{|B - A|r}{1 - |B|^2r^2} \leq \alpha$$

is true if

$$1 - \alpha + |B - A|r - (\Re A\bar{B} - \alpha|B|^2)r^2 \leq 0.$$

Solving the quadratic inequality yields

$$r = R_M = \frac{2(\alpha - 1)}{|B - A| + |(2\alpha - 1)B - A|}.$$

The result is sharp for the function given in (4.2.2). \square

COROLLARY 4.2.2. [27] Let $A, B \in \mathcal{R}$, $A < B < 1$, $|B| \leq 1$ and $\alpha > 1$. If $f \in S^*[A, B]$, then the function f is in $M(\alpha)$ for $|z| \leq R_M(\alpha)$ where

$$R_M(\alpha) = \begin{cases} 1 & (\frac{1-A}{1-B} \leq \alpha) \\ \frac{\alpha-1}{\alpha B-A} & (\frac{1-A}{1-B} \geq \alpha). \end{cases}$$

THEOREM 4.2.3. Let $A, B, C, D \in \mathcal{C}$, $A \neq B$, $|B| \leq 1$ and $D \neq C$, $|D| \leq 1$. If $f \in S^*[C, D]$, then the $S^*[A, B]$ -radius $R_{[A,B]}$ of f is given by

$$R_{[A,B]} = \min \left\{ \frac{|A-B|}{|C-D| + |AD-BC|}, 1 \right\}.$$

PROOF. Let $P(z) = (1 + Az)/(1 + Bz)$ and $Q(z) = (1 + Cz)/(1 + Dz)$. Since

$$\frac{zf'(z)}{f(z)} \prec Q(z),$$

the $S^*[A, B]$ -radius is determined from the number R such that $0 < R \leq 1$ and $Q(Rz) \prec P(z)$ for z in U . Let $H(z) = P^{-1}(Q(z))$. We know that

$$P^{-1}(z) = \frac{z-1}{A-Bz}$$

and

$$P^{-1}(Q(z)) = \frac{\frac{1+Cz}{1+Dz} - 1}{A - B\left(\frac{1+Cz}{1+Dz}\right)}.$$

Hence

$$H(z) = \frac{(C-D)z}{(A-B) + (AD-BC)z}$$

and

$$|H(Rz)| \leq \frac{|C-D|R}{|A-B| - |AD-BC|R} \leq 1$$

for

$$|z| = R \leq \frac{|A-B|}{|C-D| + |AD-BC|}. \quad \square$$

COROLLARY 4.2.3. Let $A, B, C, D \in \mathcal{C}$, $A \neq B$, $|B| \leq 1$ and $D \neq C$, $|D| \leq 1$. Then the class $S^*[C, D]$ is a subclass of $S^*[A, B]$ if and only if

$$|AD-BC| \leq |A-B| - |C-D|.$$

The above corollary is an extension of the fact that $S^*(\alpha) \subset S^*(\beta)$ if and only if $\alpha \geq \beta$.

4.3. RADIUS OF STRONG STARLIKENESS

A function $f(z) \in \mathcal{A}$ is strongly starlike of order γ , $0 < \gamma \leq 1$ if

$$\frac{zf'(z)}{f(z)} \prec \left(\frac{1+z}{1-z} \right)^\gamma.$$

This is equivalent to the condition

$$\left| \arg \left\{ \frac{zf'(z)}{f(z)} \right\} \right| \leq \frac{\pi}{2} \gamma.$$

In other words the values of $zf'(z)/f(z)$ are in the same sector of $|y| \leq \tan(\gamma\pi/2)x$, $x \geq 0$. In this section we compute the radius of strong starlikeness for the class $S^*[A, B]$. We will use the following lemma:

LEMMA 4.3.1. [10] *If $R_a \leq (\Re a) \sin(\pi\gamma/2) - (\Im a) \cos(\pi\gamma/2)$, $\Im a \geq 0$, for $a \in \mathcal{C}$, then the disk $|w - a| \leq R_a$ is contained in the sector $|\arg w| \leq \pi\gamma/2$, $0 < \gamma \leq 1$.*

THEOREM 4.3.1. *Let $0 < \gamma \leq 1$ and $\Im(A\bar{B}) \leq 0$. If $f \in S^*[A, B]$, then the function f is strongly starlike of order γ in $|z| < R(\gamma)$ where $R(\gamma) = \min\{1, R_\gamma\}$, and*

$$R_\gamma = \frac{2 \sin(\pi\gamma/2)}{|B - A| + [|B - A|^2 + 4 \sin^2(\pi\gamma/2) \Re(A\bar{B}) - 4 \cos(\pi\gamma/2) \sin(\pi\gamma/2) \Im(A\bar{B})]^{1/2}}.$$

PROOF. Lemma 4.2.1 yields

$$\left| \frac{zf'(z)}{f(z)} - a \right| \leq R_a,$$

where

$$a = \frac{1 - A\bar{B}r^2}{1 - |B|^2r^2} \quad \text{and} \quad R_a = \frac{|B - A|r}{1 - |B|^2r^2}.$$

Since the $\Im(A\bar{B}) \leq 0$, it follows from Lemma 4.3.1 that

$$\frac{|B - A|r}{1 - |B|^2 r^2} \leq \frac{1 - \Re(A\bar{B})r^2}{1 - |B|^2 r^2} \sin\left(\frac{\pi\gamma}{2}\right) + \frac{\Im(A\bar{B})r^2}{1 - |B|^2 r^2} \cos\left(\frac{\pi\gamma}{2}\right)$$

or

$$\left[\Im(A\bar{B}) \cos\left(\frac{\pi\gamma}{2}\right) - \Re(A\bar{B}) \sin\left(\frac{\pi\gamma}{2}\right) \right] r^2 - |B - A|r + \sin\left(\frac{\pi\gamma}{2}\right) \geq 0.$$

Since $\sin(\frac{\pi\gamma}{2}) \geq 0$, the above quadratic inequality yields

$$r = \frac{|B - A| \pm \sqrt{|B - A|^2 - 4[\Im(A\bar{B}) \cos(\pi\gamma/2) - \Re(A\bar{B}) \sin(\pi\gamma/2)][\sin(\pi\gamma/2)]}}{2[\Im(A\bar{B}) \cos(\pi\gamma/2) - \Re(A\bar{B}) \sin(\pi\gamma/2)]}.$$

By taking the conjugate and simplifying it, we get

$$r = R_\gamma = \frac{2 \sin(\pi\gamma/2)}{|B - A| + [|B - A|^2 + 4 \sin^2(\pi\gamma/2) \Re(A\bar{B}) - 4 \cos(\pi\gamma/2) \sin(\pi\gamma/2) \Im(A\bar{B})]^{1/2}}.$$

Thus f is strongly starlike of order γ in $|z| < R(\gamma)$ where $R(\gamma) = \min\{1, R_\gamma\}$. \square

Silvia [45] defined the class $SP(\alpha, A, B)$ consisting of functions f in \mathcal{A} satisfying

$$e^{i\alpha} \frac{zf'(z)}{f(z)} \prec \cos \alpha \frac{1 + Az}{1 + Bz} + i \sin \alpha, \quad z \in U,$$

with $0 \leq \alpha < 1$, $-1 \leq B < A \leq 1$.

COROLLARY 4.3.1. *Let $f \in SP(\alpha, A, B)$, $-1 \leq B < A \leq 1$, $0 \leq \alpha < 1$, $0 < \rho \leq 1$, and $B \sin 2\alpha \leq 0$. Then the function f is strongly starlike of order ρ in $|z| < R(\rho)$ where $R(\rho) = \min\{1, R_\rho\}$, and*

$$R_\rho = \frac{2 \sin \delta}{(A - B) \cos \alpha + \sqrt{(A - B)^2 \cos^2 \alpha + 4B^2 \sin^2 \delta + 4B(A - B) \sin \delta \cos \alpha \sin(\alpha + \delta)}}$$

where $\delta = \pi\rho/2$.

PROOF. The function f is in $SP(\alpha, A, B)$ if

$$e^{i\alpha} \frac{zf'(z)}{f(z)} \prec \cos \alpha \frac{1 + Az}{1 + Bz} + i \sin \alpha,$$

or

$$\frac{zf'(z)}{f(z)} \prec e^{-i\alpha} \left[\cos \alpha \frac{1 + Az}{1 + Bz} + i \sin \alpha \right].$$

Therefore,

$$\frac{zf'(z)}{f(z)} \prec \frac{1 + (A \cos \alpha + iB \sin \alpha)e^{-i\alpha}z}{1 + Bz}.$$

Replacing A with $(A \cos \alpha + Bi \sin \alpha)e^{-i\alpha}$ in Theorem 4.3.1 yields

$$R_\rho = \frac{2 \sin \delta}{(A - B) \cos \alpha + \sqrt{(A - B)^2 \cos^2 \alpha + 4B^2 \sin^2 \delta + 4B(A - B) \sin \delta \cos \alpha \sin(\alpha + \delta)}}$$

where $\delta = \pi\rho/2$. □

REMARK 4.3.1. Corollary 4.3.1 was also obtained by Gangadharan et al.[10].

However there was a slight mistake in their result. Refer [46].

4.4. RADIUS OF PARABOLIC STARLIKENESS

In [31] the class S_p of parabolic starlike functions was generalized by introducing a parameter β , $-1 \leq \beta < 1$. The class $S_p(\beta)$ is a subclass of \mathcal{A} consisting of functions $f \in \mathcal{A}$ satisfying

$$\left| \frac{zf'(z)}{f(z)} - 1 \right| < \Re \left\{ \frac{zf'(z)}{f(z)} \right\} - \beta \quad (|z| < 1).$$

Observe that the values of the functional $\frac{zf'(z)}{f(z)}$ lies in the parabolic region

$$(4.4.1) \quad \Omega := \left\{ w = u + iv : v^2 < 2(1 - \beta) \left(u - \frac{\beta + 1}{2} \right) \right\}.$$

In this section, the $S_p(\beta)$ -radius of functions in $S^*[A, B]$ for $A, B \in \mathcal{R}$, $A < B$ and $|B| \leq 1$ is determined.

THEOREM 4.4.1. *Let $\beta < 1$, $A < B$, and $|B| \leq 1$. Let R_1 be given by*

$$R_1 := \min \left\{ 1, \frac{2(1 - \beta)}{B - A + \sqrt{(B - A)^2 + 4B^2(1 - \beta)^2}} \right\},$$

R_2 be the largest number in $(0, 1]$ such that $1 \geq (B(1 + \beta) - 2A)r + \beta$ for all $r \in [0, R_2]$, and R_3 be the largest number in $(0, 1]$ such that $A + B(1 - 2\beta) \geq 2B^3(1 - \beta)r^2$ for all $r \in [0, R_3]$. If $f \in S^*[A, B]$, then f satisfies

$$\Re \left\{ \frac{zf'(z)}{f(z)} \right\} > \left| \frac{zf'(z)}{f(z)} - 1 \right| + \beta \quad (|z| < R)$$

where

$$R = \begin{cases} R_2 & \text{if } R_2 \leq R_1 \\ R_3 & \text{if } R_2 > R_1. \end{cases}$$

PROOF. Since

$$\frac{zf'(z)}{f(z)} \prec \frac{1 + Az}{1 + Bz},$$

it follows from Lemma 4.2.1 that

$$(4.4.2) \quad \left| \frac{zf'(z)}{f(z)} - \frac{1 - ABr^2}{1 - B^2r^2} \right| \leq \frac{(B - A)r}{1 - B^2r^2}, \quad (|z| \leq r < 1).$$

By letting $w(z) = \frac{zf'(z)}{f(z)} = u + iv$, the points on the boundary of the disk in (4.4.2) are given by

$$w(z) = \frac{1 - ABr^2}{1 - B^2r^2} + \frac{(B - A)r}{1 - B^2r^2} e^{i\theta}$$

and hence we have

$$(4.4.3) \quad \Re w(z) = \frac{(1 - ABr^2) + (B - A)r \cos \theta}{1 - B^2r^2}, \quad \Im w(z) = \frac{(B - A)r \sin \theta}{1 - B^2r^2}.$$

For $f \in S_p(\beta)$, we have

$$\Re \frac{zf'(z)}{f(z)} > \left| \frac{zf'(z)}{f(z)} - 1 \right| + \beta$$

or equivalently

$$(4.4.4) \quad u > |(u + iv) - 1| + \beta.$$

Squaring (4.4.4) and rewriting yields

$$v^2 < 2u(1 - \beta) + \beta^2 - 1$$

or

$$(4.4.5) \quad (\Im w(z))^2 < 2(1 - \beta) \left(\Re w(z) - \frac{1 + \beta}{2} \right).$$

Substituting (4.4.3) into (4.4.5) gives

$$\left[\frac{(B - A)r \sin \theta}{1 - B^2r^2} \right]^2 < 2 \left[\frac{1 - ABr^2 + (B - A)r \cos \theta}{1 - B^2r^2} \right] (1 - \beta) + \beta^2 - 1.$$

Letting $x = \cos \theta$ and simplifying leads to

$$T(x) := (B - A)^2 r^2 x^2 + 2(1 - \beta)(B - A)(1 - B^2 r^2)rx + 2(1 - \beta)(1 - B^2 r^2)(1 - AB r^2) - (1 - \beta^2)(1 - B^2 r^2)^2 - (B - A)^2 r^2 \geq 0.$$

We need to find $r = R$ such that $T(x) \geq 0$ for all $x \in [-1, 1]$.

Since

$$T'(x) = 2(B - A)^2 r^2 x + 2(1 - \beta)(B - A)(1 - B^2 r^2)r$$

we see that $T'(x) = 0$ for

$$x = x_0 = -\frac{(1 - \beta)(1 - B^2 r^2)}{(B - A)r}.$$

Since $\beta < 1$, $A < B$ and $|B| \leq 1$, we have $x_0 < 0$. If $x_0 \leq -1$, we require $T(-1) \geq 0$ and if $-1 < x_0 < 0$, we require $T(x_0) \geq 0$. Note that $x_0 \leq -1$ implies

$$1 - B^2 r^2 - \beta + \beta B^2 r^2 \geq (B - A)r$$

which is equivalent to

$$R_1 = r \leq \frac{2(1 - \beta)}{B - A + \sqrt{(B - A)^2 + 4B^2(1 - \beta)}}.$$

The condition $T(-1) \geq 0$ is equivalent to

$$(B - A)^2 r^2 - 2(1 - \beta)(B - A)(1 - B^2 r^2)r + 2(1 - \beta)(1 - B^2 r^2)(1 - AB r^2) - (1 - \beta^2)(1 - B^2 r^2)^2 - (B - A)^2 r^2 \geq 0$$

or

$$2(1 - \beta)[-(B - A)r + (1 - AB r^2)] \geq (1 - \beta^2)(1 - B^2 r^2).$$

Therefore

$$1 \geq (B(1 + \beta) - 2A)r + \beta.$$

Also, $T(x_0) \geq 0$ implies

$$(B - A)^2 r^2 \left(-\frac{(1 - \beta)(1 - B^2 r^2)}{(B - A)r} \right) + 2(1 - \beta)(B - A)(1 - B^2 r^2)r \left(-\frac{(1 - \beta)(1 - B^2 r^2)}{(B - A)r} \right) + 2(1 - \beta)(1 - B^2 r^2)(1 - AB r^2) - (1 - \beta^2)(1 - B^2 r^2)^2 - (B - A)^2 r^2 \geq 0$$

which yields

$$A + B(1 - 2\beta) \geq 2B^3(1 - \beta)r^2.$$

For $A, B \in \mathcal{R}$, $A < B$ and $|B| \leq 1$, let R_1 , R_2 and R_3 be as in the hypothesis. If $R_2 \leq R_1$, then the disk (4.4.2) will be inside the parabolic region (4.4.1) if and only if $r \leq R_2$. If $R_2 > R_1$, then the disk (4.4.2) will be inside the parabolic region (4.4.1) if and only if $r \leq R_3$. This completes our proof. \square

In the special case $\beta = 0$, the following results is obtained:

COROLLARY 4.4.1. [27] *For $A, B \in \mathcal{R}$, $A < B$ and $|B| \leq 1$, let R_1 be given by*

$$R_1 := \min \left\{ 1, \frac{2}{B - A + \sqrt{(B - A)^2 + 4B^2}} \right\}.$$

and let R_2 be the largest number in $(0, 1]$ such that $1 \geq (B - 2A)r$ for all $r \in [0, R_2]$

and R_3 be the largest number in $(0, 1]$ such that $A + B \geq 2B^2r^2$ for all $r \in [0, R_3]$.

If $f \in S^[A, B]$, then the S_p -radius is given by*

$$R = \begin{cases} R_2 & \text{if } R_2 \leq R_1 \\ R_3 & \text{if } R_2 > R_1. \end{cases}$$

CONCLUSION

- A modest attempt has been made in this thesis to introduce and study certain classes of univalent functions defined on the unit disc U .
- Several subclasses of meromorphic functions are introduced by means of convolution. Class relation as well as inclusion and convolution properties are established.
- Coefficient estimate, growth theorem, covering theorem, distortion theorem, closure theorems and radius properties are obtained for a generalized subclass of univalent functions with negative coefficients.
- Radius of starlikeness of order α , radius of strong starlikeness and radius of parabolic starlikeness are determined for the class $S^*[A, B]$ where A and B are complex constants.
- **Open Problem:** An analytic convex function in U is necessarily starlike or equivalently $K \subset S^*$. For the meromorphic subclass, is it true that the class $\mathcal{M}_g^k(h) \subset \mathcal{M}_g^s(h)$?

REFERENCES

- [1] O. P. Ahuja, Hadamard products of analytic functions defined by Ruscheweyh derivatives, in *Current topics in analytic function theory*, 13–28, World Sci. Publ., River Edge, NJ, 1992.
- [2] J. W. Alexander, Functions which map the interior of the unit circle upon simple regions, *Ann. of Math. (2)* **17** (1915), no. 1, 12–22.
- [3] R.M. Ali; V. Ravichandran; S.K. Lee, Subclasses of multivalent starlike and convex functions, *Bull. Belgian Math. Soc. Simon Stevin.* (to appear)
- [4] H. S. Al-Amiri, Prestarlike functions of order α and type β with negative coefficients, *Ann. Univ. Mariae Curie-Skłodowska Sect. A* **39** (1985), 1–11 (1988).
- [5] R. W. Barnard and C. Kellogg, Applications of convolution operators to problems in univalent function theory, *Michigan Math. J.* **27** (1980), no. 1, 81–94.
- [6] L. de Branges, A proof of the Bieberbach conjecture, *Acta Math.* **154** (1985), no. 1-2, 137–152.
- [7] P. L. Duren, *Univalent functions*, Springer, New York, 1983.
- [8] P. Eenigenburg, S. S. Miller, P. T. Mocanu and M. O. Reade, On a Briot-Bouquet differential subordination, *Rev. Roumaine Math. Pures Appl.* **29** (1984), no. 7, 567–573.
- [9] B. A. Frasin, Generalization of partial sums of certain analytic and univalent functions, *Appl. Math. Lett.* **21** (2008), no. 7, 735–741.
- [10] A. Gangadharan, V. Ravichandran and T. N. Shanmugam, Radii of convexity and strong starlikeness for some classes of analytic functions, *J. Math. Anal. Appl.* **211** (1997), no. 1, 301–313.
- [11] A. W. Goodman, On uniformly convex functions, *Ann. Polon. Math.* **56** (1991), no. 1, 87–92.
- [12] A. W. Goodman, *Univalent functions. Vol. I*, Mariner, Tampa, FL, 1983.
- [13] E. Kadioğlu, On subclass of univalent functions with negative coefficients, *Appl. Math. Comput.* **146** (2003), no. 2-3, 351–358.
- [14] M. Kamali and S. Akbulut, On a subclass of certain convex functions with negative coefficients, *Appl. Math. Comput.* **145** (2003), no. 2-3, 341–350.
- [15] S. Kanas and A. Wisniowska, Conic regions and k -uniform convexity, *J. Comput. Appl. Math.* **105** (1999), no. 1-2, 327–336.
- [16] W. C. Ma and D. Minda, Uniformly convex functions, *Ann. Polon. Math.* **57** (1992), no. 2, 165–175.
- [17] W. C. Ma and D. Minda, A unified treatment of some special classes of univalent functions, in *Proceedings of the Conference on Complex Analysis (Tianjin, 1992)*, 157–169, Int. Press, Cambridge, MA.
- [18] S. Owa, An application of the Ruscheweyh derivatives. II, *Publ. Inst. Math. (Beograd) (N.S.)* **38(52)** (1985), 99–110.
- [19] S. Owa, On new classes of analytic functions with negative coefficients, *Internat. J. Math. Math. Sci.* **7** (1984), no. 4, 719–730.
- [20] S. Owa and J. Nishiwaki, Coefficient estimates for certain classes of analytic functions, *JIPAM. J. Inequal. Pure Appl. Math.* **3** (2002), no. 5, Article 72, 5 pp. (electronic).
- [21] S. Owa and H. M. Srivastava, Some generalized convolution properties associated with certain subclasses of analytic functions, *JIPAM. J. Inequal. Pure Appl. Math.* **3** (2002), no. 3, Article 42, 13 pp. (electronic).
- [22] K. S. Padmanabhan and R. Parvatham, Some applications of differential subordination, *Bull. Austral. Math. Soc.* **32** (1985), no. 3, 321–330.
- [23] R. Parvatham and S. Radha, On α -starlike and α -close-to-convex functions with respect to n -symmetric points, *Indian J. Pure Appl. Math.* **17** (1986), no. 9, 1114–1122.

- [24] G. Pólya and I. J. Schoenberg, Remarks on de la Vallée Poussin means and convex conformal maps of the circle, *Pacific J. Math.* **8** (1958), 295–334.
- [25] C. Pommerenke, *Univalent functions*, Vandenhoeck & Ruprecht, Göttingen, 1975.
- [26] V. Ravichandran, Functions starlike with respect to n -ply symmetric, conjugate and symmetric conjugate points, *J. Indian Acad. Math.* **26** (2004), no. 1, 35–45.
- [27] V. Ravichandran, M. Hussain Khan, S. Silverman and K. G. Subramanian, Radius problems for a class of analytic functions, *Demonstratio Math.* **39** (2006), no. 1, 67–74.
- [28] V. Ravichandran, F. Rønning and T. N. Shanmugam, Radius of convexity and radius of starlikeness for some classes of analytic functions, *Complex Variables Theory Appl.* **33** (1997), no. 1-4, 265–280.
- [29] V. Ravichandran and T. N. Shanmugam, Radius problems for analytic functions, *Chinese J. Math.* **23** (1995), no. 4, 343–351.
- [30] F. Rønning, A survey on uniformly convex and uniformly starlike functions, *Ann. Univ. Mariae Curie-Skłodowska Sect. A* **47** (1993), 123–134.
- [31] F. Rønning, On starlike functions associated with parabolic regions, *Ann. Univ. Mariae Curie-Skłodowska Sect. A* **45** (1991), 117–122 (1992).
- [32] F. Rønning, Some radius results for univalent functions, *J. Math. Anal. Appl.* **194** (1995), no. 1, 319–327.
- [33] F. Rønning, Uniformly convex functions and a corresponding class of starlike functions, *Proc. Amer. Math. Soc.* **118** (1993), no. 1, 189–196.
- [34] T. Rosy, K. G. Subramanian and G. Murugusundaramoorthy, Neighbourhoods and partial sums of starlike functions based on Ruscheweyh derivatives, *JIPAM. J. Inequal. Pure Appl. Math.* **4** (2003), no. 4, Article 64, 8 pp. (electronic).
- [35] S. Ruscheweyh, *Convolutions in geometric function theory*, Presses Univ. Montréal, Montreal, Que., 1982.
- [36] St. Ruscheweyh and T. Sheil-Small, Hadamard products of Schlicht functions and the Pólya-Schoenberg conjecture, *Comment. Math. Helv.* **48** (1973), 119–135.
- [37] G. Ş. Sălăgean, Convolution properties of some classes of analytic functions with negative coefficients, in *Proc. Int. Symp on New Development of Geometric Function Theory and its Applications (Malaysia 2008)*, 12–16.
- [38] G. Ş. Sălăgean, Subclasses of univalent functions, in *Complex analysis—fifth Romanian-Finnish seminar, Part 1 (Bucharest, 1981)*, 362–372, *Lecture Notes in Math.*, 1013, Springer, Berlin.
- [39] S. M. Sarangi and B. A. Uralegaddi, The radius of convexity and starlikeness for certain classes of analytic functions with negative coefficients. I, *Atti Accad. Naz. Lincei Rend. Cl. Sci. Fis. Mat. Natur. (8)* **65** (1978), no. 1-2, 38–42 (1979).
- [40] S. Shamani, R. M. Ali, V. Ravichandran and S. K. Lee, Convolution and differential subordination for multivalent functions (2008), preprint.
- [41] T. N. Shanmugam, Convolution and differential subordination, *Internat. J. Math. Math. Sci.* **12** (1989), no. 2, 333–340.
- [42] T. N. Shanmugam and V. Ravichandran, Certain properties of uniformly convex functions, in *Computational methods and function theory 1994 (Penang)*, 319–324, World Sci. Publ., River Edge, NJ.
- [43] T. N. Shanmugam, S. Sivasubramanian and M. Darus, On a subclass of k -uniformly convex functions with negative coefficients, *Int. Math. Forum* **1** (2006), no. 33-36, 1677–1689.
- [44] H. Silverman, Univalent functions with negative coefficients, *Proc. Amer. Math. Soc.* **51** (1975), 109–116.
- [45] E. M. Silvia, A brief overview of subclasses of spiral-like functions, in *Current topics in analytic function theory*, 328–336, World Sci. Publ., River Edge, NJ.
- [46] E. M. Silvia, Subclasses of spiral-like functions, *Tamkang J. Math.* **14** (1983), no. 2, 161–169.
- [47] B. A. Uralegaddi, M. D. Ganigi and S. M. Sarangi, Univalent functions with positive coefficients, *Tamkang J. Math.* **25** (1994), no. 3, 225–230.