

## Submanifolds of a Lorentzian Para-Sasakian Manifold

<sup>1</sup>U.C. DE, <sup>2</sup>ADNAN AL-AQEEL AND <sup>3</sup>A.A. SHAIKH

<sup>1</sup>Department of Mathematics, University of Kalyani  
Kalyani- 741235, West Bengal, India

<sup>2</sup>Department of Mathematics and Computer Science, Kuwait University

<sup>3</sup>Department of Mathematics, North Bengal University,  
Darjeeling, West Bengal, India

<sup>1</sup>ucde@klyuniv.ernet.in, <sup>2</sup>math@sci.kuniv.edu.kw

**Abstract.** Recently, Matsumoto [1] introduced the notion of Lorentzian para-contact structure and studied its several properties. The object of the present paper is to study the submanifolds of Lorentzian para-Sasakian manifolds.

2000 Mathematics Subject Classification: 53C25, 53C40

Key words and phrases: Lorentzian Para-Sasakian manifold, totally umbilical submanifold, anti-invariant distribution.

### 1. Introduction

Let  $\bar{M}$  be a real  $n$ -dimensional manifold of class  $C^\infty$  endowed with an endomorphism  $\phi$  of the tangent bundle, a  $C^\infty$ -vector field  $\xi$  which is called the structure vector field, a 1-form  $\eta$  and a Lorentzian metric  $g$  with signature  $(-, +, +, +)$  satisfy

$$(1.1) \quad \phi^2 X = X + \eta(X)\xi, \quad \eta(\xi) = -1,$$

$$(1.2) \quad g(\phi X, \phi Y) = g(X, Y) + \eta(X)\eta(Y), \quad g(X, \xi) = \eta(X)$$

for any  $X, Y \in T\bar{M}$ , where  $T\bar{M}$  is the tangent bundle of  $\bar{M}$ . Then such a structure  $(\phi, \xi, \eta, g)$  is termed as Lorentzian para-contact structure and the manifold  $\bar{M}$  with a Lorentzian para-contact structure is called a Lorentzian para-contact manifold [1].

Also, in a Lorentzian para-contact structure the following relations hold:

$$(1.3) \quad \phi\xi = 0, \quad \eta \circ \phi = 0, \quad \text{rank}(\phi) = n - 1.$$

A Lorentzian para-contact manifold  $M$  is called a Lorentzian para-Sasakian (LP-Sasakian) manifold if

$$(1.4) \quad (\bar{\nabla}_X \phi)(Y) = g(X, Y)\xi + \eta(Y)X + 2\eta(X)\eta(Y)\xi,$$

and

$$(1.5) \quad \bar{\nabla}_X \xi = \phi X$$

for any  $X, Y \in T\bar{M}$ , where  $\bar{\nabla}$  is the Riemannian connection with respect to  $g$ . Again, if we put  $\Omega(X, Y) = g(X, \phi Y)$ , then  $\Omega$  is a symmetric (0,2) tensor field [1]. Thus we have from (1.5)

$$(1.6) \quad \Omega(X, Y) = (\bar{\nabla}_X \eta)Y.$$

Also, from (1.4), it follows that

$$(1.7) \quad (\bar{\nabla}_Z \Omega)(X, Y) = g(X, (\bar{\nabla}_Z \phi)Y) = g((\bar{\nabla}_Z \phi)X, Y),$$

$$(1.8) \quad (\bar{\nabla}_Z \Omega)(X, Y) = g(Y, Z)\eta(X) + g(X, Z)\eta(Y) + 2\eta(X)\eta(Y)\eta(Z)$$

for any  $X, Y, Z \in T\bar{M}$ .

LP-Sasakian manifolds have also been studied by K. Matsumoto and Mihai [2], Mihai and Rosca [3] and Matsumoto, De and Shaikh [4]. Let  $M$  be a Riemannian submanifold of a semi-Riemannian manifold  $\bar{M}$ . Then the Gauss and Weingarten formulae are given by

$$(1.9) \quad \bar{\nabla}_X Y = \nabla_X Y + h(X, Y),$$

$$(1.10) \quad \bar{\nabla}_X N = -A_N X + \nabla_X^\perp N$$

for each  $X, Y \in TM$  and each  $N \in T^\perp M$ , where  $\nabla$  is the Levi-Civita connection on  $M$ ,  $\nabla^\perp$  is the normal connection on the normal bundle  $T^\perp M$ ,  $h$  is the second fundamental form of  $M$  and  $A_N$  is the shape operator with respect to the normal section  $N$ . Then we know that

$$(1.11) \quad g(h(X, Y), N) = g(A_N X, Y)$$

for each  $X, Y \in TM$  and  $N \in T^\perp M$ . We denote by the same symbol  $g$  both metrics on  $\bar{M}$  and  $M$ .

**Definition 1.1.** A submanifold  $M$  is said to be

(i) totally geodesic in  $\bar{M}$  if

$$(1.12) \quad h = 0 \quad \text{or equivalently} \quad A_N = 0$$

for any  $N \in T^\perp M$ .

(ii) minimal in  $\bar{M}$  if the mean curvature vector  $H$  satisfies

$$(1.13) \quad H \stackrel{\text{def}}{=} \frac{Tr(h)}{\dim M} = 0$$

and (iii) totally umbilical if

$$(1.14) \quad h(X, Y) = g(X, Y)H.$$

In Section 2 of this paper, we obtain some properties of submanifolds of an LP-Sasakian manifold. In the last section, we prove the main result concerning the non-existence of an anti-invariant distribution on the submanifold of an LP-Sasakian manifold which include the results of [5] as particular case. Sub-manifolds of an LP-Sasakian manifold have been studied by two of the present authors [5], Prasad [6], Kalpana and Guha [7] and others. Throughout this paper, we shall assume the following notation:

- (a)  $M$  is a submanifold of an LP-Sasakian manifold  $\bar{M}$ ,
- (b)  $\{\xi\}$  is the 1- dimensional distribution spanned by  $\xi$ ,
- (c)  $TM$  and  $T^\perp M$  are the tangent and normal bundles of  $M$ , respectively, and
- (d)  $D^\perp$  is an anti- invariant distribution (i.e.  $\phi D^\perp \subset T^\perp M$ ) of  $M$  such that  $D^\perp \cap \{\xi\} = 0$ .

**2. Submanifolds of an LP-Sasakian manifold**

We first prove the following lemma:

**Lemma 2.1.** *For a submanifold  $M$  of an LP-Sasakian manifold  $\bar{M}$ , we have*

$$(2.1) \quad \phi X = \nabla_X \xi + h(X, \xi), \quad \xi \in TM,$$

$$(2.2) \quad \phi X = -A_\xi X + \nabla_X^\perp \xi, \quad \xi \in T^\perp M,$$

$$(2.3) \quad \eta(A_N X) = 0, \quad \xi \in T^\perp M,$$

$$(2.4) \quad \eta(A_N X) = g(\phi X, N), \quad \xi \in TM$$

for  $X \in TM$  and  $N \in T^\perp M$ .

*Proof.* From (1.5) and (1.9), we get (2.1). Also, from (1.5) and (1.10), we obtain (2.2). Again, in view of (1.2), (2.3) is obvious. Lastly, for  $\xi \in TM$ , we get

$$\eta(A_N X) = g(\xi, A_N X) = -g(\xi, \bar{\nabla}_X N) = g(\bar{\nabla}_X \xi, N) = g(\phi X, N),$$

where (1.2), (1.5) and (1.10) have been used. These complete the proof of our lemma. □

**Theorem 2.1.** *Let  $M$  be a submanifold of an LP-Sasakian manifold  $\bar{M}$  such that the structure vector field  $\xi$  is tangent to  $M$ . Then  $M$  is invariant (i.e.  $\phi TM \subset TM$ ) if and only if  $h(X, \xi) = 0$ , and  $M$  is anti-invariant (i.e.  $\phi TM \subset T^\perp M$ ) if and only if  $\nabla_X \xi = 0$ .*

Since it is trivial from (2.1), we omit to prove our theorem.

**Theorem 2.2.** *If  $M$  is a totally umbilical submanifold of an LP-Sasakian manifold  $\bar{M}$  such that the structure vector field  $\xi$  is tangent to  $M$ , then*

- (i)  $M$  is necessarily minimal and consequently totally geodesic and
- (ii)  $M$  is an invariant submanifold of  $\bar{M}$  and  $\nabla_X \xi \neq 0$ .

*Proof.* Let  $M$  be totally umbilical. Using (1.2), (1.3) and (2.1) in (1.14), we get

$$0 = h(\xi, \xi) = g(\xi, \xi)H = -H.$$

Hence, in view of (1.13) and (1.14), we obtain (i).

The second part follows from Theorem 2.1 and the above (i). □

**Theorem 2.3.** *A submanifold  $M$  of an LP-Sasakian manifold  $\bar{M}$  such that the structure vector field  $\xi$  is normal to  $M$  is anti-invariant in  $\bar{M}$  if and only if  $A_\xi X = 0$ . Consequently, if  $M$  is totally geodesic, then it is anti-invariant.*

*Proof.* Since  $\xi$  is normal to  $M$ , by virtue of (1.10) and (2.2), we get

$$g(\phi X, Y) = -g(A_\xi X, Y) = -g(h(X, Y), \xi), \quad X, Y \in TM,$$

which provides the proof of our theorem.  $\square$

### 3. Non-existence of an anti-invariant distribution

First, we prove the following:

**Lemma 3.1.** *For a submanifold  $M$  of an LP-Sasakian manifold  $\bar{M}$ , we have*

$$(3.1) \quad (\bar{\nabla}_Z \Omega)(X, Y) = -g(A_{\phi Y} X, Z) - \Omega(X, \nabla_Z Y) - \Omega(X, h(Z, Y))$$

for  $Y \in D^\perp$ ,  $X, Z \in TM$ .

$$(3.2) \quad (\bar{\nabla}_Z \Omega)(X, Y) = -g(A_{\phi X} Y + A_{\phi Y} X, Z)$$

for  $X, Y \in D^\perp$ ,  $Z \in TM$ .

*Proof.* Let  $Y \in D^\perp$ ,  $Z \in TM$ . Then, by virtue of (1.11) and the fact  $\phi Y \in T^\perp M$ , we get

$$(3.3) \quad (\bar{\nabla}_Z \phi)Y = -A_{\phi Y} Z + \nabla_Z^\perp \phi Y - \phi(\bar{\nabla}_Z Y).$$

Using this equation in (1.7), we can easily derive (3.1).

Next, in the special case of  $X \in D^\perp$ , since  $\phi X \in T^\perp M$ , (3.1) in view of (1.5) and (1.11) yields (3.2).  $\square$

**Lemma 3.2.** *Let  $M$  be a submanifold of an LP-Sasakian manifold  $\bar{M}$  and  $D^\perp \perp \{\xi\}$ . Then we get*

$$(3.4) \quad (\bar{\nabla}_Z \Omega)(X, X) = 0$$

for  $X \in D^\perp$  and  $Z \in TM$ . And consequently

$$(3.5) \quad A_{\phi X} X = 0$$

for  $X \in D^\perp$ .

*Proof.* Since  $D^\perp \perp \{\xi\}$ , we have  $\eta(X) = 0$  for any  $X \in D^\perp$  and hence in view of (1.8), we get (3.4). Again (3.5) follows from (3.2) and (3.4).  $\square$

**Theorem 3.1.** *There does not exist any anti-invariant distribution  $D^\perp$  on a submanifold  $M$  of an LP-Sasakian manifold  $\bar{M}$  if  $\xi$  is tangent to  $M$  and  $D^\perp \perp \{\xi\}$ .*

*Proof.* Since  $D^\perp \perp \{\xi\}$ , we get  $\eta(X) = 0$  for any  $X \in D^\perp$ . Thus, from (1.2), (2.4) and (3.5), we have

$$0 = \eta(A_{\phi X} X) = g(\phi X, \phi X) = g(X, X)$$

for any  $X \in D^\perp$ . This means  $D^\perp = \{0\}$ .

This proves the theorem.  $\square$

A submanifold  $M$  of an LP-Sasakian manifold  $\bar{M}$  is said to be semi-invariant submanifold [8] if the following conditions are satisfied

- (i)  $TM = D \oplus D^\perp \oplus \{\xi\}$ , where  $D$ ,  $D^\perp$  are orthogonal differentiable distributions on  $M$  and  $\{\xi\}$  is the 1-dimensional distribution spanned by  $\xi$ ,
- (ii) The distribution  $D$  is invariant by  $\phi$ , that is,  $\phi D_x = D_x$  for each  $x \in M$ ,
- (iii) The distribution  $D^\perp$  is anti-invariant under  $\phi$ , that is  $\phi D^\perp \subset T_x M^\perp$  for each  $x \in M$ .

If both the distribution  $D$  and  $D^\perp$  are non-zero then the semi-invariant submanifold is called a proper semi-invariant submanifold.

Hence by virtue of Theorem 3.1, we have the following:

**Corollary 3.1.** *An LP-Sasakian manifold does not admit any proper semi-invariant submanifold.*

**Acknowledgment.** The authors are thankful to the referee for his valuable suggestions in the improvement of the paper.

## References

- [1] K. Matsumoto, On Lorentzian para contact manifolds, *Bull. Yamagata Univ. Nat. Sci.* **12** (1989), 151–156.
- [2] K. Matsumoto and I. Mihai, On a certain transformation in a Lorentzian para-Sasakian manifold, *Tensor, N. S.* **47** (1988), 189–197.
- [3] I. Mihai and R. Rosca, On Lorentzian P-Sasakian manifolds, *Classical Analysis, World Scientific Publ., Singapore* (1992), 155–169.
- [4] K. Matsumoto, U. C. De and A. A. Shaikh, On Lorentzian Para-Sasakian manifolds, *Rendiconti del seminario Matematico di Messina Serie II supplemento al n. 3* (1999).
- [5] U. C. De, A. A. Shaikh, Non-existence of proper semi-invariant submanifolds of a Lorentzian para-Sasakian manifold, *Bull. Malays. Math. Soc.(2)* **22** (1999), 179–183.
- [6] B. Prasad, Semi-invariant submanifolds of a Lorentzian Para-Sasakian manifold, *Bull. Malays. Math. Soc. (2)* **21** (1998), 21–26.
- [7] Kalpana and G. Guha, Semi-invariant submanifolds of a Lorentzian Para-Sasakian manifold, *Ganit, J. Bangladesh Math. Soc.* **13** (1993), 71–76.
- [8] A. Bejancu and N. Papaghiuc, Semi-invariant submanifold of a Sasakian manifold, *An. Stiint. Univ. "Al. I. Cuza" Iasi* **27** (1981), 163–170.