

Research Article

Initial Coefficients of Biunivalent Functions

See Keong Lee,¹ V. Ravichandran,² and Shamani Supramaniam¹

¹ School of Mathematical Sciences, Universiti Sains Malaysia (USM), 11800 Penang, Malaysia

² Department of Mathematics, University of Delhi, Delhi 110 007, India

Correspondence should be addressed to See Keong Lee; sklee@cs.usm.my

Received 24 January 2014; Accepted 12 March 2014; Published 8 April 2014

Academic Editor: Om P. Ahuja

Copyright © 2014 See Keong Lee et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

An analytic function f defined on the open unit disk is biunivalent if the function f and its inverse f^{-1} are univalent in \mathbb{D} . Estimates for the initial coefficients of biunivalent functions f are investigated when f and f^{-1} , respectively, belong to some subclasses of univalent functions. Some earlier results are shown to be special cases of our results.

1. Introduction

Let \mathcal{S} be the class of all univalent analytic functions f in the open unit disk $\mathbb{D} := \{z \in \mathbb{C} : |z| < 1\}$ and normalized by the conditions $f(0) = 0$ and $f'(0) = 1$. For $f \in \mathcal{S}$, it is well known that the n th coefficient is bounded by n . The bounds for the coefficients give information about the geometric properties of these functions. Indeed, the bound for the second coefficient of functions in the class \mathcal{S} gives rise to the growth, distortion and covering theorems for univalent functions. In view of the influence of the second coefficient in the geometric properties of univalent functions, it is important to know the bounds for the (initial) coefficients of functions belonging to various subclasses of univalent functions. In this paper, we investigate this coefficient problem for certain subclasses of biunivalent functions.

Recall that the Koebe one-quarter theorem [1] ensures that the image of \mathbb{D} under every univalent function $f \in \mathcal{S}$ contains a disk of radius $1/4$. Thus, every univalent function f has an inverse f^{-1} satisfying $f^{-1}(f(z)) = z$, ($z \in \mathbb{D}$), and

$$f(f^{-1}(w)) = w, \quad \left(|w| < r_0(f), r_0(f) \geq \frac{1}{4}\right). \quad (1)$$

A function $f \in \mathcal{S}$ is *biunivalent* in \mathbb{D} if both f and f^{-1} are univalent in \mathbb{D} . Let σ denote the class of biunivalent functions defined in the unit disk \mathbb{D} . Lewin [2] investigated this class σ and obtained the bound for the second coefficient of the biunivalent functions. Several authors subsequently studied similar problems in this direction (see [3, 4]). A function

$f \in \sigma$ is *bistarlike* or *strongly bistarlike* or *biconvex* of order α if f and f^{-1} are both starlike, strongly starlike, or convex of order α , respectively. Brannan and Taha [5] obtained estimates for the initial coefficients of bistarlike, strongly bistarlike, and biconvex functions. Bounds for the initial coefficients of several classes of functions were also investigated in [6–24].

An analytic function f is *subordinate* to an analytic function g , written $f(z) \prec g(z)$, if there is an analytic function $w : \mathbb{D} \rightarrow \mathbb{D}$ with $w(0) = 0$ satisfying $f(z) = g(w(z))$. Ma and Minda [25] unified various subclasses of starlike (\mathcal{S}^*) and convex functions (\mathcal{C}) by requiring that either the quantity $zf'(z)/f(z)$ or $1 + zf''(z)/f'(z)$ is subordinate to a more general superordinate function φ with positive real part in the unit disk \mathbb{D} , $\varphi(0) = 1$, $\varphi'(0) > 0$, φ maps \mathbb{D} onto a region starlike with respect to 1 and symmetric with respect to the real axis. The class $\mathcal{S}^*(\varphi)$ of Ma-Minda starlike functions with respect to φ consists of functions $f \in \mathcal{S}$ satisfying the subordination $zf'(z)/f(z) \prec \varphi(z)$. Similarly, the class $\mathcal{C}(\varphi)$ of Ma-Minda convex functions consists of functions $f \in \mathcal{S}$ satisfying the subordination $1 + zf''(z)/f'(z) \prec \varphi(z)$. Ma and Minda investigated growth and distortion properties of functions in $\mathcal{S}^*(\varphi)$ and $\mathcal{C}(\varphi)$ as well as Fekete-Szegő inequalities for $\mathcal{S}^*(\varphi)$ and $\mathcal{C}(\varphi)$. Their proof of Fekete-Szegő inequalities requires the univalence of φ . Ali et al. [7] investigated Fekete-Szegő problems for various other classes and their proof does not require the univalence or starlikeness of φ . In particular, their results are valid even if one just assumes the function φ to have a series expansion

of the form $\varphi(z) = 1 + B_1z + B_2z^2 + \dots$, $B_1 > 0$. So, in this paper, we assume that φ has series expansion $\varphi(z) = 1 + B_1z + B_2z^2 + \dots$, B_1, B_2 are real, and $B_1 > 0$. A function f is Ma-Minda bistarlike or Ma-Minda biconvex if both f and f^{-1} are, respectively, Ma-Minda starlike or convex. Motivated by the Fekete-Szegő problem for the classes of Ma-Minda starlike and Ma-Minda convex functions [25], Ali et al. [26] recently obtained estimates of the initial coefficients for biunivalent Ma-Minda starlike and Ma-Minda convex functions.

The present work is motivated by the results of Kędzierawski [27] who considered functions f belonging to certain subclasses of univalent functions while their inverses f^{-1} belong to some other subclasses of univalent functions. Among other results, he obtained the following coefficient estimates.

Theorem 1 (see [27]). *Let $f \in \sigma$ with Taylor series $f(z) = z + a_2z^2 + \dots$ and $g = f^{-1}$. Then,*

$$|a_2| \leq \begin{cases} 1.5894 & \text{if } f \in \mathcal{S}, g \in \mathcal{S}, \\ \sqrt{2} & \text{if } f \in \mathcal{S}^*, g \in \mathcal{S}^*, \\ 1.507 & \text{if } f \in \mathcal{S}^*, g \in \mathcal{S}, \\ 1.224 & \text{if } f \in \mathcal{C}, g \in \mathcal{S}. \end{cases} \quad (2)$$

We need the following classes investigated in [6, 7, 26].

Definition 2. Let $\varphi : \mathbb{D} \rightarrow \mathbb{C}$ be analytic and $\varphi(z) = 1 + B_1z + B_2z^2 + \dots$ with $B_1 > 0$ and $B_2 \in \mathbb{R}$. For $\alpha \geq 0$, let

$$\mathcal{M}(\alpha, \varphi) := \left\{ f \in \mathcal{S} : (1 - \alpha) \frac{zf'(z)}{f(z)} + \alpha \left(1 + \frac{zf''(z)}{f'(z)} \right) < \varphi(z) \right\},$$

$\mathcal{L}(\alpha, \varphi)$

$$:= \left\{ f \in \mathcal{S} : \left(\frac{zf'(z)}{f(z)} \right)^\alpha \left(1 + \frac{zf''(z)}{f'(z)} \right)^{1-\alpha} < \varphi(z) \right\},$$

$$\mathcal{P}(\alpha, \varphi) := \left\{ f \in \mathcal{S} : \frac{zf'(z)}{f(z)} + \alpha \frac{z^2f''(z)}{f(z)} < \varphi(z) \right\}. \quad (3)$$

In this paper, we obtain the estimates for the second and third coefficients of functions f when

- (i) $f \in \mathcal{P}(\alpha, \varphi)$ and $g := f^{-1} \in \mathcal{P}(\beta, \psi)$, or $g \in \mathcal{M}(\beta, \psi)$, or $g \in \mathcal{L}(\beta, \psi)$,
- (ii) $f \in \mathcal{M}(\alpha, \varphi)$ and $g \in \mathcal{M}(\beta, \psi)$, or $g \in \mathcal{L}(\beta, \psi)$,
- (iii) $f \in \mathcal{L}(\alpha, \varphi)$ and $g \in \mathcal{L}(\beta, \psi)$.

2. Coefficient Estimates

In the sequel, it is assumed that φ and ψ are analytic functions of the form

$$\begin{aligned} \varphi(z) &= 1 + B_1z + B_2z^2 + B_3z^3 + \dots \quad (B_1 > 0), \\ \psi(z) &= 1 + D_1z + D_2z^2 + D_3z^3 + \dots \quad (D_1 > 0). \end{aligned} \quad (4)$$

Theorem 3. *Let $f \in \sigma$ and $g = f^{-1}$. If $f \in \mathcal{P}(\alpha, \varphi)$, $g \in \mathcal{P}(\beta, \psi)$ and f is of the form*

$$f(z) = z + \sum_{n=2}^{\infty} a_n z^n, \quad (5)$$

then

$$\begin{aligned} |a_2| &\leq \left(B_1 D_1 \sqrt{B_1(1+3\beta) + D_1(1+3\alpha)} \right) \\ &\quad \times \left(\left| \sigma B_1^2 D_1^2 - (1+2\alpha)^2(1+3\beta)(B_2 - B_1) D_1^2 \right. \right. \\ &\quad \left. \left. - (1+2\beta)^2(1+3\alpha)(D_2 - D_1) B_1^2 \right| \right)^{-1/2}, \\ 2\sigma |a_3| &\leq B_1(3+10\beta) + D_1(1+2\alpha) + (3+10\beta) |B_2 - B_1| \\ &\quad + \frac{(1+2\beta)^2 B_1^2 |D_2 - D_1|}{D_1^2(1+2\alpha)}, \end{aligned} \quad (6)$$

(7)

where $\sigma := 2 + 7\alpha + 7\beta + 24\alpha\beta$.

Proof. Since $f \in \mathcal{P}(\alpha, \varphi)$ and $g \in \mathcal{P}(\beta, \psi)$, $g = f^{-1}$, then there exist analytic functions $u, v : \mathbb{D} \rightarrow \mathbb{D}$, with $u(0) = v(0) = 0$, satisfying

$$\begin{aligned} \frac{zf'(z)}{f(z)} + \frac{\alpha z^2 f''(z)}{f(z)} &= \varphi(u(z)), \\ \frac{wg'(w)}{g(w)} + \frac{\beta w^2 g''(w)}{g(w)} &= \psi(v(w)). \end{aligned} \quad (8)$$

Define the functions p_1 and p_2 by

$$\begin{aligned} p_1(z) &:= \frac{1+u(z)}{1-u(z)} = 1 + c_1z + c_2z^2 + \dots, \\ p_2(z) &:= \frac{1+v(z)}{1-v(z)} = 1 + b_1z + b_2z^2 + \dots, \end{aligned} \quad (9)$$

or, equivalently,

$$\begin{aligned} u(z) &= \frac{p_1(z) - 1}{p_1(z) + 1} = \frac{1}{2} \left(c_1z + \left(c_2 - \frac{c_1^2}{2} \right) z^2 + \dots \right), \\ v(z) &= \frac{p_2(z) - 1}{p_2(z) + 1} = \frac{1}{2} \left(b_1z + \left(b_2 - \frac{b_1^2}{2} \right) z^2 + \dots \right). \end{aligned} \quad (10)$$

Then, p_1 and p_2 are analytic in \mathbb{D} with $p_1(0) = 1 = p_2(0)$. Since $u, v : \mathbb{D} \rightarrow \mathbb{D}$, the functions p_1 and p_2 have positive real part in \mathbb{D} , and $|b_i| \leq 2$ and $|c_i| \leq 2$. In view of (8) and (10), it is clear that

$$\begin{aligned} \frac{zf'(z)}{f(z)} + \frac{\alpha z^2 f''(z)}{f(z)} &= \varphi \left(\frac{p_1(z) - 1}{p_1(z) + 1} \right), \\ \frac{wg'(w)}{g(w)} + \frac{\beta w^2 g''(w)}{g(w)} &= \psi \left(\frac{p_2(w) - 1}{p_2(w) + 1} \right). \end{aligned} \quad (11)$$

Using (10) together with (4), it is evident that

$$\begin{aligned} \varphi\left(\frac{p_1(z)-1}{p_1(z)+1}\right) &= 1 + \frac{1}{2}B_1c_1z \\ &\quad + \left(\frac{1}{2}B_1\left(c_2 - \frac{c_1^2}{2}\right) + \frac{1}{4}B_2c_1^2\right)z^2 + \dots, \\ \psi\left(\frac{p_2(w)-1}{p_2(w)+1}\right) &= 1 + \frac{1}{2}D_1b_1w \\ &\quad + \left(\frac{1}{2}D_1\left(b_2 - \frac{b_1^2}{2}\right) + \frac{1}{4}D_2b_1^2\right)w^2 + \dots. \end{aligned} \tag{12}$$

Since f has the Maclaurin series given by (5), a computation shows that its inverse $g = f^{-1}$ has the expansion

$$g(w) = f^{-1}(w) = w - a_2w^2 + (2a_2^2 - a_3)w^3 + \dots \tag{13}$$

Since

$$\begin{aligned} \frac{zf'(z)}{f(z)} + \frac{\alpha z^2 f''(z)}{f(z)} &= 1 + a_2(1 + 2\alpha)z \\ &\quad + (2(1 + 3\alpha)a_3 - (1 + 2\alpha)a_2^2)z^2 + \dots, \\ \frac{wg'(w)}{g(w)} + \frac{\beta w^2 g''(w)}{g(w)} &= 1 - (1 + 2\beta)a_2w \\ &\quad + ((3 + 10\beta)a_2^2 - 2(1 + 3\beta)a_3)w^2 + \dots, \end{aligned} \tag{14}$$

it follows from (11) and (12) that

$$a_2(1 + 2\alpha) = \frac{1}{2}B_1c_1, \tag{15}$$

$$2(1 + 3\alpha)a_3 - (1 + 2\alpha)a_2^2 = \frac{1}{2}B_1\left(c_2 - \frac{c_1^2}{2}\right) + \frac{1}{4}B_2c_1^2, \tag{16}$$

$$-(1 + 2\beta)a_2 = \frac{1}{2}D_1b_1, \tag{17}$$

$$(3 + 10\beta)a_2^2 - 2(1 + 3\beta)a_3 = \frac{1}{2}D_1\left(b_2 - \frac{b_1^2}{2}\right) + \frac{1}{4}D_2b_1^2. \tag{18}$$

It follows from (15) and (17) that

$$b_1 = -\frac{B_1(1 + 2\beta)}{D_1(1 + 2\alpha)}c_1. \tag{19}$$

Equations (15), (16), (18), and (19) lead to

$$\begin{aligned} a_2^2 &= B_1^2 D_1^2 [B_1(1 + 3\beta)c_2 + D_1(1 + 3\alpha)b_2] \\ &\quad \times \left(2\left[\sigma B_1^2 D_1^2 - (1 + 2\alpha)^2(1 + 3\beta)(B_2 - B_1)D_1^2\right. \right. \\ &\quad \left. \left. - (1 + 2\beta)^2(1 + 3\alpha)(D_2 - D_1)B_1^2\right]\right)^{-1}, \end{aligned} \tag{20}$$

where $\sigma := 2 + 7\alpha + 7\beta + 24\alpha\beta$, which, in view of $|b_2| \leq 2$ and $|c_2| \leq 2$, gives us the desired estimate on $|a_2|$ as asserted in (6).

By using (16), (18), and (19), we get

$$\begin{aligned} 2\sigma a_3 &= \frac{1}{2}[B_1(3 + 10\beta)c_2 + D_1(1 + 2\alpha)b_2] \\ &\quad + \frac{c_1^2}{4}\left[(3 + 10\beta)(B_2 - B_1) + \frac{(1 + 2\beta)^2 B_1^2 (D_2 - D_1)}{D_1^2(1 + 2\alpha)}\right], \end{aligned} \tag{21}$$

and this yields the estimate given in (7). \square

Remark 4. When $\alpha = \beta = 0$ and $B_1 = B_2 = 2, D_1 = D_2 = 2$, then (6) reduces to Theorem 1. When $\beta = \alpha$ and $\psi = \varphi$, Theorem 3 reduces to [26, Theorem 2.2].

Theorem 5. Let $f \in \sigma$ and $g = f^{-1}$. If $f \in \mathcal{P}(\alpha, \varphi)$ and $g \in \mathcal{M}(\beta, \psi)$, then

$$\begin{aligned} |a_2| &\leq \left(B_1 D_1 \sqrt{B_1(1 + 2\beta) + D_1(1 + 3\alpha)}\right) \\ &\quad \times \left(\left|\sigma B_1^2 D_1^2 - (1 + 2\alpha)^2(1 + 2\beta)(B_2 - B_1)D_1^2\right. \right. \\ &\quad \left. \left. - (1 + \beta)^2(1 + 3\alpha)(D_2 - D_1)B_1^2\right|\right)^{-1/2}, \end{aligned} \tag{22}$$

$$\begin{aligned} 2\sigma |a_3| &\leq B_1(3 + 5\beta) + D_1(1 + 2\alpha) + (3 + 5\beta)|B_2 - B_1| \\ &\quad + \frac{(1 + \beta)^2 B_1^2 |D_2 - D_1|}{D_1^2(1 + 2\alpha)}, \end{aligned} \tag{23}$$

where $\sigma := 2 + 7\alpha + 3\beta + 11\alpha\beta$.

Proof. Let $f \in \mathcal{P}(\alpha, \varphi)$ and $g \in \mathcal{M}(\beta, \psi), g = f^{-1}$. Then, there exist analytic functions $u, v : \mathbb{D} \rightarrow \mathbb{D}$, with $u(0) = v(0) = 0$, such that

$$\begin{aligned} \frac{zf'(z)}{f(z)} + \frac{\alpha z^2 f''(z)}{f(z)} &= \varphi(u(z)), \\ (1 - \beta)\frac{wg'(w)}{g(w)} + \beta\left(1 + \frac{wg''(w)}{g'(w)}\right) &= \psi(v(w)). \end{aligned} \tag{24}$$

Since

$$\begin{aligned} & \frac{zf'(z)}{f(z)} + \frac{\alpha z^2 f''(z)}{f(z)} \\ &= 1 + a_2(1 + 2\alpha)z \\ & \quad + (2(1 + 3\alpha)a_3 - (1 + 2\alpha)a_2^2)z^2 + \dots, \end{aligned} \tag{25}$$

$$\begin{aligned} & (1 - \beta) \frac{wg'(w)}{g(w)} + \beta \left(1 + \frac{wg''(w)}{g'(w)} \right) \\ &= 1 - (1 + \beta)a_2w \\ & \quad + ((3 + 5\beta)a_2^2 - 2(1 + 2\beta)a_3)w^2 + \dots, \end{aligned}$$

(12) and (24) yield

$$a_2(1 + 2\alpha) = \frac{1}{2}B_1c_1, \tag{26}$$

$$2(1 + 3\alpha)a_3 - (1 + 2\alpha)a_2^2 = \frac{1}{2}B_1 \left(c_2 - \frac{c_1^2}{2} \right) + \frac{1}{4}B_2c_1^2, \tag{27}$$

$$-(1 + \beta)a_2 = \frac{1}{2}D_1b_1, \tag{28}$$

$$(3 + 5\beta)a_2^2 - 2(1 + 2\beta)a_3 = \frac{1}{2}D_1 \left(b_2 - \frac{b_1^2}{2} \right) + \frac{1}{4}D_2b_1^2. \tag{29}$$

It follows from (26) and (28) that

$$b_1 = -\frac{B_1(1 + \beta)}{D_1(1 + 2\alpha)}c_1. \tag{30}$$

Hence, (26), (27), (29), and (30) lead to

$$\begin{aligned} a_2^2 &= (B_1^2D_1^2 [B_1(1 + 2\beta)c_2 + D_1(1 + 3\alpha)b_2]) \\ & \quad \times (2 [\sigma B_1^2D_1^2 - (1 + 2\alpha)^2(1 + 2\beta)(B_2 - B_1)D_1^2 \\ & \quad - (1 + 2\beta)^2(1 + 3\alpha)(D_2 - D_1)B_1^2])^{-1}, \end{aligned} \tag{31}$$

which gives us the desired estimate on $|a_2|$ as asserted in (22) when $|b_2| \leq 2$ and $|c_2| \leq 2$.

Further, (27), (29), and (30) give

$$\begin{aligned} 2\sigma a_3 &= \frac{1}{2} [B_1(3 + 5\beta)c_2 + D_1(1 + 2\alpha)b_2] \\ & \quad + \frac{c_1^2}{4} \left[(3 + 5\beta)(B_2 - B_1) + \frac{(1 + \beta)^2B_1^2(D_2 - D_1)}{D_1^2(1 + 2\alpha)} \right], \end{aligned} \tag{32}$$

and this yields the estimate given in (23). \square

Theorem 6. Let $f \in \sigma$ and $g = f^{-1}$. If $f \in \mathcal{P}(\alpha, \varphi)$ and $g \in \mathcal{L}(\beta, \psi)$, then

$$\begin{aligned} |a_2| &\leq (B_1D_1 \sqrt{2 [B_1(3 - 2\beta) + D_1(1 + 3\alpha)]}) \\ & \quad \times (|\sigma B_1^2D_1^2 - 2(1 + 2\alpha)^2(3 - 2\beta)(B_2 - B_1)D_1^2 \\ & \quad - 2(2 - \beta)^2(1 + 3\alpha)(D_2 - D_1)B_1^2|)^{-1/2}, \\ |\sigma a_3| &\leq \frac{1}{2}B_1(\beta^2 - 11\beta + 16) + D_1(1 + 2\alpha) \\ & \quad + \frac{1}{2}(\beta^2 - 11\beta + 16)|B_2 - B_1| \\ & \quad + \frac{(2 - \beta)^2B_1^2|D_2 - D_1|}{D_1^2(1 + 2\alpha)}, \end{aligned} \tag{33}$$

where $\sigma := 10 + 36\alpha - 7\beta - 25\alpha\beta + \beta^2 + 3\alpha\beta^2$.

Proof. Let $f \in \mathcal{P}(\alpha, \varphi)$ and $g \in \mathcal{L}(\beta, \psi)$, $g = f^{-1}$. Then, there are analytic functions $u, v : \mathbb{D} \rightarrow \mathbb{D}$, with $u(0) = v(0) = 0$, satisfying

$$\begin{aligned} & \frac{zf'(z)}{f(z)} + \frac{\alpha z^2 f''(z)}{f(z)} = \varphi(u(z)), \\ & \left(\frac{wg'(w)}{g(w)} \right)^\beta \left(1 + \frac{wg''(w)}{g'(w)} \right)^{1-\beta} = \psi(v(w)). \end{aligned} \tag{34}$$

Using

$$\begin{aligned} & \frac{zf'(z)}{f(z)} + \frac{\alpha z^2 f''(z)}{f(z)} \\ &= 1 + a_2(1 + 2\alpha)z \\ & \quad + (2(1 + 3\alpha)a_3 - (1 + 2\alpha)a_2^2)z^2 + \dots, \\ & \left(\frac{wg'(w)}{g(w)} \right)^\beta \left(1 + \frac{wg''(w)}{g'(w)} \right)^{1-\beta} \\ &= 1 - (2 - \beta)a_2w \\ & \quad + \left(\left(8(1 - \beta) + \frac{1}{2}\beta(\beta + 5) \right) a_2^2 - 2(3 - 2\beta)a_3 \right) w^2 \\ & \quad + \dots, \end{aligned} \tag{35}$$

and (12) and (34) will yield

$$\begin{aligned} a_2(1 + 2\alpha) &= \frac{1}{2}B_1c_1, \\ 2(1 + 3\alpha)a_3 - (1 + 2\alpha)a_2^2 &= \frac{1}{2}B_1 \left(c_2 - \frac{c_1^2}{2} \right) + \frac{1}{4}B_2c_1^2, \\ -(2 - \beta)a_2 &= \frac{1}{2}D_1b_1, \end{aligned}$$

$$\begin{aligned} & \left[8(1-\beta) + \frac{\beta}{2}(\beta+5) \right] a_2^2 - 2(3-2\beta)a_3 \\ &= \frac{1}{2}D_1 \left(b_2 - \frac{b_1^2}{2} \right) + \frac{1}{4}D_2 b_1^2. \end{aligned} \tag{36}$$

Further implication of (36) and applying the fact that $|b_2| \leq 2$ and $|c_2| \leq 2$ give the estimates in (33). \square

Theorem 7. Let $f \in \sigma$ and $g = f^{-1}$. If $f \in \mathcal{M}(\alpha, \varphi)$, $g \in \mathcal{M}(\beta, \psi)$, then

$$\begin{aligned} |a_2| &\leq \left(B_1 D_1 \sqrt{B_1(1+2\beta) + D_1(1+2\alpha)} \right) \\ &\quad \times \left(\left| \sigma B_1^2 D_1^2 - (1+\alpha)^2(1+2\beta)(B_2 - B_1) D_1^2 \right. \right. \\ &\quad \left. \left. - (1+\beta)^2(1+2\alpha)(D_2 - D_1) B_1^2 \right| \right)^{-1/2}, \\ 2\sigma |a_3| &\leq B_1(3+5\beta) + D_1(1+3\alpha) + (3+5\beta)|B_2 - B_1| \\ &\quad + \frac{(1+\beta)^2(1+3\alpha)B_1^2|D_2 - D_1|}{D_1^2(1+\alpha)^2}, \end{aligned} \tag{37}$$

where $\sigma := 2 + 3\alpha + 3\beta + 4\alpha\beta$.

Proof. For $f \in \mathcal{M}(\alpha, \varphi)$ and $g \in \mathcal{M}(\beta, \psi)$, $g = f^{-1}$, there exist analytic functions $u, v : \mathbb{D} \rightarrow \mathbb{D}$, with $u(0) = v(0) = 0$, satisfying

$$\begin{aligned} (1-\alpha) \frac{zf'(z)}{f(z)} + \alpha \left(1 + \frac{zf''(z)}{f'(z)} \right) &= \varphi(u(z)), \\ (1-\beta) \frac{wg'(w)}{g(w)} + \beta \left(1 + \frac{wg''(w)}{g'(w)} \right) &= \psi(v(w)). \end{aligned} \tag{38}$$

Since

$$\begin{aligned} (1-\alpha) \frac{zf'(z)}{f(z)} + \alpha \left(1 + \frac{zf''(z)}{f'(z)} \right) &= 1 + (1+\alpha)a_2z \\ &\quad + (2(1+2\alpha)a_3 - (1+3\alpha)a_2^2)z^2 + \dots, \\ (1-\beta) \frac{wg'(w)}{g(w)} + \beta \left(1 + \frac{wg''(w)}{g'(w)} \right) &= 1 - (1+\beta)a_2w \\ &\quad + ((3+5\beta)a_2^2 - 2(1+2\beta)a_3)w^2 + \dots, \end{aligned} \tag{39}$$

then (12) and (38) yield

$$\begin{aligned} a_2(1+\alpha) &= \frac{1}{2}B_1c_1, \\ 2(1+2\alpha)a_3 - (1+3\alpha)a_2^2 &= \frac{1}{2}B_1 \left(c_2 - \frac{c_1^2}{2} \right) + \frac{1}{4}B_2c_1^2, \\ -(1+\beta)a_2 &= \frac{1}{2}D_1b_1, \\ (3+5\beta)a_2^2 - 2(1+2\beta)a_3 &= \frac{1}{2}D_1 \left(b_2 - \frac{b_1^2}{2} \right) + \frac{1}{4}D_2b_1^2. \end{aligned} \tag{40}$$

Further implication of (40) and applying the fact that $|b_2| \leq 2$ and $|c_2| \leq 2$ give the estimates in (37). \square

Remark 8. When $\beta = \alpha$ and $\psi = \varphi$, Theorem 7 reduces to [26, Theorem 2.3].

The following theorems give the estimates for the second and third coefficients of functions f when (i) $f \in \mathcal{M}(\alpha, \varphi)$ and $g \in \mathcal{L}(\beta, \psi)$ and (ii) $f \in \mathcal{L}(\alpha, \varphi)$ and $g \in \mathcal{L}(\beta, \psi)$. The proofs are similar as for the theorems above; hence, they are omitted here.

Theorem 9. Let $f \in \sigma$ and $g = f^{-1}$. If $f \in \mathcal{M}(\alpha, \varphi)$ and $g \in \mathcal{L}(\beta, \psi)$, then

$$\begin{aligned} |a_2| &\leq \left(B_1 D_1 \sqrt{2[B_1(3-2\beta) + D_1(1+2\alpha)]} \right) \\ &\quad \times \left(\left| \sigma B_1^2 D_1^2 - 2(1+\alpha)^2(3-2\beta)(B_2 - B_1) D_1^2 \right. \right. \\ &\quad \left. \left. - 2(2-\beta)^2(1+2\alpha)(D_2 - D_1) B_1^2 \right| \right)^{-1/2}, \\ |\sigma a_3| &\leq \frac{B_1}{2}(\beta^2 - 11\beta + 16) + D_1(1+3\alpha) \\ &\quad + \frac{1}{2}(\beta^2 - 11\beta + 16)|B_2 - B_1| \\ &\quad + \frac{(2-\beta)^2(1+3\alpha)B_1^2|D_2 - D_1|}{D_1^2(1+\alpha)^2}, \end{aligned} \tag{41}$$

where $\sigma := 10 + 14\alpha - 7\beta + \beta^2 + 2\alpha\beta^2 - 10\alpha\beta$.

Theorem 10. Let $f \in \sigma$ and $g = f^{-1}$. If $f \in \mathcal{L}(\alpha, \varphi)$ and $g \in \mathcal{L}(\beta, \psi)$, then

$$\begin{aligned} |a_2| &\leq \left(B_1 D_1 \sqrt{2[B_1(3-2\beta) + D_1(3-2\alpha)]} \right) \\ &\quad \times \left(\left| \sigma B_1^2 D_1^2 - 2(2-\alpha)^2(3-2\beta)(B_2 - B_1) D_1^2 \right. \right. \\ &\quad \left. \left. - 2(2-\beta)^2(3-2\alpha)(D_2 - D_1) B_1^2 \right| \right)^{-1/2}, \end{aligned} \tag{42}$$

$$\begin{aligned}
2|\sigma a_3| \leq & B_1(\beta^2 - 11\beta + 16) + D_1(8 - 5\alpha - \alpha^2) \\
& + (\beta^2 - 11\beta + 16)|B_2 - B_1| \\
& + \frac{(2 - \beta)^2(\alpha^2 + 5\alpha - 8)B_1^2|D_2 - D_1|}{D_1^2(2 - \alpha)^2},
\end{aligned} \tag{43}$$

where $\sigma := 24 + 3\alpha^2 + 3\beta^2 - 17\alpha - 17\beta - 2\beta\alpha^2 - 2\alpha\beta^2 + 12\alpha\beta$.

Remark 11. When $\beta = \alpha$ and $\psi = \varphi$, Theorem 10 reduces to [26, Theorem 2.4].

Conflict of Interests





The authors declare that there is no conflict of interests regarding the publication of this paper.

Acknowledgments

The research of the first and last authors is supported, respectively, by FRGS Grant and MyBrain MyPhD Programme of the Ministry of Higher Education, Malaysia.

References

- [1] P. L. Duren, *Univalent Functions*, vol. 259, Springer, New York, NY, USA, 1983.
- [2] M. Lewin, "On a coefficient problem for bi-univalent functions," *Proceedings of the American Mathematical Society*, vol. 18, pp. 63–68, 1967.
- [3] D. A. Brannan, J. Clunie, and W. E. Kirwan, "Coefficient estimates for a class of star-like functions," *Canadian Journal of Mathematics*, vol. 22, pp. 476–485, 1970.
- [4] E. Netanyahu, "The minimal distance of the image boundary from the origin and the second coefficient of a univalent function in $|z| < 1$," *Archive for Rational Mechanics and Analysis*, vol. 32, pp. 100–112, 1969.
- [5] D. A. Brannan and T. S. Taha, "On some classes of bi-univalent functions," *Universitatis Babeş-Bolyai. Studia. Mathematica*, vol. 31, no. 2, pp. 70–77, 1986.
- [6] R. M. Ali, S. K. Lee, V. Ravichandran, and S. Supramaniam, "The Fekete-Szegő coefficient functional for transforms of analytic functions," *Iranian Mathematical Society. Bulletin*, vol. 35, no. 2, article 276, pp. 119–142, 2009.
- [7] R. M. Ali, V. Ravichandran, and N. Seenivasagan, "Coefficient bounds for p -valent functions," *Applied Mathematics and Computation*, vol. 187, no. 1, pp. 35–46, 2007.
- [8] B. A. Frasin and M. K. Aouf, "New subclasses of bi-univalent functions," *Applied Mathematics Letters*, vol. 24, no. 9, pp. 1569–1573, 2011.
- [9] A. K. Mishra and P. Gochhayat, "Fekete-Szegő problem for a class defined by an integral operator," *Kodai Mathematical Journal*, vol. 33, no. 2, pp. 310–328, 2010.
- [10] T. N. Shanmugam, C. Ramachandran, and V. Ravichandran, "Fekete-Szegő problem for subclasses of starlike functions with respect to symmetric points," *Bulletin of the Korean Mathematical Society*, vol. 43, no. 3, pp. 589–598, 2006.
- [11] H. M. Srivastava, "Some inequalities and other results associated with certain subclasses of univalent and bi-univalent analytic functions," in *Nonlinear Analysis*, vol. 68 of *Springer Series on Optimization and Its Applications*, pp. 607–630, Springer, Berlin, Germany, 2012.
- [12] H. M. Srivastava, A. K. Mishra, and P. Gochhayat, "Certain subclasses of analytic and bi-univalent functions," *Applied Mathematics Letters*, vol. 23, no. 10, pp. 1188–1192, 2010.
- [13] Q.-H. Xu, H.-G. Xiao, and H. M. Srivastava, "A certain general subclass of analytic and bi-univalent functions and associated coefficient estimate problems," *Applied Mathematics and Computation*, vol. 218, no. 23, pp. 11461–11465, 2012.
- [14] Q.-H. Xu, Y.-C. Gui, and H. M. Srivastava, "Coefficient estimates for a certain subclass of analytic and bi-univalent functions," *Applied Mathematics Letters*, vol. 25, no. 6, pp. 990–994, 2012.
- [15] G. Murugusundaramoorthy, N. Magesh, and V. Prameela, "Coefficient bounds for certain subclasses of bi-univalent function," *Abstract and Applied Analysis*, vol. 2013, Article ID 573017, 3 pages, 2013.
- [16] H. Tang, G.-T. Deng, and S.-H. Li, "Coefficient estimates for new subclasses of Ma-Minda bi-univalent functions," *Journal of Inequalities and Applications*, vol. 2013, article 317, 2013.
- [17] S. G. Hamidi, S. A. Halim, and J. M. Jahangiri, "Coefficient estimates for bi-univalent strongly starlike and Bazilevic functions," *International Journal of Mathematics Research*, vol. 5, no. 1, pp. 87–96, 2013.
- [18] S. Bulut, "Coefficient estimates for initial Taylor-Maclaurin coefficients for a subclass of analytic and bi-univalent functions defined by Al-Oboudi differential operator," *The Scientific World Journal*, vol. 2013, Article ID 171039, 6 pages, 2013.
- [19] S. Bulut, "Coefficient estimates for a class of analytic and bi-univalent functions," *Novi Sad Journal of Mathematics*, vol. 43, no. 2, pp. 59–65, 2013.
- [20] N. Magesh, T. Rosy, and S. Varma, "Coefficient estimate problem for a new subclass of biunivalent functions," *Journal of Complex Analysis*, vol. 2013, Article ID 474231, 3 pages, 2013.
- [21] H. M. Srivastava, G. Murugusundaramoorthy, and N. Magesh, "On certain subclasses of bi-univalent functions associated with Hohlov operator," *Global Journal of Mathematical Analysis*, vol. 1, no. 2, pp. 67–73, 2013.
- [22] M. Çağlar, H. Orhan, and N. Yağmur, "Coefficient bounds for new subclasses of bi-univalent functions," *Filomat*, vol. 27, no. 7, pp. 1165–1171, 2013.
- [23] H. M. Srivastava, S. Bulut, M. C. Çağlar, and N. Yağmur, "Coefficient estimates for a general subclass of analytic and bi-univalent functions," *Filomat*, vol. 27, no. 5, pp. 831–842, 2013.
- [24] S. S. Kumar, V. Kumar, and V. Ravichandran, "Estimates for the initial coefficients of bi-univalent functions," *Tamsui Oxford Journal of Information and Mathematical Science*. In press.
- [25] W. C. Ma and D. Minda, "A unified treatment of some special classes of univalent functions," in *Proceedings of the Conference on Complex Analysis (Tianjin, 1992)*, Conference Proceedings and Lecture Notes in Analysis, pp. 157–169, International Press, Cambridge, Mass, USA.
- [26] R. M. Ali, S. K. Lee, V. Ravichandran, and S. Supramaniam, "Coefficient estimates for bi-univalent Ma-Minda starlike and convex functions," *Applied Mathematics Letters*, vol. 25, no. 3, pp. 344–351, 2012.
- [27] A. W. Kędzierzawski, "Some remarks on bi-univalent functions," *Annales Universitatis Mariae Curie-Skłodowska. Section A. Mathematica*, vol. 39, no. 1985, pp. 77–81, 1988.



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

